

On Linear Ill-Posed Problems

V. K. Ivanov

1962

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196201.84281>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

Reports of the Academy of Sciences of the USSR
1962. Volume 145, No. 2

MATHEMATICS

V. K. IVANOV

ON LINEAR ILL-POSED PROBLEMS

(Presented by Academician S. L. Sobolev on 26 I 1962)

1. Let X be a linear metric space; Y a Banach space; A a linear continuous operator from X into Y such that A^{-1} exists but is unbounded. Many linear ill-posed problems of mathematical physics and function theory can be reduced to solving the equation

$$Ax = y, \quad (1)$$

where $y \in Y$ is the given element and $x \in X$ the sought element. Because of the unboundedness of the operator A^{-1} , a solution does not exist for all y and is unstable: arbitrarily small variations of y may correspond to arbitrarily large changes in x .

As was shown by A. N. Tikhonov ⁽¹⁾, stability can be achieved if the solution is sought in a prescribed compact set $M \subset X$. Put $AM = N$. Then, according to a well-known topological theorem, the mapping $M = A^{-1}N$ is continuous on N . Therefore, if $y \in N$ and, as y varies, we do not leave N , then x will depend continuously on y . The estimate of stability is determined by the modulus of continuity of the mapping A^{-1} on N :

$$\omega(\delta) = \sup \rho(x, x') \quad \text{for } x, x' \in N, \|Ax - Ax'\| \leq \delta. \quad (2)$$

If M is symmetric with respect to the zero of the space X , then for sufficiently small δ in (2) one may take $x' = \theta$.

M. M. Lavrent'ev ⁽²⁾ constructed an approximate method for solving equation (1), in which it is assumed that instead of the exact value y its approximation y_δ is known with accuracy δ , and the function $\omega(\delta)$, or its majorant, is known. A similar device is used in the work of F. John ⁽³⁾.

Usually there are no effective criteria that make it possible to establish whether y belongs to N ; this has to be assumed known *a priori* (see ^(1,2)). At the same time, in approximate solution, instead of y one operates with its approximate

value y_δ , which may not lie in N , so that $A^{-1}y_\delta$ may not belong to M or may have no meaning.

In connection with this it is natural to change the formulation of the problem and, instead of the exact solution of equation (1), to seek a quasisolution (see below). For the quasisolution the classical well-posedness conditions are preserved (Theorem 1), and to find it one can indicate, by modifying known methods ^(2,5), convergent processes. If, for the given y , there exists a true solution in M , then the quasisolution coincides with it; in other cases it gives the best approximation to the solution.

2. Definition. We shall call a quasisolution of equation (1) on a given compact set M of the space X , and for a given $y_0 \in N$, a point $x_0 \in M$ for which $\|Ax - y_0\|$ attains its minimum on M .

In application to the Cauchy problem for the Laplace equation, the idea of best approximation is contained in the work of S. N. Mergelyan ⁽⁴⁾.

Theorem 1. A quasi-solution of equation (1) exists for any nonempty compact set $M \subset X$ and any $y \in Y$. If M is convex and the sphere in the space Y is strictly convex, then the quasi-solution is unique and depends continuously on y .

Proof. Existence follows from the compactness of M . If M is convex, then AM is also convex; $q = Ax$, where x is a quasi-solution, is the point N closest to y (the projection of y onto N).

Under the assumptions of the theorem, the projection of a point onto a convex set is uniquely determined; hence the uniqueness of the quasi-solution. Continuous dependence follows from uniqueness and compactness.

If, under the assumptions of Theorem 1, for a given y_0 equation (1) has in M a true solution x_0 , and a sequence $\{y_n\}$ of points of Y converges to y_0 , then the sequence of the corresponding quasi-solutions $\{x_n\}$ converges to x_0 . Therefore, if y_δ is an approximate value of y_0 , then for $y_\delta \rightarrow y_0$, for the quasi-solution x_δ we shall have $x_\delta \rightarrow x_0$, independently of whether y_δ belongs to N .

If Y is Hilbert and M is convex, then from geometric considerations it follows that if y and y' are elements of Y , and q and q' are their projections onto N , then $\|q - q'\| \leq \|y - y'\|$. Therefore, if in Hilbert Y , $\|y - y'\| \leq \delta$, and x and x' are the corresponding quasi-solutions, then $\rho(x, x') \leq \omega(\delta)$.

In ⁽⁵⁾ an approximate method for solving equation (1) is constructed under the assumption that $y \in N$. It can be shown that, in the case of linearity of A , for arbitrary y , the approximations constructed there converge to the quasi-solution. Thus ⁽⁵⁾ gives an approximate method for finding quasi-solutions.

3. More definite results can be obtained by specializing the spaces X and Y . We shall assume that X and Y are Hilbert spaces, $M = \Omega_R$ is the ball $\|x\| \leq R$, and A is a completely continuous operator from X into Y . Since the ball is weakly compact in a Hilbert space and this space is metrizable

with respect to the weak topology, the results of Section 2 for the weak topology in X are applicable to the case under consideration.

Let A^* be the operator adjoint to A . Then A^*A is a self-adjoint, positive, completely continuous operator from Y into Y . Denote by $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq \dots$ the complete system of its eigenvalues, and by $u_1, u_2, \dots, u_n, \dots$ the complete orthonormal system of its eigenvectors. Let

$$A^*y = \sum_n \beta_n u_n. \quad (3)$$

Theorem 2. The quasi-solution of equation (1) on Ω_R is expressed by the formula

$$x = \sum_n \frac{\beta_n}{\lambda_n + \lambda} u_n, \quad (4)$$

where $\lambda = 0$, if

$$\sum_n \frac{\beta_n^2}{\lambda_n^2} \leq R^2; \quad (5)$$

λ is the positive root of the equation

$$\sum_n \frac{\beta_n^2}{(\lambda_n + \lambda)^2} = R^2,$$

if

$$\sum_n \frac{\beta_n^2}{\lambda_n^2} > R^2. \quad (6)$$

Proof. The finding of quasi-solutions on Ω_R reduces to finding in Ω_R a vector x minimizing on Ω_R the quadratic functional $(Ax - y, Ax - y)$. If (5) holds, the minimum is unconditional, and its finding reduces to solving the equation

$$A^*Ax = A^*y. \quad (7)$$

The solution has the form (4) with $\lambda = 0$; it is the quasi-solution (the exact solution).

If (5) is not satisfied, then one must seek the minimum under the condition $(x, x) = R^2$.

Applying the method of Lagrange multipliers, we arrive at an equation of the second kind

$$A^*Ax + \lambda x = A^*y, \quad (8)$$

whose solution gives (4) with $\lambda > 0$.

When y is given approximately, owing to errors, as a rule, (6) will hold. The presence of positive λ in the denominator ensures strong convergence of the series (4).

The method of Theorem 2 is a development of the device of M. M. Lavrent'ev (see (2)); however, unlike the latter, for the application of this method there is no need to know the function $\omega(\delta)$ defined by (2). This makes it possible to restrict oneself to weak compactness of the set M .

Ural State University
named after A. M. Gorky

Received
22 I 1962

References

- ¹ A. N. Tikhonov, DAN, **39**, No. 35, 195 (1943).
- ² M. M. Lavrent'ev, DAN, **127**, No. 1, 31 (1959).
- ³ F. John, Ann. di Mat. pura ed appl., **40**, 129 (1955).
- ⁴ S. N. Mergelyan, UMN, **11**, No. 5, 3 (1956).
- ⁵ V. K. Ivanov, DAN, **142**, No. 5 (1962).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.