

Quasianalytic Classes of Functions in the Disk

B. I. Korenblum

1965

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196501.05360>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

B. I. Korenblyum

Quasianalytic Classes of Functions in the Disk

(Presented by Academician A. N. Kolmogorov, February 8, 1965)

Mathematics

1°. In the present paper, for functions analytic in the unit disk $|z| < 1$ and infinitely differentiable on the circle $|z| = 1$, a problem analogous to the classical Hadamard–Denjoy–Carleman problem of quasianalyticity is solved.

2°. Let $\mathfrak{D}$ be the class of functions $f(z)$ infinitely differentiable in the closed unit disk K ($|z| \leq 1$). This means that at each point $z_0 \in K$ and for each function $f(z) \in \mathfrak{D}$ there is an asymptotic expansion

$$f(z) \sim \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n \quad (x \rightarrow z_0, z \in K),$$

and, for $|z_0| < 1$, this series obviously has a positive radius of convergence. The boundary values $\tilde{f}(\theta) = f(e^{i\theta})$ ($f \in \mathfrak{D}$) form a class of periodic functions of the real variable θ , which we shall denote by $\tilde{\mathfrak{D}}$.

Let $\{A_n\}_0^\infty$ be a prescribed nondecreasing sequence of positive numbers. Introduce the following subclasses of $\mathfrak{D}$ and $\tilde{\mathfrak{D}}$:

1) $\mathfrak{D}\{A_n\}$ is the class of functions $f(z) \in \mathfrak{D}$ for which

$$\max_{z \in K} |f^{(n)}(z)| \leq CA_n \quad (n = 0, 1, 2, \dots), \quad (1)$$

where the constant C depends on $f(z)$.

2) $\tilde{\mathfrak{D}}\{A_n\}$ is the class of functions $\tilde{f}(\theta) \in \tilde{\mathfrak{D}}$ for which

$$\max_{-\infty < \theta < \infty} |\tilde{f}^{(n)}(\theta)| \leq CA_n \quad (n = 0, 1, 2, \dots). \quad (2)$$

Definition 1. The class $\mathfrak{D}\{A_n\}$ is called **quasianalytic** if from $f(z) \in \mathfrak{D}\{A_n\}$ and

$$f^{(n)}(z_0) = 0 \quad (n = 0, 1, 2, \dots), \quad (3)$$

where z_0 is some point of K , it follows that $f(z) \equiv 0$.

Obviously, (3) can hold for a function $f(z) \in \mathfrak{D}$ not identically equal to zero only when $|z_0| = 1$.

Definition 2. The class $\widetilde{\mathfrak{D}}\{A_n\}$ is called **quasianalytic** if from $\tilde{f}(\theta) \in \widetilde{\mathfrak{D}}\{A_n\}$ and

$$\tilde{f}^{(n)}(\theta_0) = 0 \quad (n = 0, 1, 2, \dots), \quad (4)$$

where θ_0 is some point $(-\infty < \theta_0 < \infty)$, it follows that $\tilde{f}(\theta) \equiv 0$.

Remark. The class $\widetilde{\mathfrak{D}}\{A_n\}$, generally speaking, does **not** coincide with the class of functions $f(e^{i\theta})$ that are boundary values of functions from $\mathfrak{D}\{A_n\}$.

Definition 3. We shall say that the sequence $\{A_n\}_0^\infty$ satisfies the Carleman-Ostrowski-Mandelbrojt conditions ($K - O - M$) if any one of the following equivalent (see (1), p. 29) conditions is fulfilled:

a) If we put $\beta_n = \inf_{k \geq n} A_k^{1/k}$, then

$$\sum_{n=1}^{\infty} \frac{1}{\beta_n} = \infty. \quad (5)$$

b) If we put $T(r) = \sup_{n \geq 1} r^n / A_n$, then

$$\int_0^\infty \frac{\log T(r)}{r^2} dr = \infty. \quad (6)$$

c) either $\lim_{n \rightarrow \infty} A_n^{1/n} < \infty$, or $\lim_{n \rightarrow \infty} A_n^{1/n} = \infty$ and

$$\sum_{n=1}^{\infty} \frac{A_n^c}{A_{n+1}^c} = \infty, \quad (7)$$

where $\{A_n^c\}$ is the convex regularization by means of logarithms (see (1), p. 24) of the sequence $\{A_n\}$.

3°. The following theorem gives the solution of the problem posed.

Theorem 1. *In order that the class $\mathfrak{D}\{A_n\}$ be quasianalytic, it is necessary and sufficient that the sequence $\{\sqrt{A_n}\}$ satisfy the $K - O - M$ conditions.*

The same conditions are necessary and sufficient for the quasianalyticity of the class $\widetilde{\mathfrak{D}}\{A_n\}$.

Remark 1. In the theory of quasianalytic functions one usually considers the broader classes

$$\mathbf{D}\{A_n\} = \bigcup_{k \geq 0} \mathfrak{D}\{k^n A_n\}, \quad \widetilde{\mathbf{D}}\{A_n\} = \bigcup_{k \geq 0} \widetilde{\mathfrak{D}}\{k^n A_n\}. \quad (8)$$

It can be shown that Theorem 1 is also valid for the classes $\mathbf{D}, \widetilde{\mathbf{D}}$.

Remark 2. The class $\mathfrak{D}\{A_n\}$ is a proper part of the class $\widetilde{\mathfrak{D}}\{A_n\}$ of all periodic (with period 2π) functions satisfying inequalities (2). From Carleman's classical result ⁽²⁾ it follows that condition (5) is necessary and sufficient for the quasianalyticity of $\widetilde{\mathfrak{D}}\{A_n\}$.

The condition of Theorem 1 is substantially broader. Thus, for example, the class $\mathfrak{D}\{(n!)^2\}$ is quasianalytic, although $\widetilde{\mathfrak{D}}\{(n!)^2\}$ does not possess this property.

4°. To prove Theorem 1 it is convenient to map the disk K conformally onto the half-plane $\operatorname{Re} z \geq 0$ in such a way that the point z_0 , at which (3) or (4) holds, goes to zero. Under this mapping the sequence $\{A_n\}$ becomes another one, but the fulfillment of the $K-O-M$ conditions for $\{\sqrt{A_n}\}$ is not disturbed. Moreover, instead of inequalities of type (1), (2) it is convenient to consider analogous inequalities for mean-square norms on the straight lines $\operatorname{Re} z = c \geq 0$. Thus we arrive at a theorem which is essentially equivalent to Theorem 1.

Theorem 2. *In order that there not exist a function $f(z) \not\equiv 0$, infinitely differentiable in the half-plane $\operatorname{Re} z \geq 0$ and such that*

$$\max_{0 \leq x < \infty} \int_{-\infty}^{\infty} |f^n(x + iy)|^2 dy \leq A_n^2 \quad (n = 0, 1, 2, \dots), \quad (9)$$

$$f^{(n)}(0) = 0 \quad (n = 0, 1, 2, \dots), \quad (10)$$

* It is known that the maximum in (9) is attained at $x = 0$.

necessary and sufficient that the sequence $\{\sqrt{A_n}\}$ satisfy the $K-O-M$ conditions.

Proof. Suppose such a function $f(z)$ exists. As is known ⁽³⁾, $f(z)$ can be represented in the form

$$f(z) = \frac{1}{\sqrt{2\pi}} \int_0^{\infty} g(t) e^{-tz} dt \quad (\operatorname{Re} z \geq 0),$$

where from (9) and (10) it follows that

$$\int_0^{\infty} t^{2n} |g(t)|^2 dt \leq A_n^2, \quad \int_0^{\infty} t^n g(t) dt = 0 \quad (n = 0, 1, 2, \dots). \quad (11)$$

We have arrived at the problem of uniqueness of the solution of a certain Stieltjes-type moment problem, but for measures from L^2 . We shall show that problem (11) reduces to the classical Watson problem.*

Consider the function

$$G(z) = \frac{1}{\sqrt{2\pi}} \int_0^\infty t^{z-\frac{1}{2}} g(t) dt = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty e^{(z+\frac{1}{2})\xi} g(e^\xi) d\xi \quad (\operatorname{Re} z > 0).$$

From (11) it follows that $G(z)$ has the following properties:

- a) $M(x) = \int_{-\infty}^\infty |G(x+iy)|^2 dy$ is bounded in every finite interval $0 < x < a$;
- b) $M(n) \leq A_n^2$ ($n = 0, 1, 2, \dots$);
- c) $G(\frac{1}{2} + n) = 0$ ($n = 0, 1, 2, \dots$).

Hence it is easy to conclude that the function $\Phi(z) = G(z) \sec \pi z$ is analytic in the half-plane $\operatorname{Re} z > 0$, and

$$M_1(x) = \int_{-\infty}^\infty \operatorname{ch}^2 \pi y |\Phi(x+iy)|^2 dy$$

is bounded in every finite interval $(0, a)$, and

$$M_1(n) \leq A_n^2 \quad (n = 0, 1, 2, \dots). \quad (12)$$

Finally, consider the Fourier transform of $\Phi(z)$

$$\varphi(\xi) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^\infty \Phi(iy) e^{-iy\xi} dy. \quad (13)$$

From (12) it follows that $\varphi(\xi)$ is analytic in the strip $|\operatorname{Im} \xi| < \pi$, and its boundary values on the lines $\operatorname{Im} \xi = \pm \pi$ are square-summable. Using the analyticity of $\Phi(z)$, we may shift the path of integration in (13):

$$\varphi(\xi) = \frac{1}{\sqrt{2\pi}} e^{-n\xi} \int_{-\infty}^\infty \Phi(n+iy) e^{-iy\xi} dy.$$

By virtue of (12),

$$\begin{aligned} \int_{-\infty}^\infty |\varphi(\xi + i\eta)|^2 e^{2n\xi} d\xi &= \int_{-\infty}^\infty |\Phi(n+iy)|^2 e^{2n\eta} dy \leq \\ &\leq 4 \int_{-\infty}^\infty |\Phi(n+iy)|^2 \operatorname{ch}^2 \pi y dy \leq 4A_n^2 \quad (-\pi < \eta < \pi; n = 0, 1, 2, \dots). \end{aligned}$$

* The connection between the uniqueness of the Stieltjes moment problem and the Watson problem is known: see, for example, ⁽⁴⁾.

Thus, for every function $f(z) \neq 0$, analytic in the half-plane $\operatorname{Re} z > 0$ and satisfying (9), (10), one can construct $\psi(\zeta) = 1/\varphi(\zeta) \neq 0$, analytic in the strip $|\operatorname{Im} \zeta| < \pi$ and satisfying the conditions

$$\int_{-\infty}^{\infty} |\psi(\xi + i\eta)|^2 e^{2n\xi} d\xi \leq A_n^2 \quad (-\pi < \eta < \pi; \ n = 0, 1, 2, \dots), \quad (14)$$

and conversely. Problem (14) is easily reduced to the classical Watson problem for the strip $|\eta| < \pi$, if one considers the function

$$\psi_\delta(\xi + i\eta) = \int_{\xi}^{\xi+\delta} \psi(\xi + i\eta) d\xi$$

($\delta > 0$), which, by virtue of (14), satisfies the inequalities

$$|\psi_\delta(\xi + i\eta)| \leq C A_n e^{-n\xi} \quad (|\eta| < \pi, \ \xi > 0; \ n = 0, 1, 2, \dots). \quad (15)$$

As is known ([1], p. 55), for the existence of a nonzero solution of problem (15) it is necessary and sufficient that the sequence $\{\sqrt[n]{A_n}\}$ not satisfy the C-O-M conditions. The same is necessary and sufficient for the existence of a nonzero solution of problem (14). The theorem is proved.

Kyiv Civil Engineering Institute

Received
8 II 1965

REFERENCES

1. S. Mandelbrojt, *Adjacent Series. Regularization of Sequences. Applications*, Moscow, 1955.
2. T. Carleman, *Les fonctions quasi analytiques*, Paris, 1926.
3. N. Wiener, R. Paley, *Fourier Transform in the Complex Domain*, Moscow, 1964.
4. P. Malliavin, C. R., 238, No. 26, 2481 (1954).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.