

ON SOME LIMITING PROPERTIES OF ORTHOGONAL POLYNOMIALS

Ya. L. Geronimus

1965

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196501.56058>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.512.7

MATHEMATICS

Ya. L. Geronimus

ON SOME LIMITING PROPERTIES OF ORTHOGONAL POLYNOMIALS

(Presented by Academician S. N. Bernstein on 30 III 1965)

1. Let the polynomials $\{\varphi_n(z)\}_0^\infty$ be orthonormal on the unit circle

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} \varphi_n(z) \overline{\varphi_m(z)} d\sigma(\theta) = \delta_{nm}, \quad z = e^{i\theta} \quad (m, n = 0, 1, 2, \dots), \quad (1)$$

where the set e of points of increase of the bounded nondecreasing function $\sigma(\theta)$ is closed; suppose that on the set $e_1 \subseteq e$ there exists $\sigma'(\theta) > 0$, and $e_2 \subset e_1$ is the set of Lebesgue points of the characteristic function of the set e_1 ; by $E_2 \subset E_1 \subset E$ we denote the corresponding sets on the unit circle and by d their transfinite diameters (capacities).

Theorem. If $d(E_2) = d(E) > 0$ and $\mu(\theta)$ is the Robin distribution function for the set E , then at every point $z_0 = e^{i\theta_0}$ at which $\mu'(\theta_0)$, $\sigma'(\theta_0) > 0$ exist, we have

$$\lim_{n \rightarrow \infty} \frac{K_n(z_0, z_0)}{n+1} = 2\pi \frac{\mu'(\theta_0)}{\sigma'(\theta_0)}, \quad K_n(z, z_0) = \sum_{s=0}^n \varphi_s(z) \overline{\varphi_s(z_0)} \quad (n = 0, 1, 2, \dots). \quad (2)$$

An analogous theorem holds for polynomials orthonormal on a finite interval of the real axis. These results are a generalization of earlier results (¹⁻⁷).

2. We outline the course of the proof of the theorem. Consider the polynomial

$$\Phi_{n+1}(z, z_0) = \frac{K_n(z, z_0)(z - z_0)}{\alpha_n^2 \Phi_n(z_0)}, \quad \Phi_n(z) = \frac{\varphi_n(z)}{\alpha_n} = z^n + \dots, \quad (n = 0, 1, 2, \dots); \quad (3)$$

all its roots $\{z_k^{(n)} = e^{i\theta_k^{(n)}}\}_0^n$ are simple and lie on $|z| = 1$. Let $\psi_n(\theta, \theta_0)$ be a stepwise nondecreasing function having a positive jump $1/(n+1)$ at each point

$\{\theta_k^{(n)}\}_0^n$; consider the infinite sequence of functions $\{\psi_n(\theta, \theta_0)\}_1^\infty$, and suppose that $\theta_0 = \{\theta_0^{(n)}\}_{n=1}^\infty$ is a point of increase of each of them.

We apply the theorem of P. P. Korovkin ⁽⁸⁾ on the existence of the limit

$$\lim_{n \rightarrow \infty} M_n^{1/n} = d(E), \quad M_n = \max |\Phi_n(z)|, \quad z \in E \quad (n = 1, 2, \dots); \quad (4)$$

then, slightly modifying Walsh's theorem ^(9, §§ 7.3, 7.4), we prove the existence of the limits

$$\lim_{n \rightarrow \infty} |\Phi_n(z)|^{1/n} = \lim_{n \rightarrow \infty} |\Phi_{n+1}(z, z_0)|^{1/(n+1)} = d(E)|\varphi(z)|, \quad |z| > R > 1, \quad (5)$$

where the function $w = \varphi(z)$ maps the region F complementing the set E in the extended plane onto the region $|w| > 1$.

Hence, using Helly's theorems, we derive convergence at all continuity points of $\mu(\theta)$:

$$\lim_{n \rightarrow \infty} \psi_n(\theta, \theta_0) = \mu(\theta). \quad (6)$$

Remark. If one does not use (8), then from our earlier results (5) there follows the validity of (4), and consequently also of (2), under the more restrictive condition: $\sigma'(\theta) > 0$ almost everywhere on the set $E^{(1)}$, where $E^{(1)}$ is the sum of a finite number of arcs of the unit circle, and $E = E^{(1)} + E^{(2)}$, where $E^{(2)}$ is an isolated countable set.

3. Let us now consider a stepwise nondecreasing function $\sigma_n(\theta, \theta_0)$, having $n+1$ points of increase at the same points $\{\theta_k^{(n)}\}_0^n$, and suppose the distributions $d\sigma_n(\theta, \theta_0)$ and $d\sigma(\theta)$ have equal moments $\{c_k\}_0^n$; it is easy to see that the quantity $2\pi\{K_n(z_0, z_0)\}^{-1}$ is equal to the mass of the distribution $d\sigma_n(\theta, \theta_0)$ concentrated at the point $\theta_0^{(n)} = \theta_0$. Let $\theta_0 = \{\theta_0^{(n)}\}_{n=1}^\infty$ be a point of increase of all the functions $\{\sigma_n(\theta, \theta_0)\}_1^\infty$; again using Helly's theorems and the determinacy of the trigonometric moment problem, we prove convergence in the main,

$$\lim_{n \rightarrow \infty} \sigma_n(\theta, \theta_0) = \sigma(\theta). \quad (7)$$

4. Let a small quantity $\varepsilon_n > 0$ be chosen in such a way that both functions $\sigma(\theta)$, $\mu(\theta)$ are continuous at the points $\theta_0 \pm \varepsilon_n$, and that between these points there lies only one point of increase θ_0 of the functions $\sigma_n(\theta, \theta_0)$ and $\psi_n(\theta, \theta_0)$; we have

$$\frac{K_n(z_0, z_0)}{n+1} = 2\pi \frac{\psi_n(\theta_0 + \varepsilon_n, \theta_0) - \psi_n(\theta_0 - \varepsilon_n, \theta_0)}{\sigma_n(\theta_0 + \varepsilon_n, \theta_0) - \sigma_n(\theta_0 - \varepsilon_n, \theta_0)}; \quad (8)$$

choosing n sufficiently large and using the convergence conditions (6) and (7), we obtain

$$\frac{K_n(z_0, z_0)}{n+1} = 2\pi \frac{\mu(\theta_0 + \varepsilon_n) - \mu(\theta_0 - \varepsilon_n)}{\sigma(\theta_0 + \varepsilon_n) - \sigma(\theta_0 - \varepsilon_n)} + o(1); \quad (9)$$

dividing the numerator and denominator of the right-hand side by $2\varepsilon_n$ and using the existence of the derivatives, we arrive at (2).

5. Let us consider several examples.

If E is the unit circle, then we have

$$\lim_{n \rightarrow \infty} \frac{K_n(z_0, z_0)}{n+1} = \frac{1}{\sigma'(\theta_0)}, \quad \sigma'(\theta_0) > 0 \quad (10)$$

under the condition $d(E_2) = 1$, or $\sigma'(\theta) > 0$ almost everywhere on the interval $[-\pi, \pi]$; this latter condition is satisfied, in particular, if $\ln \sigma'(\theta) \in L_1(-\pi, \pi)$; but it also holds for Polachek polynomials ⁽⁹⁾, 397–400), for which $\ln \sigma'(\theta) \notin L_1(-\pi, \pi)$.

Now let $E^{(1)}$ be the arc $[\exp \alpha, \exp(2\pi - \alpha)]$, and let the set $E^{(2)}$ be situated on the complementary arc; in this case we have

$$\lim_{n \rightarrow \infty} \frac{K_n(z_0, z_0)}{n+1} = \frac{\sin \theta_0/2}{\sigma'(\theta_c) \sqrt{\cos^{(2)} \alpha/2 - \cos^{(2)} \theta/2}}, \quad \sigma'(\theta_0) > 0, \quad \alpha < \theta_0 < 2\pi - \alpha,$$

under the condition $d(E_2) = \cos \alpha/2$, or $\sigma'(\theta) > 0$ almost everywhere on the interval $[\alpha, 2\pi - \alpha]$.

Kharkov Aviation Institute

Received
25 III 1965

CITED LITERATURE

- ¹ P. Erdős, P. Turan, *Ann. Math.*, **4**, 510 (1940).
- ² U. Grenander, M. Rosenblatt, *Trans. Am. Math. Soc.*, No. 1, 112 (1954).
- ³ G. Szegő, *Math. Zs.*, **12**, 61 (1922).
- ⁴ N. I. Akhiezer, *Zhurn. Inst. matem. AN USSR*, No. 3, 75 (1937).
- ⁵ Ya. L. Geronimus, *Matem. sborn.*, **23** (65), No. 1, 77 (1948).
- ⁶ Ya. L. Geronimus, *DAN*, **45**, No. 4 (1949).

⁷ Ya. L. Geronimus, *Polynomials Orthogonal on a Circle and an Interval*, 1958.

⁸ P. P. Korovkin, *Uch. zap. Kaliningradsk. ped. inst.*, issue 5, 34 (1958).

⁹ J. L. Walsh, *Interpolation and Approximation by Rational Functions in the Complex Domain*, 1961.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.