

ESTIMATES OF THE SPHERICAL MATRIX NORM AND OF THE CORRESPONDING LOGARITHMIC NORM

S. M. Lozinskii

1965

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196501.75547>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 512.831

MATHEMATICS

S. M. LOZINSKII

ESTIMATES OF THE SPHERICAL MATRIX NORM AND OF THE CORRESPONDING LOGARITHMIC NORM

(Presented by Academician V. I. Smirnov on 13 IV 1965)

1. **Notation.** Up to and including Sec. 5, all notation and definitions from (1) are adopted. The symbols Φ_I , Φ_{II} , Φ_{III} denote the matrix norms induced, respectively, by the 1st, 2nd, and 3rd vector norms ((2), p. 121), i.e., the norms obtained from (1), Sec. 6, for $p = +\infty$, $p = 1$, $p = 2$. Φ_{III} is often called the spherical matrix norm. The symbol γ_{III} denotes the logarithmic matrix norm corresponding to Φ_{III} . If $A \in \mathbf{M}$, then A^* denotes the matrix Hermitian-conjugate to A . If A is a square matrix, then

$$\tilde{A} \stackrel{\text{def}}{=} \frac{1}{2}(A + A^*).$$

If Σ denotes a simple or double sum, then Σ' denotes the sum obtained from Σ by replacing by zeros the terms with equal indices. For any $A \in \mathbf{M}$ we have (see (2), pp. 125-127)

$$\Phi_I(A) = \max_{\mu} \sum_{\nu} |a_{\mu\nu}|; \quad (1)$$

$$\Phi_{II}(A) = \max_{\nu} \sum_{\mu} |a_{\mu\nu}|; \quad (2)$$

$$\Phi_{III}(A) = \sqrt{\rho(AA^*)} = \sqrt{\rho(A^*A)}, \quad (3)$$

and for a square matrix A , in addition (see (3), pp. 59-60),

$$\gamma_{III}(A) = \sigma(\tilde{A}). \quad (4)$$

2. In (1), estimates were obtained for $\Phi_{III}(A)$ and $\gamma_{III}(A)$ in terms of the elements of the matrix A , using a decomposition of the matrix into blocks. Here we give estimates that do not use such a decomposition.

3. Let Φ be a real-valued function defined on the set of Hermitian matrices having only nonnegative eigenvalues, and such that on this set $\rho(A) \leq \Phi(A)$. Define on $\mathbf{M}$ the functions L_Φ and R_Φ by the formulas

$$L_\Phi(A) \stackrel{\text{def}}{=} \sqrt{\Phi(AA^*)}; \quad (5_1)$$

$$R_\Phi(A) \stackrel{\text{def}}{=} \sqrt{\Phi(A^*A)}. \quad (5_2)$$

4. **Theorem 1.** *Under the assumptions of Sec. 3, for $A \in \mathbf{M}$ we have*

$$\Phi_{\text{III}}(A) \leq L_\Phi(A); \quad (6_1)$$

$$\Phi_{\text{III}}(A) \leq R_\Phi(A). \quad (6_2)$$

Proof. $\rho(AA^*) \leq \Phi(AA^*)$. Applying (3), we obtain (6₁). Similarly for (6₂).

If Φ is expressed explicitly in terms of the elements of the matrix, then inequalities (6) give an estimate of $\Phi_{\text{III}}(A)$ expressed explicitly in terms of the elements of the matrix A .

For example, putting in (6) $\Phi = \Phi_I$, we obtain

$$\Phi_{\text{III}}(A) \leq \min \left(\sqrt{\Phi_I(AA^*)}, \sqrt{\Phi_I(A^*A)} \right) \quad \text{for } A \in \mathbf{M}. \quad (7)$$

Using the multiplicativity of Φ_I ((³), p.56), we obtain

$$\Phi_I(AA^*) \leq \Phi_I(A)\Phi_I(A^*) = \Phi_I(A)\Phi_{II}(A),$$

which together with (7) gives

$$\Phi_{\text{III}}(A) \leq \sqrt{\Phi_I(A)\Phi_{II}(A)} \quad \text{for } A \in \mathbf{M}. \quad (8)$$

5. Define on $\mathbf{M}$ the function N by the formula

$$N(A) \stackrel{\text{def}}{=} \left(\sum_{\mu=1}^m \sum_{\nu=1}^n |a_{\mu\nu}|^2 \right)^{1/2} \quad \text{for } A \in \mathbf{M}_{m \times n}; \quad (9)$$

N , as is known, is a matrix norm, and $\Phi_{\text{III}} \leq N$ on $\mathbf{M}$.

6. From this point on (with the exception of item 13), we consider only square matrices of fixed order n , and the term matrix norm is used in the sense of a norm on $\mathbf{M}_n$. The remaining symbols and terms are as above.
7. If A is a square matrix, then the symbol D_A (S_A) denotes the matrix obtained from A by replacing the off-diagonal (diagonal) elements by zeros.

8. Let Φ be a nonnegative numerical function on $\mathbf{M}_n$. Define on $\mathbf{M}_n$ the function $\bar{\Phi}$ by the formula

$$\bar{\Phi}(A) \stackrel{\text{def}}{=} \max(|a_{11}|, \dots, |a_{nn}|) + \Phi(S_A). \quad (10)$$

9. **Theorem 2.** Let Φ and $\bar{\Phi}$ be as in item 8. Then:

- 1) If Φ is a matrix norm, then $\bar{\Phi}$ is a matrix norm, $\bar{\Phi}(E_n) = 1$, and $\gamma_{\bar{\Phi}}(A) = \max_k \operatorname{Re} a_{kk} + \Phi(S_A)$.
- 2) If $\Phi_{\text{III}} \leq \Phi$ on $\mathbf{M}_n$, then $\Phi_{\text{III}} \leq \bar{\Phi}$ on $\mathbf{M}_n$.
- 3) If Φ is a matrix norm and $\Phi_{\text{III}} \leq \Phi$ on $\mathbf{M}_n$, then $\gamma_{\text{III}} \leq \gamma_{\bar{\Phi}}$ on $\mathbf{M}_n$.

Proof. 1) This is proved by straightforward computations.

- 2) Let $\Phi_{\text{III}} \leq \Phi$ on $\mathbf{M}_n$. Fix $A \in \mathbf{M}_n$. We have $A = D_A + S_A$, which gives

$$\begin{aligned} \Phi_{\text{III}}(A) &\leq \Phi_{\text{III}}(D_A) + \Phi_{\text{III}}(S_A) \leq \\ &\leq \max_k |a_{kk}| + \Phi(S_A) = \bar{\Phi}(A). \end{aligned}$$

- 3) This follows from 2) and the definition of the logarithmic norm.

Remark. The function

$$\bar{N}(A) = \max(|a_{11}|, \dots, |a_{nn}|) + N(S_A) \quad (11)$$

will be called the norm of E. Schmidt (see ⁽⁴⁾, p. 135). By Theorem 2(1) and (3),

$$\gamma_{\text{III}}(A) \leq \gamma_{\bar{N}}(A) = \max_k \operatorname{Re} a_{kk} + N(S_A). \quad (12)$$

10. Let m be a natural number, $2 \leq m \leq n$, and let $\varkappa$ denote a fixed set of pairwise distinct integers $k_1, \dots, k_m$, where $1 \leq k_1, \dots, k_m \leq n$. Put

$$T_{\varkappa}(A) \stackrel{\text{def}}{=} \sum_{\mu=1}^m \operatorname{Re} a_{k_{\mu} k_{\mu}}; \quad (13)$$

$$N_{\varkappa}(A) \stackrel{\text{def}}{=} \left(\sum_{\mu=1}^m |a_{k_{\mu} k_{\mu}}|^2 + \sum_{\mu, \nu=1}^n |a_{\mu \nu}|^2 \right)^{1/2}; \quad (14)$$

$$\Gamma_{\varkappa}(A) \stackrel{\text{def}}{=} \left([N_{\varkappa}(A)]^2 - \frac{[T_{\varkappa}(A)]^2}{m} \right)^{1/2}; \quad (15)$$

$$\alpha_{\varkappa}(A) \stackrel{\text{def}}{=} \frac{T_{\varkappa}(A)}{m} - \frac{1}{\sqrt{m(m-1)}} \Gamma_{\varkappa}(A); \quad (16)$$

$$Z_{\varkappa}(A) = \frac{T_{\varkappa}(A)}{m} + \sqrt{\frac{m-1}{m}} \Gamma_{\varkappa}(A). \quad (17)$$

If $m = n$, then the index $\varkappa$ on $T_{\varkappa}, N_{\varkappa}, \Gamma_{\varkappa}, Z_{\varkappa}$ will sometimes be omitted. Thus, for example, $T(A) = \text{Re tr } A$.

11. **Theorem 3.** Let $A \in M_n$. Then:

1a) If $m < n$ and

$$\alpha_{\varkappa}(A) \geq |a_{jj}|, \quad \text{for } j \notin \varkappa, \quad (18)$$

then $\Phi_{\text{III}}(A) \leq Z_{\varkappa}(A)$.

1b) If $m = n$ and $T(A) \geq N(A)$, then $\Phi_{\text{III}}(A) \leq Z(A)$.

2a) If $m < n$ and

$$\alpha_{\varkappa}(A) \geq \text{Re } a_{jj} \quad \text{for } j \notin \varkappa, \quad (19)$$

then $\gamma_{\text{III}}(A) \leq Z_{\varkappa}(A)$.

2b) $\gamma_{\text{III}}(A) \leq Z(A)$.

3) If $m < n$ and $\beta \geq |a_{jj}|$ for $j \notin \varkappa$, then

$$\Phi_{\text{III}}(A) \leq \begin{cases} \beta + ([N_{\varkappa}(A)]^2 - 2\beta T_{\varkappa}(A) + m\beta^2)^{1/2}, \\ \beta + \Gamma_{\varkappa}(A) \quad \text{provided } \beta \geq |T_{\varkappa}(A)|/m. \end{cases} \quad (20)$$

4) If $m < n$ and $\beta \geq \text{Re } a_{jj}$ for $j \notin \varkappa$, then the inequality is valid which is obtained from (20) by replacing Φ on the left by γ ; moreover, in the second line of formula (20) one may replace $|T_{\varkappa}(A)|$ by $T_{\varkappa}(A)$.

Proof. Let the matrix D be obtained from D_A by replacing, for $j \in \varkappa$, the element a_{jj} by $\alpha_{\varkappa}(A)$. Then, as is easy to calculate,

$$\alpha_{\varkappa}(A) + N(A - D) = Z_{\varkappa}(A).$$

Therefore, if $m < n$ and (18) holds, then

$$\Phi_{\text{III}}(A) \leq \Phi_{\text{III}}(D) + \Phi_{\text{III}}(A - D) \leq \alpha_{\varkappa}(A) + N(A - D) = Z_{\varkappa}(A), \quad (21)$$

which proves 1a). If $m = n$, then $T(A) \geq N(A)$ is equivalent to $\alpha_{\mathfrak{N}}(A) \geq 0$ and, consequently, the conditions of 1b) give (21); this proves 1b). If $m < n$ and (19) holds, then

$$\gamma_{\text{III}}(A) \leq \gamma_{\text{III}}(D) + \gamma_{\text{III}}(A - D) \leq \alpha_{\mathfrak{N}}(A) + N(A - D) = Z_{\mathfrak{N}}(A), \quad (22)$$

which gives 2a). If $m = n$, then $\gamma_{\text{III}}(D) = \alpha_{\mathfrak{N}}(A)$ and, consequently, (22) holds, which proves 2b).

Let $m < n$, let β be a real number, and let the matrices D_{β} and $\hat{D}$ be obtained from D_A by replacing, for $j \in \mathfrak{N}$, the elements a_{jj} respectively by β and $T_{\mathfrak{N}}(A)/m$. Then, if $\beta \geq |a_{jj}|$ for $j \notin \mathfrak{N}$, then $\Phi_{\text{III}}(D_{\beta}) \leq \beta$, and $\Phi_{\text{III}}(A) \leq \Phi_{\text{III}}(D_{\beta}) + \Phi_{\text{III}}(A - D_{\beta}) \leq \beta + N(A - D_{\beta})$, whence follows the upper of inequalities (20). If also $\beta \geq |T_{\mathfrak{N}}(A)|/m$, then $\Phi_{\text{III}}(\hat{D}) \leq \beta$ and

$$\Phi_{\text{III}}(A) \leq \Phi_{\text{III}}(\hat{D}) + \Phi_{\text{III}}(A - \hat{D}) \leq \beta + N(A - \hat{D}) = \beta + \Gamma_{\mathfrak{N}}(A).$$

Assertion 3) is proved. 4) is proved analogously.

12. Obviously, $N_{\mathfrak{N}}(\tilde{A}) \leq N_{\mathfrak{N}}(A)$, and the same is true for $Z_{\mathfrak{N}}, T_{\mathfrak{N}}$; moreover, $T_{\mathfrak{N}}(A) = T_{\mathfrak{N}}(\tilde{A})$ and $\alpha_{\mathfrak{N}}(A) \leq \alpha_{\mathfrak{N}}(\tilde{A})$. Therefore, when estimating $\gamma_{\text{III}}(A)$ by Theorem 3, it is recommended first to estimate $\gamma_{\text{III}}(\tilde{A})$ by this theorem, and then to use the equality $\gamma_{\text{III}}(A) = \gamma_{\text{III}}(\tilde{A})$.

13. **Theorem 4.** $\Phi_{\text{III}}(A) \leq \sqrt{Z(AA^*)}$ for $A \in M$.

Proof. $\rho(AA^*) = \sigma(AA^*) = \Upsilon_{\text{III}}(AA^*)$. But, by Theorem 3 (26), $\Upsilon_{\text{III}}(AA^*) \geq Z(AA^*)$.

14. Put

$$P_k \stackrel{\text{def}}{=} \sum_{\nu} |a_{k\nu}|, \quad Q_k \stackrel{\text{def}}{=} \sum_{\mu} |a_{\mu k}|.$$

Theorem 5. If $0 \leq \alpha \leq 1$, then

$$\sigma(A) \leq \max_k (\text{Re } a_{kk} + P_k^{\alpha} Q_k^{1-\alpha}). \quad (23)$$

Proof. The following inequality of A. Ostrovskii is known (see (5), p. 151):

$$\rho(A) \leq \max_k (|a_{kk}| + P_k^{\alpha} Q_k^{1-\alpha}). \quad (24)$$

Replacing A in (24) by $E_n + hA$ and applying (1), Theorem 1, we obtain (23).

15. **Example** ((5), p. 148). The eigenvalues of the matrix

$$A = \begin{bmatrix} 7 + 3i & -4 - 6i & -4 \\ -1 - 6i & 7 & -2 - 6i \\ 2 & 4 - 6i & 13 - 3i \end{bmatrix}$$

are 9 , $9 + 9i$, $9 - 9i$; $\rho(A) = 9\sqrt{2} \simeq 12.7$ (+), $\sigma(A) = 9$. In (5) several estimates for $\rho(A)$ and $\sigma(A)$ were obtained, the best of which are $\rho(A) < 22.05$ and $\sigma(A) < 22.05$. Theorem 4 and Theorem 3 (2a) give, respectively, $\Phi_{\text{III}}(A) \leq \sqrt[4]{Z(AA^*)} < 18.02$ and $\Upsilon_{\text{III}}(A) \leq Z(A) < 14.2$. Consequently, $\rho(A) < 18.02$, $\sigma(A) < 14.2$. Direct computation gives $\Phi_{\text{III}}(A) > 17.24$, $\Upsilon_{\text{III}}(A) = 13.5$.

Received
28 III 1965

REFERENCES

- ¹ S. M. Lozinskii, DAN, **163**, No. 4, 1965.
- ² D. K. Faddeev, V. N. Faddeeva, *Computational Methods of Linear Algebra*, Moscow-Leningrad, 1960.
- ³ S. M. Lozinskii, *Izv. vyssh. uchebn. zaved.*, Mathematics, No. 5 (1958); No. 5 (1959).
- ⁴ J. Naas, H. L. Schmid, *Mathematisches Wörterbuch*, 2, Berlin—Leipzig, 1961.
- ⁵ M. Marcus, H. Minc, *A Survey of Matrix Theory and Matrix Inequalities*, Boston, 1964.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.