

ON EXTENSIONS AND REMAINDERS OF FINITE ORDER OF COMPLETELY REGULAR SPACES

Kh. N. Inasaridze

1966

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196601.33886>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 513.83

MATHEMATICS

Kh. N. INASARIDZE

ON EXTENSIONS AND REMAINDERS OF FINITE ORDER OF COMPLETELY REGULAR SPACES

(Presented by Academician P. S. Aleksandrov on 28 V 1965)

In this note all topological spaces considered are completely regular Hausdorff spaces, and all mappings are continuous.

A sequence

$$(C^0(X), b_1 C^0(X), b_2 C^1(X), \dots, b_{n-1} C^{n-2}(X), b_n C^{n-1}(X))$$

of spaces, where $C^0(X) = X$; $b_i C^{i-1}(X)$ is some compact extension of $C^{i-1}(X)$ and $C^i(X) = b_i C^{i-1}(X) \setminus C^{i-1}(X)$ for $1 \leq i \leq n$, is called a **compact extension of n -th order** of the space X , and $C^n(X)$ an **n -th order remainder** of the space X . In the case where all b_i are Čech extensions β_i , the extension is called the **Čech extension of n -th order** of the space X , and the remainder of n -th order $\Gamma^n(X)$ the **Čech remainder of n -th order** of the space X .

Let M_n^* be the class of all compact extensions of n -th order of the space X . We introduce in M_n^* a quasi-ordering as follows:

$$(\tilde{C}^0(X), \tilde{b}_1 \tilde{C}^0(X), \tilde{b}_2 \tilde{C}^1(X), \dots, \tilde{b}_n \tilde{C}^{n-1}(X)) \geq (C^0(X), b_1 C^0(X), b_2 C^1(X), \dots, b_n C^{n-1}(X)),$$

if there exist mappings

$$f_i : \tilde{b}_i \tilde{C}^{i-1}(X) \rightarrow b_i C^{i-1}(X), \quad i = 1, \dots, n,$$

such that f_1 is an extension of the identity mapping $e : \tilde{C}^0(X) \rightarrow C^0(X)$, and for $i = 2, \dots, n$ the mapping f_i is an extension, on $\tilde{b}_i \tilde{C}^{i-1}(X)$, of the mapping induced by f_{i-1} . If all f_i are homeomorphisms, then we shall call such extensions **equivalent**.

Let M_n be the set of classes of equivalent extensions. Using Lemma 1.5 of the paper ⁽¹⁾, one can prove the following lemma.

Lemma 1. *The set M_n is a partially ordered set whose largest element is the class of the Čech extension of n -th order.*

It follows from Lemma 1 that if some n -th order remainder of the space X has a perfect property (see ⁽¹⁾), then all its n -th order remainders have this property.

Denote by P_n the following property of a space: the n -th order remainder of the space has the perfect property P . Then from Lemma 1.5 ⁽¹⁾ and from Lemma 1 the following proposition follows immediately.

Lemma 2. *The property P_n is a perfect property.*

Denote by $R^1(X)$ the set of all points of the space X that have no compact neighborhoods, and let, for $n > 1$,

$$R^n(X) = R^1[R^{n-1}(X)]$$

(see ⁽¹⁾). We shall assume that $X = R^0(X)$.

Let K_n , where $n \geq 0$, denote the class of all spaces whose n -th order remainder is compact. By a space with compact remainder of zeroth order we mean a compact space. It can be shown that there exists a space belonging to K_n and not belonging to K_{n-1} , and that if the space X has compact

if it is a remainder of finite order, then it contains an open everywhere dense locally compact subset A , and $X \setminus A = R^1(X)$.

Since compactness is an absolute property, Lemma 2 implies the following theorem.

Theorem 1. *If f is a perfect mapping of X onto Y , then X belongs to the class K_n if and only if Y belongs to the class K_n .*

Theorem 2. *A space X belongs to the class K_n if and only if $R^{n/2}(X)$ is compact, when n is even, and $R^{(n-1)/2}(X)$ is locally compact, when n is odd.*

This theorem follows from the fact that $R^j(X)$ is a remainder of order $2j$ of the space X for $j \geq 1$.

From Theorem 2, in particular, Theorem 3.1 of Henriksen and Isbell ⁽¹⁾ follows: X is locally compact at infinity (i.e., has a compact remainder of second order) if and only if $R^1(X)$ is compact. Further, X has a locally compact remainder of second order if and only if $R^1(X)$ is locally compact.

It is easy to see that if F is a closed subset of the space X , then $R^k(F)$ is a closed subset of $R^k(X)$. Hence it follows that if $X \in K_n$, then $F \in K_n$. Therefore, if X is a compact space and A is its subset belonging to K_n , then $X \setminus A$ belongs to K_{n+1} .

Theorem 3. Let A and B be subsets of a space X , with A compact and B a space in K_n . Then, for even n , $A \cup B \in K_n$, and for odd n , $A \cup B \in K_{n+1}$.

Proof. For $n = 0$ this is known. Let bX be a compact extension of X . We have

$$\overline{A \cup B} \setminus A \cup B = (\overline{B} \setminus B) \setminus A.$$

The closure is taken in X . Let $\overline{B} \setminus B = D$. It is clear that

$$[A \cup (\overline{D} \setminus D)] \cap (D \setminus A) = \emptyset \quad \text{and} \quad [A \cup (\overline{D} \setminus D)] \cup (D \setminus A) = \overline{D} \cup A.$$

Let us prove the theorem for $n = 1, 2$. If $B \in K_1$, then $D \in K_0$ and $\overline{D} \setminus D = \emptyset$. Then $D \setminus A \in K_1$, and therefore $A \cup B \in K_2$. If $B \in K_2$, then $D \in K_1$ and $\overline{D} \setminus D \in K_0$. Since $A \cup (\overline{D} \setminus D) \in K_0$, we obtain that $D \setminus A \in K_1$, and therefore $A \cup B \in K_2$. Suppose the theorem is proved for $n = m, m + 1$; we shall prove it for $n = m + 2, m + 3$. If $B \in K_{m+2}$, then $\overline{D} \setminus D \in K_m$. By induction $A \cup (\overline{D} \setminus D) \in K_{m+1}$. Then $D \setminus A \in K_{m+2}$, and therefore $A \cup B \in K_{m+3}$. If $B \in K_{m+3}$, then $\overline{D} \setminus D \in K_{m+1}$. By induction $A \cup (\overline{D} \setminus D) \in K_{m+2}$. Therefore we obtain that $D \setminus A \in K_{m+2}$ and, hence, $A \cup B \in K_{m+3}$. The theorem is proved.

Using Theorem 3, one can prove the following propositions:

1. The direct product of a compact space and a space in K_n is a space in K_n .
2. The direct product of a locally compact space and a space in K_n is a space in K_{n+1} for even n , and a space in K_n for odd n .
3. If X is a space in K_n and A is its compact subset, then $X \setminus A$ is a space in K_{n+1} for even n , and a space in K_n for odd n .
4. If X is a locally compact space and A is its subset belonging to the class K_n , then $X \setminus A$ is a space in K_{n+1} for even n , and a space in K_{n+2} for odd n .

From 4, in particular, it follows that the complement of a locally compact set in a locally compact space is a space for which the set of points at which the space is not locally compact is locally compact.

Let f be a mapping of X onto Y . Denote by $R(f)$ the set of all points $y \in Y$ such that $f^{-1}(y)$ is noncompact. By a theorem of Stone ⁽²⁾, Theorem 88, f has a unique extension $f^* : \beta X \rightarrow \beta Y$.

Theorem 4. If f is a closed mapping of X onto Y , then

$$f^*[\Gamma^1(X)] \setminus \Gamma^1(Y) = R(f).$$

Proof. It is clear that $R(f) \subset \bar{f}^*[\Gamma^1(X)] \setminus \Gamma^1(Y)$. Let $b \in \Gamma^1(X)$ and $f^*(b) = a' \in f^*[\Gamma^1(X)] \setminus \Gamma^1(Y)$. Suppose that $f^{-1}(a') = A$ is compact. Since A is closed

in βX and $b \notin A$, there exists a continuous function $\varphi : \beta X \rightarrow I = [0, 1]$ such that $\varphi(a) = 0$ for $a \in A$ and $\varphi(b) = 1$. Let

$$Q = \{x \in X, \varphi(x) \geq 1/2\}.$$

It is clear that Q is closed in X , and therefore $f(Q) = Q'$ is closed in Y , i.e. $Q' = P \cap Y$, where P is closed in βY . Obviously, $a' \notin Q'$. Then there exists a continuous function $\psi : \beta Y \rightarrow I$ such that $\psi(a') = 0$ and $\psi(p) = 1$ for $p \in P$. Let $k = \psi f^*$. Then $k(b) = 0$. Denote by D the set of points x of the space X for which $\varphi(x) > 1/2$ and $k(x) < 1/2$. It is clear that $b \in D$. The set D is open, and therefore there exists a point $s \in X$ such that $s \in D$, i.e. $k(s) < 1/2$ and $\varphi(s) > 1/2$. Further, we have $\psi[f^*(s)] < 1/2$, and therefore $f^*(s) = f(s) \notin Q'$. Hence it follows that $s \notin Q$, and therefore $\varphi(s) < 1/2$. The contradiction obtained shows that $f^{-1}(a')$ is noncompact, i.e. $a' \in R(f)$. The theorem is proved.

Using Theorems 3 and 4, one can prove the following theorem.

Theorem 5. *Let f be a closed mapping of X onto Y such that the $R(f)$ -space is of class K_n . Then, if $X \in K_1$, then $Y \in K_{n+2}$, and if $X \in K_2$, then for even n , $Y \in K_{n+2}$, while for odd n , $Y \in K_{n+3}$.*

In particular, it follows from Theorem 5 that under a closed mapping of a locally compact space, if the set of points whose full inverse image is noncompact is compact, then the image is a space for which the set of points having no compact neighborhoods is compact.

Let f be a mapping of X onto Y , and let $f^* : bX \rightarrow bY$ be an extension of f to their compact extensions bX and bY . Let $X_1 = bX \setminus X$, $Y_1 = f^*(X_1)$, and let f_1 be the mapping of X_1 onto Y_1 induced by the mapping f^* . Then the mapping $f_1 : X_1 \rightarrow Y_1$ will be called a **remainder of first order** (or simply a **remainder**) of the mapping f . If bX and bY are Čech compactifications, then f_1 is called the **Čech remainder** of the mapping f . The mapping $f_n : X_n \rightarrow Y_n$ is called a **remainder of n -th order of the mapping f** , if f_n is a remainder of first order of a remainder of $(n-1)$ -st order of the mapping f . It is easy to see that if some remainder of a closed mapping is a perfect mapping, then all remainders are also perfect. Such a mapping f will be called **perfect at infinity**.

With the aid of Lemma 1.5 ⁽¹⁾ and Theorem 4 one can prove the following theorem.

Theorem 6. *A closed mapping of X onto Y is perfect at infinity if and only if, for every point $x \in X$ having no compact neighborhoods, the full inverse image of its image is compact.*

Theorem 7. *A closed mapping of X onto Y with a closed Čech remainder is perfect at infinity if and only if the full inverse image of every point of Y is locally compact.*

With the help of Theorems 1, 3, and 4 we obtain the following theorem.

Theorem 8. *Let f be a closed mapping of X onto Y , perfect at infinity, such that $R(f)$ is compact. Then: 1) if $X \in K_n$, then $Y \in K_n$ for even n and $Y \in K_{n+1}$ for odd n ; 2) if $Y \in K_n$, then $X \in K_{n+1}$ for even n and $X \in K_n$ for odd n .*

Tbilisi Mathematical Institute named after A. M. Razmadze
Academy of Sciences of the Georgian SSR

Received
28 V 1965

References

1. Melvin Henriksen, J. R. Isbell, Duke Math. J., **25**, No. 1, 83 (1958).
2. M. H. Stone, Trans. Am. Math. Soc., **41**, 375 (1937).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.