

ON A CLASS OF SINGULAR INTEGRAL EQUATIONS

A. Yanushauskas

1966

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196601.38621>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.948

MATHEMATICS

A. YANUSHAUSKAS

ON A CLASS OF SINGULAR INTEGRAL EQUATIONS

(Presented by Academician M. A. Lavrent'ev on May 6, 1965)

Let R^3 be three-dimensional Euclidean space; $S : \{x^2 + y^2 + z^2 = 1\}$ the unit sphere of the space R^3 . Consider the following singular integral equation:

$$f_1(x, y, z) = [\alpha(z)(1 - z^2) + z\beta(z)]u_1(x, y, z) + \int_S \frac{\alpha(z)[x(x - x_0) + y(y - y_0)] + \beta(z)(z - z_0)}{[(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2]^{3/2}} u_1(x_0, y_0, z_0) ds, \quad (1)$$

where $P = (x, y, z)$ and $Q = (x_0, y_0, z_0)$ are points of the sphere S ; $\alpha(r)$, $\beta(r)$, and $f_1(x, y, z)$ are Hölder-continuous functions, $\alpha^2 + \beta^2 \neq 0$, $\beta(1) \neq 0$.

Using the equation of the sphere S and taking as independent variables on S the variables z and $\varphi = \arctg y/x$, we reduce equation (1) to the form

$$f(z, \varphi) = [\alpha(z)(1 - z^2) + z\beta(z)]u(z, \varphi) + \frac{\alpha(z)z - \beta(z)}{2\sqrt{2}} \int_{-1}^{+1} \int_0^{2\pi} \frac{(z - z_0)u(z_0, \psi)}{(1 - zz_0)^{3/2}[1 - \lambda \cos(\psi - \varphi)]^{3/2}} d\psi dz_0 + \frac{\alpha(z)}{\sqrt{2}} \int_{-1}^{+1} \int_0^{2\pi} \frac{u(z_0, \psi)}{(1 - zz_0)^{1/2}[1 - \lambda \cos(\psi - \varphi)]^{1/2}} d\psi dz_0, \quad (2)$$

where

$$\lambda = [(1 - z^2)(1 - z_0^2)]^{1/2} / (1 - zz_0).$$

The operator

$$T(u) = \frac{\alpha(z)}{\sqrt{2}} \int_{-1}^{+1} \int_0^{2\pi} \frac{u(z_0, \psi)}{(1 - zz_0)^{1/2}[1 - \lambda \cos(\psi - \varphi)]^{1/2}} d\psi dz_0$$

is completely continuous; therefore, first of all, we consider the equation

$$g(z, \varphi) = [\alpha(z)(1 - z^2) + z\beta(z)]u(z, \varphi) + \frac{\alpha(z)z - \beta(z)}{2\sqrt{2}} \int_{-1}^{+1} \int_0^{2\pi} \frac{(z - z_0)u(z_0, \psi)}{(1 - zz_0)^{3/2}[1 - \lambda \cos(\psi - \varphi)]^{3/2}} d\psi dz_0. \quad (3)$$

We shall seek solutions of equations (2) and (3) in the class of Hölder-continuous functions. Since $g(z, \varphi)$ and $u(z, \varphi)$ are Hölder-continuous on the sphere S , they can be represented by uniformly convergent series of the form ⁽¹⁾

$$u(z, \varphi) = \sum_{l=0}^{\infty} [u_l(z) \cos l\varphi + v_l(z) \sin l\varphi], \quad (4a)$$

$$g(z, \varphi) = \sum_{l=0}^{\infty} [f_l(z) \cos l\varphi + h_l(z) \sin l\varphi]. \quad (4b)$$

Substituting (4a) into (3), we obtain

$$[\alpha(z)(1 - z^2) + z\beta(z)]u(z, \varphi) + \frac{\alpha(z)z - \beta(z)}{2\sqrt{2}} \int_{-1}^{+1} \frac{(z - z_0)\sqrt{\pi}}{(1 - zz_0)^{3/2}(1 - z^2)} \times \sum_{l=0}^{\infty} \frac{\Gamma(l + \frac{3}{2})}{2^{l-1}\Gamma(l+1)} \lambda^l F\left(l/2 + \frac{1}{4}, \frac{l}{2} - \frac{1}{4}; l+1; \lambda^2\right) \times [u_l(z_0) \cos l\varphi + v_l(z_0) \sin l\varphi] dz_0 = g(z, \varphi). \quad (5)$$

Equation (5) can be rewritten in the following form:

$$g(z, \varphi) = [\alpha(z)(1 - z^2) + z\beta(z)]u(z, \varphi) + \frac{\alpha(z)z - \beta(z)}{2} \sum_{l=0}^{\infty} \int_{-1}^{+1} \frac{\lambda^l (1 - zz_0)^{1/2}}{z - z_0} [u_l(z_0) \cos l\varphi + v_l(z_0) \sin l\varphi] dz_0 + \int_{-1}^{+1} \int_0^{2\pi} H(z, z_0, \psi - \varphi) u(z_0, \psi) d\psi dz_0, \quad (6)$$

where

$$H(z, z_0, \psi - \varphi) =$$

$$= \frac{1}{z - z_0} \sum_{l=0}^{\infty} \lambda^l \left[\frac{\sqrt{\pi} \Gamma(l + \frac{3}{2})}{2^{l-1} \Gamma(l+1)} F\left(l/2 + \frac{1}{4}, l/2 - \frac{1}{4}; l+1; \lambda^2\right) - 1 \right] \cos l(\psi - \varphi).$$

The operator

$$A(u) = \int_{-1}^{+1} \int_0^{2\pi} H(z, z_0, \psi - \varphi) u(z_0, \psi) d\psi dz_0$$

is completely continuous.

Consider the equation

$$g(z, \varphi) = [\alpha(z)(1 - z^2) + z\beta(z)]u(z, \varphi) + \frac{\alpha(z)z - \beta(z)}{2} \sum_{l=0}^{\infty} \int_{-1}^{+1} \frac{\lambda^l (1 - zt)^{1/2}}{z - t} [u_l(t) \cos l\varphi + v_l(t) \sin l\varphi] dt. \quad (7)$$

Separating the variables in equation (7), we obtain

$$f_l(z) = [\alpha(z)(1 - z^2) + z\beta(z)]u_l(z) + \frac{\alpha(z)z - \beta(z)}{2} \int_{-1}^{+1} \frac{\lambda^l (1 - zt)^{1/2}}{z - t} u_l(t) dt \quad (8)$$

(an analogous equation is also obtained for $v_l(z)$). To equation (8) the theory developed in (2) is applicable.

Introduce the following notation:

$$Q(z) = -[\alpha(z)z - \beta(z)]\pi i/2, \quad P(z) = \alpha(z)(1 - z^2) + z\beta(z),$$

$$G(z) = \frac{P(z) - Q(z)(1 - z^2)^{1/2}}{P(z) + Q(z)(1 - z^2)^{1/2}},$$

$$P^*(z) = \frac{P(z)}{\{[P(z)]^2 - [Q(z)]^2(1 - z^2)\}},$$

$$Q^*(z) = \frac{Q(z)(1 - z^2)^{1/2}}{\{[P(z)]^2 - [Q(z)]^2(1 - z^2)\}}.$$

Choose that branch of the logarithm for which $\ln G(-1) = 0$; then we have $\ln G(+1) = 2n\pi i$, where $2n\pi$ is the increment of the argument of the function $G(z)$ on the interval $-1 \leq z \leq 1$, and n is an integer. Consider the integral operator

$$K^*(h) = P^*(z)h(z) - \frac{Q^*(z)Z(z)}{\pi i} \int_{-1}^{+1} \frac{h(t)}{Z(t)(t-z)} dt,$$

where $Z(z) = \omega(z)(z-1)^n$; $\omega(z)$ is a completely determined function depending only on P and Q ; $\omega(z) \neq 0$ for $-1 \leq z \leq 1$.

Let $n \leq 0$. From (8) we have

$$u_l(z) = K^*(f_l) + \mathcal{L}_l(u), \quad (9)$$

where

$$\mathcal{L}_l(u) = \frac{1}{\pi i} \int_{-1}^{+1} K^* \left(Q(r) \frac{\lambda^l (1-zt)^{1/2} - (1-z^2)^{1/2}}{z-t} \right) u(t) dt.$$

As $l \rightarrow \infty$, the operators $\mathcal{L}_l$ converge to the operator

$$A(u) = -K^* \left(\frac{Q(z)\sqrt{1-z^2}}{\pi i} \int_{-1}^{+1} \frac{u(t)}{t-z} dt \right) + Q^*(z)Q(z)(1-z^2)^{1/2}u(z).$$

The singular integral equation

$$f(z) = u(z) + A(u)$$

is always solvable and has a unique solution. It follows from this that equation (9), for sufficiently large l , is always solvable and has a unique solution. For the solution of equation (8) we have the representation ⁽²⁾

$$u_l(z) = P(z)f_l(z) + \int_{-1}^{+1} \frac{N_l(z,t)}{t-z} f_l(t) dt + \sum_{i=1}^m C_i \omega_i(z), \quad (10)$$

where $N_l(z, t)$ is a completely determined bounded function; C_i are arbitrary constants; ω_i are solutions of the homogeneous equation (8), and, for the validity of representation (10), it is necessary and sufficient that the function $f_l(z)$ satisfy $-n + m$ orthogonality conditions

$$\int_{-1}^{+1} f_l(z) \psi_{li}(z) dz = 0, \quad i = 0, 1, \dots, -n + m - 1, \quad (11)$$

where $\psi_{li}(z)$ are linearly independent solutions of the adjoint homogeneous equation.

From the convergence of the operators $\mathcal{L}_l$ to the operator A , and from the fact that the equation $f(z) = u(z) + A(u)$ is always solvable and has a unique solution, it follows that the functions $N_l(z, t)$ are bounded uniformly with respect to l . Consider the series

$$N(z, t, \psi - \varphi, \tau) = \frac{1}{z - t} \sum_{l=0}^{\infty} N_l(z, t) \tau^l \cos l(\psi - \varphi), \quad 0 \leq \tau < 1.$$

The solution of equation (7) is written in the following form:

$$u(z, \varphi) = P(z)g(z, \varphi) + \lim_{\tau \rightarrow 1} \int_{-1}^{+1} \int_0^{2\pi} N(z, t, \psi - \varphi, \tau) g(t, \psi) d\psi dt, \quad (12)$$

and, in order that a solution (12) exist, it is necessary and sufficient that the function $g(z, \varphi)$ satisfy a countable set of conditions

orthogonality

$$\int_{-1}^{+1} \int_0^{2\pi} \mu_{lj}(z) e^{il\varphi} g(z, \varphi) d\varphi dz = 0, \quad (13)$$

$$j = 0, 1, \dots, -n + m - 1; \quad l = 0, 1, 2, \dots$$

The operator

$$N(g) = \lim_{\tau \rightarrow 1} \int_{-1}^{+1} \int_0^{2\pi} N(z, t, \psi - \varphi, \tau) g(t, \psi) d\psi dt$$

is a bounded operator; therefore the following theorem is valid.

Theorem 1. If $n < 0$, then for the solvability of equation (3) it is necessary and sufficient that the function $g(z, \varphi)$ satisfy a countable set of orthogonality conditions. The homogeneous equation corresponding to equation (3) has no more than a finite number of linearly independent solutions. Equation (3) is normally solvable in the sense of Hausdorff.

The equation

$$h(z, \varphi) = [\alpha(z)(1 - z^2) + z\beta(z)] v(z, \varphi) - \\ - \frac{1}{2\sqrt{2}} \int_{-1}^{+1} \int_0^{2\pi} \frac{[t\alpha(t) - \beta(t)](z - t)}{(1 - tz)^{3/2}[1 - \lambda \cos(\psi - \varphi)]^{3/2}} v(t, \psi) d\psi dt \quad (14)$$

will be called the adjoint equation to equation (3). Equation (14) is also normally solvable in the sense of Hausdorff (3).

If $n > 0$ for equation (3), then for the adjoint equation (14) $n < 0$; therefore, also in the case $n > 0$, equation (3) is normally solvable in the sense of Hausdorff.

Theorem 2. Equation (3) is normally solvable in the sense of Hausdorff. For $n < 0$, for the solvability of equation (3) it is necessary and sufficient to impose on the function $g(z, \varphi)$ a countable set of orthogonality conditions; the corresponding homogeneous equation has no more than a finite number of linearly independent solutions. For $n > 0$, the homogeneous equation corresponding to equation (3) has infinitely many linearly independent solutions, and for the solvability of equation (3) it is necessary and sufficient to impose on the function $g(z, \varphi)$ no more than a finite number of orthogonality conditions. For $n = 0$, equation (3) is Fredholm.

Theorem 2 is also valid for the more general integral equation

$$h(z, \varphi) = [\alpha(z)(1 - z^2) + z\beta(z)] u(z, \varphi) + T(u) + \\ + \int_{-1}^{+1} \int_0^{2\pi} \frac{[z\alpha(z) - \beta(z)](z - t)}{2\sqrt{2}(1 - zt)^{3/2}[1 - \lambda \cos(\psi - \varphi)]^{3/2}} u(t, \psi) d\psi dt, \quad (15)$$

where the operator $T(u)$ is completely continuous. A special case of equation (15) is equation (2).

Institute of Mathematics
Siberian Branch of the Academy of Sciences of the USSR

Received
6 V 1965

REFERENCES

1. N. K. Bari, *Trigonometric Series*, Moscow, 1961.
2. N. I. Muskhelishvili, *Singular Integral Equations*, Moscow, 1962.
3. F. Hausdorff, *Set Theory*, Moscow-Leningrad, 1937.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.