

ON INVERSE THEOREMS OF UNIFORM APPROXIMATION

N. A. Lebedev

1966

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196601.42023>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.537

MATHEMATICS

N. A. LEBEDEV

ON INVERSE THEOREMS OF UNIFORM APPROXIMATION

(Presented by Academician V. I. Smirnov on February 12, 1966)

This note is devoted to a certain generalization of the inverse theorems of V. K. Dzyadyk ^(1,2).

Let $\bar{B}$ be a bounded set of points z of type $\mathfrak{M}_s^*$ (see ⁽³⁾, pp. 79-81) and let L be the boundary of $\bar{B}$. The complement of L in the extended plane consists of domains B^j , $j = 1, 2, \dots, s$, not belonging to $\bar{B}$, and of no more than a countable number of domains B^j , $j = s+1, s+2, \dots$, belonging to $\bar{B}$. Further we assume that every point z of the boundary L^j of the domain B^j is accessible from B^j , $j = 1, 2, \dots$.

Let a_j be a fixed point in B^j , $j = 1, 2, \dots, s$. We shall consider approximation of a function $f(z)$, given on L , by rational functions (r.f.) of the form

$$R(z) = \sum_{j=1}^s \sum_{\nu=1}^{n_j} \frac{A_{j,\nu}}{(z-a_j)^\nu} + A, \quad A_{j,n_j} \neq 0 \quad (1)$$

(if some $a_j = \infty$, then $1/(z-a_j)^\nu$ in (1) must be replaced by z^ν). The number $n = \max_{j=1,2,\dots,s} n_j$ will be called the degree of the r.f. (1).

Let: $g_j(z, \zeta)$ be the Green function of the domain B^j , $j = 1, 2, \dots, s$;

$L_\rho^j = \{z : g_j(z, a_j) = \ln \rho\}$, $\rho > 1$; $L_\rho = \bigcup_{j=1}^s L_\rho^j$; $d(z, u)$ be the distance from the point $z \in L$ to L_{1+u} , $u > 0$. Let $A(z)$, $z \in L$, be a continuous positive function. Introduce the function $A^*(z)$, putting $A^*(z) = A(z)$ for $z \in L$ and $A^*(z) = \exp G(z, B^j)$ for $z \in B^j$, $j = 1, 2, \dots$, where $G(z, B^j)$ is a function harmonic in B^j and continuous on $\bar{B}^j$ such that $G(z, B^j) = \ln A(z)$ for $z \in L^j$. The function $A^*(z)$ is continuous on the extended plane Z , and

$$\sup_{z \in Z} A^*(z) = \sup_{z \in L} A(z).$$

Put

$$A^*(z, y) = \sup_{|\zeta - z| = y} A^*(\zeta), \quad z \in L, \quad y > 0.$$

Lemma. Let the r.f. (1) of degree not exceeding n satisfy the condition $|R(z)| \leq A(z)$ for $z \in L$. Then:

- 1) $|R(z)| \leq A^*(z)$ for $z \in \bar{B}$ and $|R(z)| \leq A^*(z)\rho^n$ for $z \in L_\rho$, $\rho > 1$;
- 2) for any $u > 0$ and $z \in L$ we have

$$|R^{(\nu)}(z)| \leq \nu! e^u A^*(z, d(z, u/n)) d(z, u/n)^{-\nu}, \quad \nu = 1, 2, \dots;$$

- 3) for any $u > 0$, $z_1 \in L$, $z_2 \in L$, $|z_1 - z_2| < d(z_1, u/n)$, we have

$$|R(z_1) - R(z_2)| \leq \frac{2}{\pi} K(k) e^u A^*\left(z_1, d\left(z_1, \frac{u}{n}\right)\right) k, \quad k = \frac{|z_1 - z_2|}{d(z_1, u/n)},$$

where $K(k)$ is the complete elliptic integral of the first kind.

For the ideas of the proof see ⁽¹⁾, pp. 730–732 (additionally see ⁽³⁾, pp. 83, 97).

Let $\omega(x)$, $x > 0$, $\omega(0) = 0$, be a nondecreasing continuous function. Put $A(z) = A_u(z) = \omega(d(z, u))$, $z \in L$, $u > 0$. If there exists a number $M \leq 1$ such that $A^*(z, d(z, u)) = A_u^*(z, d(z, u)) \leq M A_u(z)$ for every $z \in L$ and every $u > 0$, then we write $\omega(x) \in \{\bar{B}, M\}$. If $\omega_1(x) \in$

$\{\bar{B}, M_1\}$; $\omega_2(x) \in \{\bar{B}, M_2\}$ and $\alpha > 0$ and $C > 0$ are constants, then $C\omega_1(x)^\alpha \in \{\bar{B}, M_1^\alpha\}$ and $\omega_1(x)\omega_2(x) \in \{\bar{B}, M_1 M_2\}$.

Theorem. Let $f(z)$ be a function defined on the boundary L of the set $\bar{B}$; let λ , $\lambda > 0$, be a fixed number, r a nonnegative integer; $x^r \omega(x) \in \{\bar{B}, M\}$ and $\omega(\lambda x) \leq \lambda' \omega(x)$ for all $x > 0$ and $\lambda' \geq 1$; suppose there exists a sequence of rational functions $R_n(z)$ of degree not exceeding n , $n = m, m+1, \dots$, of the form (1), such that

$$|f(z) - R_n(z)| \leq d(z, \lambda/n)^r \omega(d(z, \lambda/n)), \quad z \in L. \quad (2)$$

Then:

- 1) if $r = 0$ and $0 < x \leq e^{-1} \inf_{z \in L} d(z, \lambda/m) = x_0$, we have

$$\omega(x, f) \leq \omega(x, R_m) + C_0 M x \int_x^{l_m} \frac{\omega(t)}{t^2} dt, \quad (3)$$

where $l_m = \sup_{z \in L} d(z, \lambda/m)$ and C_0 is an absolute constant ($0 < C_0 < 4.2e^{\lambda_0}$, $\lambda_0 = \max\{(m+1)\lambda/m, \lambda+2\}$) ($\omega(x, \varphi)$ is the modulus of continuity of the function $\varphi(z)$ on L);

2) if $r > 0$; $\nu = 1, 2, \dots, r$,

$$|f^{(\nu)}(z) - R_n^{(\nu)}(z)| \leq C_r^{(\nu)} M \Omega_{r-\nu}(d(z, \lambda/n)), \quad z \in L, \quad n = m, m+1, \dots, \quad (4)$$

$$\Omega_p(x) = \int_0^x t^{p-1} \omega(t) dt, \quad 0 < C_r^{(\nu)} < \frac{4\nu! e^{\lambda_0+q}}{q(e^q - 1)}, \quad q = r - \nu + 1.$$

Remark 1. For $\nu = r$, from (4) we have

$$|f^{(r)}(z) - R_n^{(r)}(z)| \leq C_r^{(r)} M \Omega_0 \left(d \left(z, \frac{\lambda_*}{n+m} \right) \right), \quad z \in L, \quad n = m, m+1, \dots,$$

where $\lambda_* = \frac{m+r}{m} \lambda$. If $\Omega_0(x) \in \{\bar{B}, M_*\}$, then the first assertion of the theorem can be applied to $f^{(r)}(z)$.

Remark 2. In the first part of the theorem one can also give an estimate of $\omega(x, f)$ for $x > x_0$.

Proof (see additionally (2), pp. 371-374).

1) $r = 0$. Let $0 < x \leq x_0$, $z_1 \in L$, $z_2 \in L$, and $|z_1 - z_2| = x$. We have

$$\begin{aligned} f(z_2) - f(z_1) &= [f(z_2) - R_{n_1}(z_2)] - [f(z_1) - R_{n_1}(z_1)] \\ &+ \sum_{j=1}^{k-1} \{ [R_{n_j}(z_2) - R_{n_{j+1}}(z_2)] - [R_{n_j}(z_1) - R_{n_{j+1}}(z_1)] \} \\ &+ [R_{n_k}(z_2) - R_{n_k}(z_1)], \end{aligned} \quad (5)$$

where n_j , $m = n_k < n_{k-1} < \dots < n_1 < \infty$, are natural numbers.

We turn to the choice of n_j . Let n be the largest natural number such that

$$\max\{d(z_1, \lambda/n), d(z_2, \lambda/n)\} \geq ex,$$

and suppose, for example, that $d(z_1, \lambda/n) \geq d(z_2, \lambda/n)$. If $d(z_1, \lambda/n) \leq e^2 x$, then put $n_1 = n$. If $d(z_1, \lambda/n) > e^2 x$, then put $n_1 = n + 1$ and $n_2 = n$. If some $n_j > m$, $j \geq 1$, has been found, then we find the largest natural n such that $d(z_1, \lambda/n) \geq ed(z_1, \lambda/n_j)$, and if either $d(z_1, \lambda/n) \leq e^2 d(z_1, \lambda/n_j)$, or

$$n \geq \frac{m}{m+1} n_j,$$

then put $n_{j+1} = n$. In the contrary case put $n_{j+1} = n + 1$ and $n_{j+2} = n$. We continue this process until we obtain (or are forced to take) some $n_k = m$. For brevity denote

$$d_0 = x, \quad d_j = d(z_1, \lambda/n_j).$$

There are three possible cases: a) $ed_j \leq d_{j+1} \leq e^2 d_j$; b) $e^2 d_j < d_{j+1}$, but $n_j/n_{j+1} \leq (m+1)/m$; c) $d_{j+1} < ed_j$, but $d_{j+2} > e^2 d_j$ and $n_{j+1}/n_{j+2} \leq (m+1)/m$, or $j+1 = k$.

Let us estimate the right-hand side in (5). For $\nu = 1, 2$ we have $|f(z_\nu) - R_{n_1}(z_\nu)| \leq \omega(d(z_\nu, \lambda/n_1)) \leq \omega(d_1)$. If $d_1 < ex$, then $\omega(d_1) \leq \omega(ex)$, and if $ex \leq d_1 \leq e^2 x$, then $\omega(d_1) \leq \frac{d_1}{ex} \omega(ex) \leq e\omega(ex)$, and, consequently,

$$|[f(z_2) - R_{n_1}(z_2)] - [f(z_1) - R_{n_1}(z_1)]| \leq 2e\omega(ex).$$

$$|R_{n_j}(z_1) - R_{n_{j+1}}(z_1)| \leq |f(z_1) - R_{n_j}(z_1)| + |f(z_1) - R_{n_{j+1}}(z_1)| \leq 2\omega(d_{j+1}),$$

and, by virtue of the lemma (in our case $k \leq e^{-1}$ and $\frac{2}{\pi}K(k) < 1.05$), we have

$$\begin{aligned} I_j &= |[R_{n_j}(z_2) - R_{n_{j+1}}(z_2)] - [R_{n_j}(z_1) - R_{n_{j+1}}(z_1)]| \leq \\ &\leq 2.1 Me^{\lambda u} \omega(d_{j+1}) d(z_1, \lambda u/n_j)^{-1} x \end{aligned}$$

for $0 < u \leq n_j/n_{j+1}$. If $n_j/n_{j+1} \leq (m+1)/m$, then put $u = n_j/n_{j+1}$. In the contrary case put $u = 1$ and use the fact that $d_{j+1} \leq e^2 d_j$. In both cases we obtain the estimate $I_j \leq 2.1e^{\lambda_0} M \omega(d_{j+1}) d_{j+1}^{-1} x$.

From what has been stated and from (5) we have

$$|f(z_2) - f(z_1)| \leq 2e^2 \frac{\omega(ex)}{ex} x + 2.1e^{\lambda_0} M x \sum_{j=2}^k \frac{\omega(d_j)}{d_j} + \omega(x, R_m). \quad (6)$$

Let us estimate $\omega(d_j)/d_j$. If $d_j \geq ed_{j-1}$, then

$$\int_{d_{j-1}}^{d_j} \frac{\omega(t)}{t^2} dt \geq \frac{\omega(d_j)}{d_j} \int_{d_{j-1}}^{d_j} \frac{dt}{t} \geq \frac{\omega(d_j)}{d_j}.$$

If $d_j < ed_{j-1}$, then $d_{j-1} \geq ed_{j-2}$, and therefore

$$\int_{d_{j-2}}^{d_j} \frac{\omega(t)}{t^2} dt \geq \frac{\omega(d_j)}{d_j} \int_{d_{j-2}}^{d_{j-1}} \frac{dt}{t} \geq \frac{\omega(d_j)}{d_j}.$$

$$\int_x^{ex} \frac{\omega(t)}{t^2} dt \geq \frac{\omega(ex)}{ex}.$$

Now from (6) it follows that

$$|f(z_2) - f(z_1)| \leq \omega(x, R_m) + 4.2e^{\lambda_0} Mx \int_x^{d_k} \frac{\omega(t)}{t^2} dt,$$

and, since $d_k \leq l_m$, the case $r = 0$ has been considered.

2) $r > 0$. Choose a natural number $n_0 \geq m$ and represent $f^{(\nu)}(z)$ as

$$f^{(\nu)}(z) - R_{n_0}^{(\nu)}(z) = \sum_{j=1}^{\infty} [R_{n_j}^{(\nu)}(z) - R_{n_{j-1}}^{(\nu)}(z)], \quad z \in L, \quad (7)$$

where $m \leq n_0 < n_1 < n_2 < \dots$ are natural numbers chosen analogously to what was set out in the first part of the proof. Further, relying on the lemma, we estimate the right-hand side in (7) and obtain (4).

For the functions $\omega(x)$ and sets $\overline{B}$ considered by V. K. Dzyadyk, it was in fact proved by him that $\omega(x) \in \{\overline{B}, M\}$. It is not difficult to indicate broader classes of sets $\overline{B}$ for which $\omega(x) \in \{\overline{B}, M\}$ ($\omega(0) = 0$, $\omega(\lambda x) \leq \lambda \omega(x)$ for all $x > 0$ and $\lambda \geq 1$), but a complete analysis could not be carried out.

The foregoing can easily be somewhat generalized.

Received
10 II 1966

CITED LITERATURE

1. V. K. Dzyadyk, *Izv. AN SSSR, Ser. Mat.*, **23**, 697 (1959).
2. V. K. Dzyadyk, *Ukr. Mat. Zh.*, **15**, No. 4 (1963).
3. V. I. Smirnov, N. A. Lebedev, *Constructive Theory of Functions of a Complex Variable*, "Nauka," 1964.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.