

ON SPACES COABSOLUTE WITH METRIC SPACES

V. I. Ponomarev

1966

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196601.55653>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 513.831

MATHEMATICS

V. I. PONOMAREV

ON SPACES COABSOLUTE WITH METRIC SPACES

(Presented by Academician P. S. Aleksandrov on 14 V 1965)

Two topological spaces will be called **coabsolute** if their absolutes (see ^(14,8-10,13)) are homeomorphic to one another. As was proved in ^(14,8-10,13), the coabsoluteness of two spaces is necessary and sufficient for the existence of an irreducible, perfect (in general, many-valued) mapping of one of these spaces onto the other.

Definition 1. A system Σ of open sets of a space X is called **dense** in this space if every open $H \subseteq X$ contains some $U \in \Sigma$. The least cardinality of a dense system in X will be called the **π -weight of the space X** .

Main theorem. *In order that a space X admit a perfect irreducible single-valued mapping onto some metrizable space, it is necessary and sufficient that the space X be a paracompact p -space* and that in X there exist a dense system Σ of open sets decomposing into a countable sum**

$$\Sigma = \bigcup_{i=1}^{\infty} \Sigma_i$$

of disjoint systems Σ_i , each of which is locally finite in X . If a paracompact p -space X admits a many-valued perfect irreducible mapping onto some metrizable space, then there exists a metrizable space Y (in general, another one) and a single-valued perfect irreducible mapping $f : X \rightarrow Y$.

For the proof of this theorem we shall need some auxiliary notions and propositions, which are also of independent interest.

1. Product of mappings. Suppose mappings $f_\alpha : X \rightarrow Y_\alpha$ are given from one and the same space X onto spaces Y_α . The mapping

$$f : X \rightarrow Y \subseteq \prod_{\alpha} Y_{\alpha},$$

which assigns to each point $x \in X$ the point $\{f_\alpha x\}$ of the product $\prod_\alpha Y_\alpha$, is called the **product** of the mappings $f_\alpha : X \rightarrow Y_\alpha$. By P_f we shall denote the collection of all distinguished subsets ^{**} of the space X under the mapping $f : X \rightarrow Y$. The following assertions hold:

Lemma 1. *If $f : X \rightarrow Y$ is the product of mappings $f_\alpha : X \rightarrow Y_\alpha$, then $f^{-1}fx = \bigcap_\alpha f_\alpha^{-1}f_\alpha x$ for all $x \in X$ and $P_f \supseteq \bigcup_\alpha P_{f_\alpha}$.*

Lemma 2. *Let ω be some open covering of the space X ,*

* Paracompacts that admit a perfect mapping onto a metrizable space were studied in detail by A. V. Arhangel'skii ^(1,2) (see also ⁽¹¹⁾). We shall call these spaces paracompact p -spaces. In particular, in ^(1,2,11) intrinsic criteria are given for a space X to be a paracompact p -space. In the paper ⁽⁴⁾ it is proved (in our terminology) that a paracompact space complete in the sense of Čech is a paracompact p -space.

** A set $A \subseteq X$ is called distinguished under a mapping $f : X \rightarrow Y$ if $A = f^{-1}fA$.

and among the mappings $f_\alpha : X \rightarrow Y_\alpha$ there is at least one ω -mapping with respect to this ω . Then the product $f : X \rightarrow Y \subseteq \prod_\alpha Y_\alpha$ will necessarily be an ω -mapping with respect to the covering ω .

Lemma 3. Suppose that among the mappings $f_\alpha : X \rightarrow Y_\alpha$ there is at least one perfect mapping. Then the product $f : X \rightarrow Y$ of the mappings f_α is also perfect.

Lemma 4*. Let X be a paracompact p -space, and let ω be an arbitrary open covering of it. Then there exists a metrizable space Y_ω and a perfect ω -mapping $f_\omega : X \rightarrow Y_\omega$.

Lemma 5. Let X be a paracompact p -space, and let $U \subseteq X$ be an open subset of type F_σ . Then there exists a metrizable space Y_U and a perfect mapping $f_U : X \rightarrow Y_U$, under which the set U is distinguished, i.e. $U = f^{-1}fU$.

n. 2. Dense systems of open sets.

Lemma 6. Let $f : X \rightarrow Y$ be a one-to-one perfect irreducible mapping, and let $\Sigma = \{U\}$ be a system of open sets dense in X . Then the system ^{**} $f^\# \Sigma = \{f^\# U\}$ of open sets in Y is dense in Y .

Lemma 7. Let $f : X \rightarrow Y$ be a one-to-one perfect irreducible mapping, and let $\Sigma = \{V\}$ be a system of open sets dense in Y . Then the system $f^{-1}\Sigma = \{f^{-1}V\}$ is dense in X .

Lemma 8. Let $f : X \rightarrow Y$ be a many-valued (in particular, one-valued) perfect irreducible mapping (see (9)). Then the π -weight of the space X is equal to the π -weight of the space Y .

n. 3. Locally finite systems.

Lemma 9. Let $\Sigma = \{U_\alpha\}$ be a locally finite in X system of open sets of type F_σ . Then the open set

$$\tilde{\Sigma} = \bigcup_{\alpha} U_\alpha$$

—the body of the system Σ —also has type F_σ .

Lemma 10. Let Σ be a system of open sets dense in X and σ -locally finite*** (in all of X). Then for every open set $U \subseteq X$ there is an open set $U' \subseteq U$, everywhere dense in U , of type F_σ .

Lemma 11. If in the space X there is a system Σ of open sets that is dense and σ -locally finite (in all of X), then there is a dense σ -locally finite (also in all of X) system Σ' of open sets of type F_σ .

Lemma 12. Let $f : X \rightarrow Y$ be a one-to-one perfect irreducible mapping, and let $\Sigma = \{U\}$ be a locally finite in X system of open sets; then the system $f^\# \Sigma = \{f^\# U\}$ of open nonempty sets is locally finite in Y .

n. 4. Construction of mappings.

Basic Lemma 13. Let $\Sigma = \{U_\alpha\}$ be a disjoint locally finite in X system of open sets of type F_σ . Then there exists a metrizable space Y_Σ and a perfect mapping $f_\Sigma : X \rightarrow Y_\Sigma$, under which all sets $U_\alpha \in \Sigma$ are distinguished.

Proof. By Lemma 9 the open set $\tilde{\Sigma}$ has type F_σ , and the closed set $\Phi = X \setminus \tilde{\Sigma}$ has type G_δ . Let

$$\Phi = \bigcap_{i=1}^{\infty} \Gamma_i.$$

By Ω_i denote the covering of the whole space X , consisting of all

* The proof of this lemma is contained in (3).

** In order that a one-to-one mapping $f : X \rightarrow Y$ be simultaneously closed and irreducible, it is necessary and sufficient that for every open $U \subseteq X$ the set

$$f^\# U = \mathcal{E}(y \in Y, f^{-1}y \subseteq U)$$

be nonempty and open in Y (proved in (9, 12)).

*** A system Σ is called σ -locally finite in X if

$$\Sigma = \bigcup_{i=1}^{\infty} \Sigma_i,$$

where the Σ_i are locally finite in X . The systems Σ_i need not be coverings of the whole space X .

sets $U_\alpha \in \Sigma$ and also the sets Γ_i . By Lemma 4 there exists a metrizable space Y_i and a perfect Ω_i -map $f_i : X \rightarrow Y_i$.

Consider the product $f : X \rightarrow Y \subseteq \prod_{i=1}^{\infty} Y_i$ of the mappings $f_i : X \rightarrow Y_i$. By Lemma 2 the mapping f is an Ω_i -map for every i , and by Lemma 3 it is perfect. It remains now to prove that every $U_\alpha \in \Sigma$ is distinguished for the mapping f . Let $x_0 \in U_\alpha$ be arbitrary. We must prove that $f^{-1}fx_0 \subseteq U_\alpha$. Choose i_0 such that $x_0 \in \Gamma_{i_0}$. Since f is an Ω_{i_0} -map, there exists $U \in \Omega_{i_0}$ such that $f^{-1}fx_0 \subseteq U$, and this U is necessarily one of the sets $U_\alpha \in \Sigma$ (here the disjointness of the system Σ is essential). We have $f^{-1}fx_0 \subseteq U_\alpha$, as was required to prove.

Main Lemma 14. *Let in a paracompact p -space X there be a system Σ of open sets of type F_σ , decomposing into a countable sum*

$$\Sigma = \bigcup_{i=1}^{\infty} \Sigma_i$$

of disjoint systems Σ_i , each of which is locally finite in X . Then there exists a metrizable space Y_Σ and a perfect mapping $f_\Sigma : X \rightarrow Y_\Sigma$ under which all sets $U \in \Sigma$ are distinguished.

Proof. By Lemma 13, for each Σ_i there exists a metrizable space Y_i and a perfect mapping $f_i : X \rightarrow Y_i$, under which every set $U \in \Sigma_i$ is distinguished. Consider the product $f : X \rightarrow Y$ of the mappings $f_i : X \rightarrow Y_i$. The space Y is metrizable, and the mapping f is perfect by Lemma 3. Further, by Lemma 1, all sets $U \in \Sigma$ are distinguished. The lemma is proved.

5. Proof of the Main Theorem. a) Let X be a paracompact p -space, and let $\Sigma = \{U_\alpha\}$ be a dense system in X of open sets decomposing into a countable sum

$$\Sigma = \bigcup_{i=1}^{\infty} \Sigma_i$$

of disjoint systems Σ_i locally finite in X . By Lemma 10, in X there exists a dense system $\Sigma' = \{U'_\alpha\}$ of open sets U'_α of type F_σ , with each U'_α everywhere dense in U_α and $U'_\alpha \subseteq U_\alpha$. Denote by Σ'_i for each $\Sigma_i = \{U_\lambda^i\} \subseteq \Sigma$ the collection of all $U_\lambda'^i$. We obtain

$$\Sigma' = \bigcup_{i=1}^{\infty} \Sigma'_i,$$

and each system Σ'_i is disjoint and locally finite in X . By Lemma 14, consider a metrizable space Y and a perfect mapping $f : X \rightarrow Y$, under which all $U_\alpha'^i \in \Sigma'$ are distinguished. We shall prove that the mapping f is irreducible. Indeed, it is enough to prove that in every open set $\Gamma \subseteq X$ there is contained the complete preimage $f^{-1}y$ of some point $y \in Y$. But the system Σ' is dense in X . Therefore there exists $U'_\alpha \in \Sigma'$ such that $U'_\alpha \subseteq \Gamma$, and the set U'_α is distinguished for the mapping f .

- b) Let the space X admit a one-to-one perfect mapping $f : X \rightarrow Y$ onto some metrizable space Y . By Bing's metrization criterion (7), in Y there exists a σ -discrete base Σ ,

$$\Sigma = \bigcup_{i=1}^{\infty} \Sigma_i,$$

where the Σ_i are discrete in Y . Every base of a space is necessarily a dense system. Therefore, by Lemma 7, the system $f^{-1}\Sigma$ is dense in X ; moreover, the systems $f^{-1}\Sigma_i$ are disjoint and locally finite in X , and all $U \in f^{-1}\Sigma$ even have type F_σ .

- c) Let the paracompact p -space X admit a many-to-one perfect irreducible mapping $f : X \rightarrow Y$ onto some metrizable space Y . Then there exist a space Z and one-to-one perfect irreducible mappings $p_X : Z \rightarrow X$ and $p_Y : Z \rightarrow Y$ such that

$$f = p_Y^\# p_X^{-1}$$

(see (8°)). Consider in Y a σ -discrete base $\Sigma = \{V\}$. Then, by Lemmas 6 and 7, the system $p_X^\# p_Y^{-1}\Sigma$ will be a dense system of open sets of the space X ; moreover, if

$$\Sigma = \bigcup_{i=1}^{\infty} \Sigma_i,$$

Σ_i are discrete, then $p_X^\# p_Y^{-1}\Sigma = \bigcup_{i=1}^{\infty} p_X^\# p_Y^{-1}\Sigma_i$, and the $p_X^\# p_Y^{-1}\Sigma_i$ are disjoint and locally finite in X (by Lemma 12). Thus, in the space X there exists a dense system $p_X^\# p_Y^{-1}\Sigma$, decomposing into a countable sum of disjoint systems locally finite in X . And then, as has already been proved, there exists a metrizable space Y' and a one-to-one perfect irreducible mapping $g : X \rightarrow Y'$. The theorem is completely proved.

§ 6. Consequences.

Theorem 2. Let X be a bicom pactum. Then, in order that X admit a one-to-one irreducible mapping onto some compactum, it is necessary and sufficient that the π -weight of this bicom pactum be countable. If a bicom pactum X admits a many-valued irreducible mapping onto some compactum, then it also admits a one-to-one irreducible mapping onto some (generally speaking, different) compactum.

Theorem 3. In order that a completely normal bicom pactum X admit an irreducible mapping onto a compactum, it is necessary and sufficient that it contain an everywhere dense countable set.

Theorem 4. In order that a bicom pactum be coabsolute with the Cantor perfect set, it is necessary and sufficient that it have no isolated points and have countable π -weight.

Theorem 5. In order that a space X admit a one-to-one perfect irreducible mapping onto some metric space with complete metric, it is necessary and sufficient that X be a Čech-complete paracompact space and have a dense system Σ of open sets decomposing into a countable sum of disjoint and locally finite systems.

If a space X admits a many-valued perfect irreducible mapping onto some metrizable space with complete metric, then X is a Čech-complete paracompact space and there exists a one-to-one perfect irreducible mapping of it onto some metrizable space with complete metric.

R. Engelking and A. Pełczyński proved ⁽⁶⁾ that a dyadic bicom pactum admitting a one-to-one irreducible mapping onto some compactum is metrizable. We shall supplement this result with the following propositions, which follow easily from our main theorem:

Theorem 6. If a dyadic bicom pactum is coabsolute with some compactum, then it is necessarily metrizable.

Theorem 7. Let X be a dyadic bicom pactum, and let X_0 be its everywhere dense subset coabsolute with some metric space Y_0 . Then Y_0 and X (and, of course, $X_0 \subseteq X$) have a countable base.

From this there immediately follows a theorem of B. Efimov ⁽⁵⁾: all dyadic bicom pact extensions of a space with a countable base are compacta.

Moscow State University
named after M. V. Lomonosov

Received
21 I 1965

CITED LITERATURE

- ¹ A. V. Arhangel'skii, DAN, **151**, No. 4, 751 (1963).
- ² A. V. Arhangel'skii, Matem. sborn., **67** (109), No. 1, 55 (1965).
- ³ V. L. Klyushin, DAN, **159**, No. 4, 734 (1964).
- ⁴ Z. Frolik, Bull. Acad. polon. sci., sér. math., **8**, No. 11–12, 743 (1960).
- ⁵ B. Efimov, DAN, **151**, No. 5 (1963).
- ⁶ R. Engelking, A. Pełczyński, Colloq. Math., **11**, No. 1, 55 (1963).
- ⁷ R. H. Bing, Canad. J. Math., **3**, 175 (1951).
- ⁸ V. I. Ponomarev, DAN, **143**, No. 1 (1962).
- ⁹ V. I. Ponomarev, Matem. sborn., **60** (102), 89 (1963).
- ¹⁰ V. I. Ponomarev, DAN, **149**, No. 1, 25 (1963).
- ¹¹ V. I. Ponomarev, DAN, **153**, No. 5, 1013 (1963).
- ¹² V. I. Ponomarev, DAN, **132**, No. 6, 1269 (1960).
- ¹³ C. Iliadis, DAN, **149**, No. 1, 22 (1963).
- ¹⁴ A. M. Gleason, Illinois J. Math., **2**, 4A, 482 (1958).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.