

METRIC PROPERTIES OF HARMONIC FUNCTIONS

M. L. Gerver

1966

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196601.68501>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.947.42

MATHEMATICS

M. L. GERVER

METRIC PROPERTIES OF HARMONIC FUNCTIONS

(Presented by Academician I. G. Petrovskii, September 30, 1965)

In this note we shall formulate two theorems on harmonic functions of two variables. An imprecise, but brief, formulation of Theorem 1 is: *harmonic functions have few long level lines.*

The exact formulation and the outline of the proof are given in § 1. In Theorem 2 a bounded harmonic function that is small on a set of sufficiently large measure is estimated. This theorem is a generalization of Hadamard's theorem on three circles. Both theorems admit a generalization to the case of elliptic equations with variable coefficients. Theorem 1 is readily carried over to the multidimensional case. Theorem 2 (both in the plane and in space) can also be obtained by a completely different method as a consequence of the results of M. Lavrent'ev⁽⁴⁾.

Throughout what follows, K_R denotes the disk $x^2 + y^2 < R^2$.

§ 1. On the length of a level line. Let $u(x, y)$ be a bounded function harmonic in K_R . The set of points $(x, y) \in K_{R/2}$ at which $u(x, y) = t$ will be denoted by E_t . For any t , the set E_t is the union of a finite or countable number of pairwise nonintersecting analytic simple arcs. We denote the sum of the lengths of these arcs by $l(t)$. Let n be a natural number. The set of those t for which $l(t) > 16Rn^2$ will be denoted by $B_{u,n}$. Let U_M be the set of all functions u harmonic in K_R and bounded in absolute value by the constant M .

Theorem 1. *For every $M > 0$ there exists an N such that, for any $n > N$ and any function $u \in U_M$, the measure of $B_{u,n}$ is less than 2^{-n} .*

The example of the function $r^n \sin n\varphi$ in the disk K_1 shows that this estimate cannot be greatly improved. The method of proof of Theorem 1 is close to that used in ⁽¹⁾.

We outline the proof. Let $u \in U_M$. Fix $t \in B_{u,n}$ so that E_t contains no zeros of $\text{grad } u$. Let $(x, y) \in E_t$. If the tangent to E_t at the point (x, y) forms with the x -axis an angle not exceeding $\pi/4$, assign the point (x, y) to the set X . The set $E_t \setminus X$ will be denoted by Y .

From X we select a subset X_1 by means of the following condition. A point $(x, y) \in X$ is assigned to X_1 if there exists a segment of length $R/2n$, parallel to the y -axis, containing the point (x, y) and intersecting E_t in at least n points. Similarly, if some segment of length $R/2n$, parallel to the x -axis and passing through a point $(x, y) \in Y$, intersects E_t in at least n points, then we assign (x, y) to the set Y_1 .

The proof of Theorem 1 is carried out in three stages:

1. It is established that at the points of the set $X_1 \cup Y_1$, for $n > N(M)$, the inequality

$$|\operatorname{grad} u| < \frac{2^{-n}}{R}$$

holds.

2. It is verified that the linear measure of $X_1 \cup Y_1$ is greater than $l(t)/2$.
3. From 1 and 2 it follows that

$$\operatorname{mes} B_{u,n} < 2^{-n}.$$

§ 2. A generalization of the three-circles theorem. One of the possible formulations of the theorem on three circles for harmonic functions is as follows*:

* This theorem can easily be obtained from the well-known Hadamard theorem on three circles for analytic functions of a complex variable ⁽²⁾.

If $u(x, y)$ is a function harmonic in K_1 , and if $|u| < 1$ in K_1 , $|u| < \varepsilon^2$ in K_{r^2} , then $|u| < C\varepsilon$ in K_r (ε and r are arbitrary numbers between 0 and 1/2, C is an absolute constant).

How does this estimate change if, in the formulation of the theorem, the circle K_{r^2} is replaced by an arbitrary set $\mathfrak{M}$ of the same measure ($\operatorname{mes}_2 \mathfrak{M} = \pi r^4$)?

The answer to this question is given by

Theorem 2. There exists an $s > 0$ such that for every set $\mathfrak{M}$ of measure πr^4 , situated in K_r , from the inequalities

$$|u| < 1 \text{ in } K_1, \quad |u| < \varepsilon^s \text{ on } \mathfrak{M}$$

it follows that

$$|u| < \varepsilon \text{ in } K_r.$$

(As above, $\Delta u = 0$ in K_1 , $0 < \varepsilon, r < 1/2$.)

Let us note that it is enough to prove Theorem 2 for $r = r_0$, in order to be sure of its validity for $r = r_0^n$ (where n is any natural number). The case $r = r_0^n$ is reduced to the case $r = r_0$ if one maps the circle K_1 onto itself by means of the transformation $\sqrt[n]{z}$ ($z = x + iy$).

Consequently, it is enough to establish that the following is true.

Theorem 2'. There exists an $s > 0$ such that for every set $\mathfrak{M}$ of measure $\pi/16$, situated in $K_{1/2}$, for every function $u(x, y)$ harmonic in K_1 , and for every sufficiently large natural number N , from the inequalities

$$|u| < 1 \text{ in } K_1, \quad |u| < 2^{-sN} \text{ on } \mathfrak{M}$$

it follows that

$$|u| < 2^{-N} \text{ in } K_{1/2}.$$

We outline the plan of the proof of Theorem 2'.

1. We shall call the following construction the k -division of the circle K_R . Consider k concentric circumferences with center 0 and radii $R/k, 2R/k, \dots, iR/k, \dots, R$. Denote by $H_2, \dots, H_k$ the concentric rings of width R/k enclosed between these circumferences. We divide the ring H_i ($1 < i \leq k$) into $2i - 1$ equal regions $D_{i1}, \dots, D_{i,2i-1}$; to D_{ij} belong those points (r, φ) of the ring H_i for which

$$\frac{2\pi}{2i-1}(j-1) < \varphi < \frac{2\pi}{2i-1}j, \quad 1 \leq j \leq 2i-1.$$

All the regions D_{ij} , obviously, are of equal area: $\text{mes}_2 D_{ij} = \pi R^2/k^2$. We shall call $H_2, \dots, H_k$ the rings of the k -division, and $D_{21}, \dots, D_{k,2k-1}$ the cells of the k -division.

2. By the hypothesis of the theorem, $|u| < 2^{-sN}$ on $\mathfrak{M}$. Denote by $\mathfrak{M}_1$ the set of points of $K_{1/2}$ where $|u| < 2^{1-sN}$. With the help of Theorem 1 it is not difficult to prove that there exists an absolute constant k_0 with the following property:

Let $k \geq k_0(sN)^2$. Carry out the k -division of the circle $K_{1/2}$. Then at least $k^2/8$ cells of the division belong entirely to $\mathfrak{M}_1$.

3. Now, using the theorem on three circles, one can reduce Theorem 2' to the following lemma:

Lemma 1. Let q and n be natural numbers. Carry out the 2^{q+4} -division of the circle $K_{1/2}$. Let $u(x, y)$ be a harmonic function in the circle K_1 , satisfying the conditions: 1) $|u| < 1$ in K_1 ; 2) among the rings of the division situated between the circumferences of radii $1/32$ and $1/2$, there are $2q$ or more rings containing cells of the division where $|u| < 2^{-2q+n}$ (at least one such cell in each); 3) $\max_{K_{1/2}} |u| > 2^{-2q}$.

Then $n < A$, where A is an absolute constant.

The proof of Lemma 1 recalls the proof of Lemma 1.6.1 in (3).

Moscow State University
named after M. V. Lomonosov

Received
29 IX 1965

REFERENCES

1. E. M. Landis, DAN, 79, No. 3, 393 (1951).
2. A. I. Markushevich, *Theory of analytic functions*, M.—L., 1950.
3. E. M. Landis, UMN, 14, issue 1 (85), 21 (1959).
4. M. M. Lavrent'ev, On certain incorrect problems of mathematical physics, Novosibirsk, 1962.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.