

ONE PROPERTY OF A SOLUTION OF THE EQUATION

$$\backslash(u_{xx}u_{yy}-u_{xy}^2)=1\backslash)$$

B. D. Annin

1966

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196601.79941>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.944

MATHEMATICS

B. D. ANNIN

ONE PROPERTY OF A SOLUTION OF THE EQUATION $u_{xx}u_{yy} - u_{xy}^2 = 1$

(Presented by Academician A. D. Aleksandrov on 6 IX 1965)

In the present paper, for a solution of the equation $u_{xx}u_{yy} - u_{xy}^2 = 1$, an expression is given for the first-order derivatives on the boundary of a convex domain in terms of a single function of one variable.

Consider the Monge—Ampère equation of elliptic type

$$u_{xx}u_{yy} - u_{xy}^2 = 1 \quad (1)$$

in a finite simply connected domain G , bounded by a convex curve Γ , given in the form

$$x = \varphi(s), \quad y = \psi(s), \quad 0 \leq s \leq s^*. \quad (2)$$

Here s is arc length, s^* is the length of the contour Γ , and as s increases the contour Γ is traversed in the positive direction (the domain G remains on the left).

Let $f(s)$ be a function defined for $0 \leq s \leq s^*$. We shall assume that: a) the functions $\varphi(s), \psi(s), f(s)$ belong to the class $C^{k,\lambda}$, $k \geq 4$, $0 < \lambda < 1$; b) at $s = 0$ and $s = s^*$ the values of the functions $\varphi(s), \psi(s), f(s)$ coincide together with their derivatives up to order k inclusive; c) the curvature at all points of Γ is not less than $\chi_0 = \text{const} > 0$.

There exist ^(1,2) two and only two solutions of equation (1) of class $C^{k,\mu}(G + \Gamma)$, $0 < \mu \leq \lambda$, which take on Γ the value $f(s)$

$$u|_{\Gamma} = f(s). \quad (3)$$

For one of these solutions the functions u_{xx} and u_{yy} are positive in $G + \Gamma$, for the other they are negative.

Consider the solution for which u_{xx}, u_{yy} are positive. Let $(x_1, y_1), (x_2, y_2)$ be two distinct points belonging to $G + \Gamma$; then the inequality ⁽³⁾

$$(x_2 - x_1)[u_x(x_2, y_2) - u_x(x_1, y_1)] + (y_2 - y_1)[u_y(x_2, y_2) - u_y(x_1, y_1)] > 0 \quad (4)$$

is valid.

Introduce new variables ξ, η and a new function Φ by the formulas

$$\xi = u_x + x, \quad \eta = u_y + y, \quad (x, y) \in G + \Gamma; \quad (5)$$

$$\Phi = xu_x + yu_y - u + (x^2 + y^2)/2. \quad (6)$$

From (4) follows the inequality

$$(x_2 - x_1)^2 + (y_2 - y_1)^2 < (\xi_2 - \xi_1)^2 + (\eta_2 - \eta_1)^2. \quad (7)$$

Here the points $(\xi_1, \eta_1), (\xi_2, \eta_2)$ of the ξ, η -plane correspond to the points $(x_1, y_1), (x_2, y_2) \in G + \Gamma$ by virtue of formulas (5).

It follows from inequality (7) that formulas (5) effect a homeomorphic mapping of $G + \Gamma$ onto some domain $D + L$ of the ξ, η -plane. The equation of L —the image of Γ —can be written in the form $(0 \leq s \leq s^*)$

$$\xi_L(s) = u_x(\varphi(s), \psi(s)) + \varphi(s), \quad \eta_L(s) = u_y(\varphi(s), \psi(s)) + \psi(s). \quad (8)$$

* The derivative of order k satisfies the Hölder condition with exponent λ .

In view of inequality (4), the first derivatives with respect to s of the functions $\xi_L(s)$ and $\eta_L(s)$ cannot vanish simultaneously. The Jacobian of the mapping (5), in view of (1), is positive in $G + L$; therefore, as s increases the curve L is traversed in the positive direction.

Let l denote the length of an arc of the contour L , and let l^* be the length of L ; in so doing, as the origin of measurement we choose the point corresponding to $s = 0$, and we shall assume that, as l increases, the curve L is traversed in the positive direction. From the equation $l = l(s)$, $0 \leq s \leq s^*$, one can determine the function of class $C^{k-1, \mu}$, $s = s(l)$, $0 \leq l \leq l^*$.

The function $\Phi(\xi, \eta)$ belongs to the class $C^{k-1, \mu}(D + L)$ and satisfies in D the equation*

$$\Phi_{\xi\xi} + \Phi_{\eta\eta} = 1.$$

At the same time, in $D + L$ the equalities

$$x = \Phi_{\xi}, \quad y = \Phi_{\eta}, \quad u = \xi\Phi_{\xi} + \eta\Phi_{\eta} - \frac{1}{2}(\Phi_{\xi}^2 - \Phi_{\eta}^2). \quad (9)$$

hold. Thus, on L the equalities (8) and the equalities

$$\partial\Phi/\partial\xi|_L = \varphi(s), \quad \partial\Phi/\partial\eta|_L = \psi(s). \quad (10)$$

are satisfied.

Let the function $w_1(\zeta) = \xi + i\eta$ map conformally the unit disk $K + C : |\zeta| = 1$ of the plane of the complex variable ζ onto the domain $D + L$, and let τ be the polar angle in the ζ -plane. We shall assume that the point determined by the equalities $|\zeta| = 1$, $\tau = 0$ corresponds to the point of the contour L taken as the origin for measuring the arc length l , and that the point $\zeta = 0$ corresponds to the point (ξ_0, η_0) , which under the transformation (5) is the image of some point $(x_0, y_0) \in G$. The function (*) $w_1(\zeta) \in C^{k-1, \mu}(K + C)$, and the function $l = l(\tau)$, $0 \leq \tau \leq 2\pi$, belongs to the class $C^{k-1, \mu}$.

Denote

$$s = s(l(\tau)) \equiv g(\tau), \quad 0 \leq \tau \leq 2\pi. \quad (11)$$

The function $g(\tau)$ realizes a homeomorphic mapping of $[0, 2\pi]$ onto $[0, s^*]$ and belongs to the class $C^{k-1, \mu}$.

Consider the function (*) of ζ of class $C^{k-1, \mu}(K + C)$

$$(-2\Phi_{\eta} + \eta) + i(-2\Phi_{\xi} + \xi) = w_2(\zeta).$$

In view of equalities (10) and (11), for $0 \leq \tau \leq 2\pi$ we have

$$\begin{aligned} \operatorname{Re} \left[\frac{1}{2}w_1(\zeta) + \frac{i}{2}w_2(\zeta) \right]_{\zeta=e^{i\tau}} &= \varphi(g(\tau)), \\ \operatorname{Re} \left[-\frac{i}{2}w_1(\zeta) - \frac{1}{2}w_2(\zeta) \right]_{\zeta=e^{i\tau}} &= \psi(g(\tau)). \end{aligned}$$

Hence, using the Schwarz formula, we find (*)

$$w_1(\zeta) = \frac{1}{2\pi} \int_{-\pi}^{\pi} [\varphi(g(\omega)) + i\psi(g(\omega))] \frac{e^{i\omega} + \zeta}{e^{i\omega} - \zeta} d\omega + a, \quad (12)$$

$$w_2(\zeta) = -\frac{1}{2\pi} \int_{-\pi}^{\pi} [i\varphi(g(\omega)) + \psi(g(\omega))] \frac{e^{i\omega} + \zeta}{e^{i\omega} - \zeta} d\omega + b.$$

Here $a = u_x(x_0, y_0) + iu_y(x_0, y_0)$, $b = u_y(x_0, y_0) + iu_x(x_0, y_0)$. Setting $\zeta = e^{it}$ in the first formula (12) and taking account of formulas (11) and (8), we obtain

* The indices ξ, η denote partial derivatives.

representation of the boundary derivatives of first order of the function $u(x, y)$ in terms of $g(\tau)$

$$\left. \frac{\partial u}{\partial x} \right|_{\Gamma} = -\frac{1}{2\pi} \int_{-\pi}^{\pi} \psi(g(\omega)) \operatorname{ctg} \frac{\tau - \omega}{2} d\omega + u_x(x_0, y_0), \quad (13)$$

$$0 \leq \tau \leq 2\pi$$

$$\left. \frac{\partial u}{\partial y} \right|_{\Gamma} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \varphi(g(\omega)) \operatorname{ctg} \frac{\tau - \omega}{2} d\omega + u_y(x_0, y_0).$$

Remark 1. As is seen from formulas (3), (11), (13), the function $g(\tau)$ satisfies the singular integral equation ($0 \leq \tau \leq 2\pi$)

$$f'(g(\tau)) - \varphi'(g(\tau))u_x(x_0, y_0) - \psi'(g(\tau))u_y(x_0, y_0) =$$

$$= \frac{1}{2\pi} \int_{-\pi}^{\pi} [\varphi(g(\omega))\psi'(g(\tau)) - \varphi'(g(\tau))\psi(g(\omega))] \operatorname{ctg} \frac{\tau - \omega}{2} d\omega;$$

$$f'(s) = df(s)/ds; \quad \varphi'(s) = d\varphi(s)/ds, \quad \psi'(s) = d\psi(s)/ds.$$

Remark 2. Formulas (9) and (12) give a parametric representation of the solution of the Dirichlet problem for equation (1).

Institute of Hydrodynamics
Siberian Branch of the Academy of Sciences of the USSR

Received
1 IX 1965

CITED LITERATURE

1. S. N. Bernstein, *Collected Works*, 3, Publishing House of the Academy of Sciences of the USSR, 1960.
2. I. Ya. Bakel' man, DAN, 157, No. 2, 247 (1964).
3. P. Hartman, A. Wintner, *Am. J. Math.*, 75, No. 3 (1953).
4. M. A. Lavrent' ev, B. V. Shabat, *Methods of the Theory of Functions of a Complex Variable*, Moscow, 1958.
5. L. A. Galin, *Prikl. matem. i mekh.*, 13, issue 3, 285 (1949).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.