

ON MEASURES IN LINEAR SPACES

G. E. Shilov

1966

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196601.88627>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.397

MATHEMATICS

G. E. SHILOV

ON MEASURES IN LINEAR SPACES

(Presented by Academician A. N. Kolmogorov on 26 X 1965)

Let X be a real linear space and let X' be some linear system of linear functionals on X . A natural number n , a collection of n functionals $f_1, \dots, f_n$ from X' , and a Borel set A in the real n -dimensional space R^n determine in X the cylindrical set

$$C = C(f_1, \dots, f_n, A) = \{x \in X : [(f_1, x), \dots, (f_n, x)] \in A\} \quad (1)$$

with base A and generators $f_1, \dots, f_n$.

We shall say that a **premeasure** is given on the totality $\mathfrak{A}$ of all cylindrical sets if to each set $C \in \mathfrak{A}$ there is assigned a nonnegative number μC in such a way that: a) μC depends only on the set C itself and not on its representation in the form (1); b) on the totality $\mathfrak{A}_f$ of all cylindrical sets $C \in \mathfrak{A}$ with fixed collection $f = (f_1, \dots, f_n)$, the function μC is countably additive (countable additivity on all of $\mathfrak{A}$ is not assumed); c) $\mu X = 1$.

We pose the question under what conditions a premeasure μ can be extended to a measure—a countably additive set function defined on a σ -ring of subsets $E \subset X$ containing all cylindrical sets.

If X is the space of **all** real functions $x = x(t)$ on some set T , and the functionals $f \in X'$ are generated by the values of x at individual points $t \in T$, then a positive answer is the content of a theorem of A. N. Kolmogorov ⁽¹⁾. There are a number of conditions due to various authors that clarify this question under certain assumptions concerning the existence and properties of a topology on X ^(2,3). The following simple result pertains to spaces without topology. Suppose that X' is a total system of functionals on X , i.e., from the equalities $(f, x) = 0$, satisfied for a fixed $x \in X$ for all $f \in X'$, it follows that $x = 0$. Then X can be regarded as a space of real functions $x(f) = (f, x)$, where f ranges over X' or even only over some total subset T in X' . Let $\hat{X}$ be the space of **all** real functions $x = x(f)$, $f \in T$, with the premeasure of cylindrical sets (with functionals f , equal to the values of the functions x at the points $f \in T$) given by the formula

$$\hat{\mu}\{x \in \hat{X} : [x(f_1), \dots, x(f_n)] \in A\} = \mu\{x \in X : [(f_1, x), \dots, (f_n, x)] \in A\}.$$

By Kolmogorov's theorem, the premeasure $\hat{\mu}$ is completed in $\hat{X}$ to a countably additive measure defined on some σ -ring of subsets of the set $\hat{X}$.

Theorem 1. *The premeasure μ is completed in X to a measure if and only if the image of X under the mapping $x \rightarrow x(f) \in \hat{X}$ fills, in the space $\hat{X}$, a set of outer $\hat{\mu}$ -measure 1.*

A premeasure μ is called **Gaussian** if to each collection of functionals $f = (f_1, \dots, f_n)$ there correspond a positive quadratic form $Q_f(\xi, \xi)$, $\xi \in R^n$, and a constant c_f such that, for any Borel set $A \subset R^n$,

$$\mu C(f_1, \dots, f_n, A) = c_f \int_A \exp[-Q_f(\xi, \xi)] d\xi_1 \dots d\xi_n.$$

If, for the space X with Gaussian premeasure μ , a countable subsystem $g_1, g_2, \dots$ has been selected in X' , then, applying the orthogonalization process to it, one can pass to a new countable subsystem $e_1, e_2, \dots$ in X' (with the same linear hull), for which

$$\mu C(e_1, \dots, e_n, A) = \frac{1}{\sqrt{\pi^n}} \int_A \exp \left[-\sum_1^n \xi_k^2 \right] d\xi_1 \dots d\xi_n.$$

If the system $g_1, g_2, \dots$ is total, then the system $e_1, e_2, \dots$ is also total; in this case the space $\hat{X}$ is the space Ω of all real sequences $x = (x_1, x_2, \dots)$ with the canonical Gaussian measure

$$\omega\{x \in \Omega : (x_1, \dots, x_n) \in A\} = \frac{1}{\sqrt{\pi^n}} \int_A \exp \left[-\sum_1^n \xi_k^2 \right] d\xi_1 \dots d\xi_n.$$

By virtue of Theorem 1, the Gaussian premeasure μ extends on X to a measure if and only if the image of X under the mapping

$$x \rightarrow [(e_1, x), (e_2, x), \dots] \in \Omega$$

fills in Ω a set of outer measure 1. Therefore, for applications it is essential to have a sufficiently broad supply of subsets of Ω of full measure. We give several examples (the original formulation of which was given earlier in the language of probability theory):

Example 1 (B. V. Gnedenko ⁽⁴⁾).

$$\omega \left\{ x \in \Omega : \lim_{n \rightarrow \infty} \frac{|x_n|}{\sqrt{\ln n}} = 1 \right\} = 1.$$

Example 2 (A. N. Kolmogorov ⁽⁵⁾). Let a function $g(\xi)$, $-\infty < \xi < \infty$, be given such that

$$\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} g(\xi) e^{-\xi^2} d\xi = a, \quad \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} g^2(\xi) e^{-\xi^2} d\xi < \infty.$$

Then

$$\omega \left\{ x \in \Omega : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_1^n g(x_k) = a \right\} = 1.$$

Example 3 (A. N. Kolmogorov and A. Ya. Khinchin ⁽⁶⁾). Let positive numbers $a_1, a_2, \dots$ be given; set

$$E_{\langle a_n \rangle} = \left\{ x \in \Omega : \sum_1^\infty a_n x_n^2 < \infty \right\}.$$

Then

$$\omega E_{\langle a_n \rangle} = \begin{cases} 1, & \text{if } \sum_1^\infty a_n < \infty, \\ 0, & \text{if } \sum_1^\infty a_n = \infty. \end{cases}$$

Let $f_1, f_2, \dots$ be a numerical sequence. By Kolmogorov's theorem ⁽⁷⁾, the series

$$[f, x] \equiv \sum_1^\infty f_n x_n$$

converges on Ω in the mean square

(and also almost everywhere) if and only if $\sum_1^\infty f_n^2 < \infty$. In this case

$$\omega \{ x \in \Omega : [f, x] > c \} = \frac{1}{\sqrt{\pi}} \int_{c/(\sum_1^\infty f_n^2)^{1/2}}^\infty e^{-\xi^2} d\xi.$$

The expression $[f, x]$ is the general form of a linear measurable functional on Ω ⁽⁸⁾. If several sequences $f_1^j, f_2^j, \dots$ ($j = 1, \dots, m$), orthogonal and normalized in the space l_2 , are given, then

$$\omega\{x \in \Omega : ([f^1, x], \dots, [f^m, x]) \in A\} = \frac{1}{\sqrt{\pi^m}} \int_A \exp \left[-\sum_1^m \xi_k^2 \right] d\xi_1 \dots d\xi_m.$$

It follows that a linear transformation of the space Ω into itself, given by the formulas

$$y_n = \sum_{m=1}^{\infty} u_{mn} x_m$$

with an orthogonal matrix $U = \|u_{mn}\|$, carries measurable sets $E \subset \Omega$ again into measurable sets of the same measure. Fan Dyk Tinh showed in ⁽⁸⁾ that the indicated transformation is the general form of a measurable linear transformation in Ω preserving measure. There he also indicated the general form of linear transformations $y = Bx$ that carry measurable sets into measurable sets and do not preserve measure, but have an absolutely continuous set function $\omega(BE)$.

Wiener measure. Let H be a separable Hilbert space and $e_1, e_2, \dots$ an orthonormal basis in H . Each vector $z \in H$ is represented by a sequence of numbers $z_n = (e_n, z)$ with $\sum_1^{\infty} z_n^2 < \infty$. Fix a sequence of positive numbers $\lambda_1, \lambda_2, \dots$ and associate with the point $z \in H$ the point $(\lambda_1 z_1, \lambda_2 z_2, \dots)$ of the space Ω . Introduce in H the premeasure

$$\mu\{z \in H : (z_1, \dots, z_n) \in A \subset R_n\} = \omega\{x \in \Omega : (\lambda_1 z_1, \dots, \lambda_n z_n) \in A\}.$$

The premeasure μ is extended in H to a measure if the image of H in Ω fills a set of full measure (Theorem 1). By the Kolmogorov-Khinchin criterion ⁽³⁾, this condition is satisfied if and only if the series $\sum_1^{\infty} \lambda_n^{-2}$ converges.

The measure in the space H obtained for $\lambda_n \equiv n$ will be called the **abstract Wiener measure**. If as H one takes the space $L_2(0, \pi)$ of all square-integrable functions $z(t)$ on the interval $[0, \pi]$, orthogonal to 1, and as basis vectors e_n takes the functions

$$e_n(t) = \sqrt{\frac{2}{\pi}} \cos nt,$$

then we obtain the **classical Wiener measure**. The functions $z(t) \in \bar{L}_2(0, \pi)$ that are continuous and satisfy the Hölder condition $|z(t') - z(t'')| \leq C|t' - t''|^\alpha$ with any exponent $\alpha < 1/2$ fill in $\bar{L}_2(0, \pi)$ a set W of full measure. If from the functions $z(t) \in W$ we pass to the functions $\varphi(t) = z(t) - z(0)$, equal to 0 for $t = 0$, we obtain the second classical realization of Wiener measure ⁽⁹⁾. Kolmogorov's criterion 2, after a certain modification, leads to a set of full measure (indicated by Cameron and Martin ⁽¹⁰⁾ for the special case $g(\xi) = \xi^2$):

$$\omega \left\{ \varphi(t) : \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n g \left[\frac{\varphi\left(\frac{k}{n}\pi\right) - \varphi\left(\frac{k-1}{n}\pi\right)}{\sqrt{\pi/n}} \right] = a \right\} = 1.$$

Moscow State University
named after M. V. Lomonosov

Received
28 IX 1965

REFERENCES

1. A. N. Kolmogorov, *Basic Concepts of Probability Theory*, Moscow-Leningrad, 1936, pp. 39-42.
2. Yu. V. Prokhorov, The Method of Characteristic Functionals, Proc. IV Berkeley Symp. on Math. Statistics and Probability, 1960, 2, 1961, p. 403.
3. I. M. Gelfand, N. Ya. Vilenkin, *Generalized Functions*, vol. 4, ch. IV, 1961.
4. B. V. Gnedenko, Ann. Math., 44, 423 (1943).
5. A. N. Kolmogorov, C. R., 185, 949 (1927).
6. A. N. Kolmogorov, A. Ya. Khinchin, Math. collection, 32, 668 (1925).
7. A. N. Kolmogorov, Math. Ann., 99, 309 (1928).
8. Fan Dyk Tinh, UMN, 20, No. 3 (123), 244 (1965).
9. N. Wiener, R. Paley, *Fourier Transform in the Complex Domain*, ch. 9, "Nauka," 1964.
10. R. H. Cameron, W. T. Martin, Bull. Am. Math. Soc., 53, No. 2, 130 (1947).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.