

ASYMPTOTICS OF EIGENVALUES AND EIGENFUNCTIONS OF ONE-DIMENSIONAL SINGULAR DIFFERENTIAL OPERATORS

M. V. Fedoryuk

1966

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196601.92258>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.941.92

MATHEMATICS

M. V. FEDORYUK

ASYMPTOTICS OF EIGENVALUES AND EIGENFUNCTIONS OF ONE-DIMENSIONAL SINGULAR DIFFERENTIAL OPERATORS

(Presented by Academician I. M. Vinogradov on 29 X 1965)

I. Consider the system

$$y' = \mu A(x)y, \quad (1)$$

where $A(x)$ is a square matrix of order n , y is a vector with n components, and μ is a parameter. Everywhere in what follows it is assumed that, for $x \geq 0$, the matrix $A''(x)$ is continuous (the prime denotes differentiation with respect to x ; the matrix $A(x)$ is complex-valued). In this part of the article we study the asymptotics of solutions of system (1) as $x \rightarrow +\infty$.

We describe the form of the matrix $A(x)$.

1°. $A(x) = q(x)Q(x)B(x)Q^{-1}(x)$, where $q(x) \neq 0$ for $x \geq 0$ and Q is a diagonal matrix; $(Q(x))_{ii} = q^{\alpha_i}(x)$; α_i are complex numbers.

2°. The matrix $B(+\infty)$ exists, is finite, nondegenerate, and has distinct eigenvalues η_j .

3°.

$$\lim_{x \rightarrow +\infty} (|q'(x)| |\bar{q}(x)|^{-2} + |\bar{q}(x)|^{-1} \|B'(x)\|) = 0,$$

where

$$\|B(x)\| = \max_{i,j} |b_{ij}(x)|.$$

4°. The function

$$F(x) = |q''q^{-3}| + |q''q^{-2}| + (|q|^{-1} + |q'q^{-2}|) \|B'\| + |q|^{-1} (\|B'\|^2 + \|B''\|)$$

is summable on the interval $[0, +\infty)$.

Denote by $\lambda_j(x)$ the eigenvalues of the matrix $A(x)$, by $e_j(x), e_j^*(x)$ the right and left eigenvectors of this matrix corresponding to the eigenvalue $\lambda_j(x)$, and by $T(x)$ the matrix whose j -th column is $e_j(x)$, $1 \leq j \leq n$. Put

$$\lambda_j^{(1)}(x) = -(T^{-1}(x)T'(x))_{jj} \quad (2)$$

and introduce the diagonal matrices $\Lambda(x), \Lambda_1(x)$:

$$(\Lambda)_{jj} = \lambda_j(x), \quad (\Lambda_1)_{jj} = \lambda_j^{(1)}(x). \quad (3)$$

All these notations, by virtue of 2° , have meaning for sufficiently large x .

Theorem 1. Suppose that conditions $1^\circ-4^\circ$ are satisfied,

$$\int_0^\infty |q(x)| dx = \infty, \quad (4)$$

and, for sufficiently large x , $i \neq j$,

$$|\operatorname{Re}(\lambda_i(x) - \lambda_j(x))| \geq C|q(x)|, \quad C > 0. \quad (5)$$

Then for every $\mu_0 > 0$ there exists $x(\mu_0) < \infty$ such that, for $x \geq x(\mu_0)$, $\mu \geq \mu_0$, there exists a fundamental matrix $Y(x, \mu)$ of system (1) having the form

$$Y = T(x)(E + \mu^{-1}U(x, \mu)) \exp \left(\int_{x_0}^x (\mu\Lambda(t) + \Lambda_1(t)) dt \right). \quad (6)$$

$$\|U(x, \mu)\| \leq u(x), \quad \lim_{x \rightarrow +\infty} u(x) = 0. \quad (7)$$

Let

$$\lim_{x \rightarrow +\infty} |q(x)| = +\infty, \quad \lim_{x \rightarrow +\infty} \arg q(x) = \varphi_0. \quad (8)$$

Denote

$$\eta_j' = \eta_j e^{i\varphi_0} \quad (9)$$

where η_j are the eigenvalues of the matrix $B(+\infty)$, and m^-, m^+ are the numbers of points η_j' lying respectively in the left and right half-planes, and introduce the operator

$$l = d/dx - A(x).$$

From Theorem 1 it follows:

Theorem 2. *Suppose that the conditions of Theorem 1 and condition (8) are satisfied and*

$$\operatorname{Re} \eta'_j \neq 0, \quad 1 \leq j \leq n; \quad \operatorname{Re}(\eta'_i - \eta'_j) \neq 0, \quad i \neq j. \quad (10)$$

Then, for any complex μ , the maximal number of linearly independent solutions of the system

$$ly = \mu y, \quad (11)$$

belonging to $L_2[0, +\infty)$, is equal to m^- .

Denote by D the collection of all vectors $y \in L_2[0, +\infty)$, all components of which are absolutely continuous on every finite interval $[0, a]$, $a > 0$, and such that $ly \in L_2[0, +\infty)$. By D_Ω we denote the collection of all vectors $y \in D$ such that

$$\Omega y(0) = 0, \quad (12)$$

where Ω is a constant matrix of rank m^- . By L_Ω we denote the operator in $L_2[0, +\infty)$ with domain of definition D_Ω , and $L_\Omega y = ly$ for $y \in D_\Omega$.

Theorem 3. *If the conditions of Theorem 2 are satisfied, then the spectrum of the operator L_Ω is purely discrete and has no finite limit points.*

Let us discuss the conditions of Theorem 1. If one requires that the matrix $A(x)$ be nondegenerate in a certain sense (see, for example, (1)) and that the $\lambda_j(x)$ have the same order of growth as $x \rightarrow +\infty$, then $A(x)$ has the form 1°, 2°. In particular, for the scalar equation $y^{(n)} + \mu^n p(x)y = 0$ these conditions are automatically satisfied. If the characteristic roots of the scalar equation

$$\sum_{k=0}^n q_k(x)y^{(k)} = 0, \quad q_n(x) \equiv 1,$$

have the same order of growth as $x \rightarrow +\infty$, then the matrix $A(x)$ has the form 1°, and $B(+\infty)$ exists and is finite; moreover $a_j = j - 1$. Therefore the main results of (3) are contained in Theorem 1. Theorem 1 also remains valid in the case where condition 2° is replaced by

2°'. The matrix $B(+\infty)$ has no multiple roots.

Condition 3° is a restriction on the rate of decrease of $q(x)$ as $x \rightarrow +\infty$, and condition 4° is a restriction on the regularity of the behavior of $q(x)$ and $B(x)$ as $x \rightarrow +\infty$. For example, if $\|B^{(j)}(x)\| = O(x^{-j})$, $x \rightarrow +\infty$, $j = 1, 2$, and $q(x) = x^\alpha$, $\operatorname{Re} \alpha > -1$, then conditions 3°, 4° are satisfied. These conditions are also satisfied if $q(x)$ grows like $(\ln x)^\beta$ or like $\exp(Cx^\gamma)$, $\gamma > 0$, $\operatorname{Re} C > 0$, with arbitrary β , and this asymptotic expression can be differentiated twice. Condition (4) is satisfied in these examples. Finally, condition (5) can be weakened. In particular, if $A(x)$ is real, then it is sufficient to require (5) only for those pairs $\lambda_i(x), \lambda_j(x)$ which are not complex conjugates. G. Birkhoff⁽²⁾ investigated the case where $\mu = 1$, $Q(x) = E$, $q(x) = x^r$, where $r \geq 0$ is an integer and $B(+\infty)$ has distinct eigenvalues; these results are contained in Theorem 1.

- II. Consider the eigenvalue problem on the whole axis $(-\infty, +\infty)$ for system (1). A solution $y(x, \mu_0) \neq 0$ belonging to $L_2(-\infty, +\infty)$ is called an eigenfunction, and μ_0 is called an eigenvalue.

value. A point x_0 is called a turning point of system (1) if the matrix $A(x_0)$ has multiple eigenvalues. A turning point x_0 is called simple if $A(x_0)$ has a double eigenvalue λ_0 , the remaining eigenvalues are simple, and $f_x(\lambda_0, x_0) \neq 0$, where $f(\lambda, x)$ is the characteristic polynomial of the matrix $A(x)$.

Introduce the conditions:

1. System (1) has on the real axis exactly two, and moreover simple, turning points $x_1 < x_2$, and $\lambda_m(x_j) = \lambda_{m+1}(x_j)$, $j = 1, 2$.
2. $\operatorname{Re}(\lambda_m(x) - \lambda_{m+1}(x)) \equiv 0$ for $x_1 \leq x \leq x_2$, $\operatorname{Re} \lambda_m(x) - \operatorname{Re} \lambda_{m+1}(x) < 0$ for $x > x_2$, $x < x_1$.
3. For all real x ,

$$\operatorname{Re}(\lambda_j(x) - \lambda_{j+1}(x)) < 0, \quad j \neq m, n.$$

In particular, for the scalar equation $y'' - \mu^2 q(x)y = 0$, conditions 1-3 are satisfied if $q(x)$ is real for real x , has exactly two, and moreover simple, zeros $x_1 < x_2$ on the real axis, and $q(x) > 0$ for $x > x_2$.

Denote by $S(\rho, \alpha, \beta)$ the sector $|\mu| \geq \rho$, $\alpha < \arg \mu < \beta$.

Theorem 4. Suppose the conditions of Theorem 1 and conditions 1-3 are satisfied on the real axis, $1 \leq m \leq n-1$, and the matrix $A(x)$ is regular on the interval $[x_1, x_2]$. Then there exist $\rho > 0$, $\alpha < 0 < \beta$ such that all eigenvalues of system (1) (for the problem on the whole axis) lying in $S(\rho, \alpha, \beta)$ have the form

$$\mu_k = \xi_0^{-1}(2\pi ki + \xi_1) + O(k^{-1}), \quad k_0 \leq k < +\infty. \quad (13)$$

Here

$$\xi_0 = \frac{1}{2} \int_C (\lambda_{m+1} - \lambda_m) dt, \quad \xi_1 = \frac{1}{2} \int_C (\lambda_{m+1}^{(1)} - \lambda_m^{(1)}) dt, \quad (14)$$

where C is a closed contour in the complex x -plane which contains the interval $[x_1, x_2]$ in its interior, traverses it in the positive direction, and contains no

other turning points of system (1) in its interior. The branches of the functions $\lambda_{m+1}, \lambda_m, \lambda_{m+1}^{(1)}, \lambda_m^{(1)}$ are chosen for $x \in C$, $x > x_2$.

We note that ξ_0 and ξ_1 are purely imaginary quantities and $\text{Im} \xi_0 > 0$. In addition, the matrices $B(+\infty)$ and $B(-\infty)$ may be different, and $q(x)$ may have different orders of growth as $x \rightarrow \pm\infty$.

Corollary. *Let γ_0 be a smooth curve in the complex x -plane connecting two real points $x_3 < x_4$, and let*

$$\gamma = (-\infty, x_3] \cup \gamma_0 \cup [x_4, +\infty).$$

Suppose that on γ the conditions of Theorem 1, condition 1, and the following conditions hold:

$$2'. \quad \text{Re} \left(\int_{x_1}^x (\lambda_{m+1} - \lambda_m) dt \right) \equiv 0, \quad \text{if } x \in \gamma \text{ and lies between } x_1 \text{ and } x_2;$$

$$\text{Re} \left(\int_{x_1}^x (\lambda_{m+1} - \lambda_m) dt \right) > 0 \quad \text{for } x \in \gamma \text{ to the right of } x_2, \text{ and is less than zero for } x \in \gamma \text{ to the left of } x_1.$$

$$3'. \quad \text{Re} \left(\int_{x_2}^x (\lambda_j - \lambda_{j+1}) dt \right) < 0 \quad \text{for } j \neq m, x \in \gamma \text{ and lying to the right of } x_2,$$

and this function is positive for $x \in \gamma$ to the left of x_2 .

Suppose, moreover, that the matrix $A(x)$ is regular for $x \in \overline{\gamma_0}$. Then all conclusions of Theorem 4 remain valid.

For the scalar equation $y'' - \mu^2 p(x)y = 0$, this question has been investigated more fully in ^(4,5).

Our methods also make it possible to find the asymptotics of the eigenfunctions as $\mu \rightarrow +\infty$ (in the case of Theorem 4) on the whole real axis, except for certain neighborhoods of the turning points.

Introduce the conditions:

1) System (1) has turning points on the real axis

$x_1 < x_2 < \dots < x_{2l}$, all of them are simple, and $\lambda_{m-j+1}(x_j) = \lambda_{m+j}(x_{2l-j+1})$, $j = 1, 2, \dots, l$.

2) $\text{Re}(\lambda_{m-j+1}(x) - \lambda_{m+j}(x)) \equiv 0$ for $x \in [x_j, x_{2l-j+1}]$.

- 3) $\operatorname{Re}(\lambda_j(x) - \lambda_{j+1}(x)) < 0$ for all x, j , with the exception of those indicated in 2), and $\operatorname{Re} \lambda_{m+1}(x) > 0$, $\operatorname{Re} \lambda_m(x) < 0$ for $x \in [x_1, x_{2l}]$.

Theorem 5. *Suppose that the hypotheses of Theorem 1 and conditions 1)–3) are satisfied and that the matrix $A(x)$ is regular for $x \in [x_1, x_{2l}]$. Then there exist $\rho > 0$, $\alpha < 0 < \beta$ such that all eigenvalues of system (1) lying in $S(\rho, \alpha, \beta)$ have the form*

$$\mu_{kj} = \xi_{0j}^{-1}(2\pi ki + \xi_{1j}) + O(k^{-1}), \quad j = 1, 2, \dots, l; \quad k \geq k_0. \quad (15)$$

Here ξ_{0j}, ξ_{1j} have the form (14), where the integral is taken over the contour C_j enclosing the segment $[x_j, x_{2l-j+1}]$.

The quantities ξ_{0j} and ξ_{1j} are purely imaginary, and $\operatorname{Im} \xi_{0j} > 0$. The corollary from Theorem 4 remains valid. Similar formulas are obtained for the asymptotics of the eigenvalues for a problem on a half-line (say, on the half-line $x \geq x_0$, $x_1 < x_0 < x_2$, under the hypotheses of Theorem 4).

Moscow Institute of Physics and Technology

Received

18 X 1965

REFERENCES

- ¹ L. R. Volevich, *Uspekhi Mat. Nauk*, 18, No. 3, 155 (1963).
- ² G. Birkhoff, *Trans. Am. Math. Soc.*, 10, 436 (1909).
- ³ M. V. Fedoruk, *Dokl. Akad. Nauk*, 165, No. 4 (1965).
- ⁴ Yu. N. Dnestrovsky, L. P. Kostomarov, *Zh. Vychisl. Mat. Mat. Fiz.*, 4, No. 2, 267 (1964).
- ⁵ M. V. Fedoruk, *Mat. Sb.*, 68 (110), No. 1, 68 (1965).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.