

ON CONJUGATE FUNCTIONS

L. V. Zhizhiashvili

1966

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196601.92817>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.51

MATHEMATICS

L. V. ZHIZHIASHVILI

ON CONJUGATE FUNCTIONS

(Presented by Academician I. N. Vekua on June 22, 1965)

1. Let the 2π -periodic function $f(x)$ be summable on $[-\pi, \pi]$. Denote by $\sigma[f]$ the Fourier series of the function $f(x)$, and by $s_n(x, f)$ the partial sums of the series $\sigma[f]$. Further, as usual, by the symbol $\bar{f}(x)$ we shall denote the function conjugate to $f(x)$, i.e.

$$\bar{f}(x) = -\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x+t) \operatorname{ctg} \frac{t}{2} dt,$$

which exists almost everywhere for every summable (see ⁽¹⁾, p. 528) function $f(x)$. It is known (see ⁽¹⁾, p. 557) that if $f(x)$ is a bounded 2π -periodic function, then $\bar{f}(x)$ need not be bounded. However, as was shown by P. Turán ⁽²⁾ and M. Kinukawa ⁽³⁾, if $f(x)$ is an even 2π -periodic function bounded on $(-\infty, +\infty)$, then the functions

$$\varphi(x) = \frac{1}{\operatorname{tg} x/2} \int_0^x \bar{f}(t) dt, \quad \psi(x) = \int_x^\pi \frac{\bar{f}(t)}{2 \operatorname{tg} t/2} dt \quad (|x| \leq \pi)$$

are bounded on $(-\infty, +\infty)^*$. The behavior of the partial sums of the series $\sigma[\varphi]$ and $\sigma[\psi]$ was studied by M. and S. Izumi ⁽⁴⁾. In particular, they showed that if $f(x)$ is an even bounded function and $\|s_n(x, f)\|_C = O(1)$, then $\|s_n(x, \varphi)\|_C = O(1)$, $\|s_n(x, \psi)\|_C = O(1)$. The question arises: what can be said about the continuity of the functions $\varphi(x)$ and $\psi(x)$, or about the uniform convergence of the series $\sigma[\varphi]$ and $\sigma[\psi]$?

Theorem 1. If $f(x)$ is an even 2π -periodic continuous function, then the function $\varphi(x)$ is also continuous. Moreover, if

$$f(0) = \int_0^\pi f(t) dt = 0,$$

then the function $\psi(x)$ is also continuous, and its modulus of continuity is

$$\omega(\delta, \psi) = O \left\{ \omega(\delta, f) + \delta \int_{\delta}^1 \frac{\omega(t, f)}{t^2} dt \right\},$$

where $\omega(\delta, u)$ is the modulus of continuity of the function $u(x) \in C(-\pi, \pi)$.

Theorem 2. Let $f(x)$ be an even 2π -periodic continuous function and $|s_n(0, f)| \leq M$ ($n = 1, 2, \dots$). Then the series $\sigma[f]$ converges uniformly. Moreover, if

$$f(0) = \int_0^{\pi} f(t) dt = 0,$$

then the series $\sigma[\psi]$ also converges uniformly.

Analogous assertions for the functions $\varphi(x)$ and $\psi(x)$ in the case when $f(x)$ is odd are, generally speaking, false.

2. Let us now consider a function of two variables $f(x, y)$. Suppose that it is periodic with respect to each of the variables and $f(x, y) \in$

* Here it is assumed that the functions $\varphi(x)$ and $\psi(x)$ are extended periodically with period 2π from the interval $[-\pi, \pi]$ to the entire line.

$\in L(R)$, where $\bar{R} = [-\pi, \pi, -\pi, \pi]$. Consider the conjugate functions of two variables

$$\begin{aligned} \bar{f}_1(x, y) &= -\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x+s, y) \operatorname{ctg} \frac{s}{2} ds, \\ \bar{f}_2(x, y) &= -\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x, y+t) \operatorname{ctg} \frac{t}{2} dt, \\ \bar{f}_3(x, y) &= \frac{1}{4\pi^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} f(x+s, y+t) \operatorname{ctg} \frac{s}{2} \operatorname{ctg} \frac{t}{2} ds dt. \end{aligned}$$

If $f(x, y)$ is an even function with respect to the aggregate of the two variables, i.e. $f(-x, -y) = f(x, y)$, then, generally speaking, the assertions of P. Turán ⁽²⁾ and M. Kinukawa ⁽³⁾ are false ⁽⁵⁾ for the functions $\bar{f}_i(x, y)$ ($i = 1, 2, 3$).

Let us now suppose that $f(x, y)$ is an even function with respect to each variable, i.e. $f(-x, y) = f(x, -y) = f(x, y)$. Further, let

$$\varphi(x, y) = \frac{1}{\operatorname{tg} x/2 \operatorname{tg} y/2} \int_0^x \int_0^y \bar{f}_3(s, t) ds dt, \quad \psi(x, y) = \int_x^{\pi} \int_y^{\pi} \frac{\bar{f}_3(s, t)}{4 \operatorname{tg} s/2 \operatorname{tg} t/2} ds dt,$$

$(x, y) \in R$, and

$$\varphi(x + 2\pi, y) = \varphi(x, y + 2\pi) = \varphi(x, y), \quad \psi(x + 2\pi, y) = \psi(x, y + 2\pi) = \psi(x, y)$$

for all x, y .

Lemma 1. If $f(x, y)$ is a bounded function, even with respect to each variable, then the relations

$$\begin{aligned} \varphi(x, y) &= \frac{1}{\pi^2 \operatorname{tg} x/2 \operatorname{tg} y/2} \int_0^\pi \int_0^\pi f(s, t) \log \left| \frac{\sin(x+s)/2 \sin(x-s)/2}{\sin^2 s/2} \right| \\ &\quad \times \log \left| \frac{\sin(y+t)/2 \sin(y-t)/2}{\sin^2 t/2} \right| ds dt, \\ \psi(x, y) &= \frac{1}{4\pi^2} \int_0^\pi \int_0^\pi f(s, t) \operatorname{ctg} \frac{s}{2} \operatorname{ctg} \frac{t}{2} \log \left| \frac{\sin(x+s)/2}{\sin(x-s)/2} \right| \log \left| \frac{\sin(y+t)/2}{\sin(y-t)/2} \right| ds dt \\ &\quad - \frac{\pi-x}{4\pi^2} \int_0^\pi \int_0^\pi f(s, t) \operatorname{ctg} \frac{t}{2} \log \left| \frac{\sin(y+t)/2}{\sin(y-t)/2} \right| ds dt \\ &\quad - \frac{\pi-y}{4\pi^2} \int_0^\pi \int_0^\pi f(s, t) \operatorname{ctg} \frac{s}{2} \log \left| \frac{\sin(x+s)/2}{\sin(x-s)/2} \right| ds dt \\ &\quad + \frac{(\pi-x)(\pi-y)}{16\pi^2} \int_0^\pi \int_0^\pi f(s, t) ds dt. \end{aligned}$$

hold.

On the basis of this lemma Theorems 3 and 4 are proved.

Theorem 3. If $f(x, y)$ is a bounded even function (with respect to each variable), then the functions $\varphi(x, y)$ and $\psi(x, y)$ are also bounded.

Theorem 4. If $f(x, y)$ is a continuous even function (with respect to each variable), then the function $\varphi(x, y)$ is continuous. Moreover, if

$$f(0, 0) = \int_0^\pi \int_0^\pi f(s, t) ds dt = 0,$$

then the function $\psi(x, y)$ is continuous and its modulus of continuity

$$\omega(\delta, \delta'; u) = O \left\{ \omega_1(\delta, f) + \omega_2(\delta', f) + \delta \int_\delta^1 \frac{\omega_1(s, f)}{s^2} ds + \delta' \int_{\delta'}^1 \frac{\omega_2(t, f)}{t^2} dt \right\},$$

where $\omega(\delta, \delta'; u)$, $\omega_1(\delta, u)$ and $\omega_2(\delta', u)$ are, respectively, the full and partial moduli of continuity of the function $u(x, y) \in C(R)$.

Analogous assertions are also valid for the functions

$$\bar{f}_i(x, y) \quad (i = 1, 2).$$

3. Let now $f(x, y) \in L(R)$ and

$$F(x, y) = \int_0^x \int_0^y f(s, t) ds dt.$$

Put

$$\Delta_x(F; x, y, s) = F(x + s, y) + F(x - s, y) - 2F(x, y),$$

$$\Delta_y(F; x, y, t) = F(x, y + t) + F(x, y - t) - 2F(x, y),$$

$$\Delta^2(F; x, y, s, t) = \Delta_x(\Delta_y(F; x, y, t)) = \Delta_y(\Delta_x(F; x, y, s));$$

$$\tilde{F}_1(x, y) = \lim_{\varepsilon \rightarrow 0+} \int_{\varepsilon}^{\pi} \frac{\Delta_x(F; x, y, s)}{2 \sin^2 s/2} ds, \quad \tilde{F}_2(x, y) = \lim_{\eta \rightarrow 0+} \int_{\eta}^{\pi} \frac{\Delta_y(F; x, y, t)}{2 \sin^2 t/2} dt,$$

$$\tilde{F}_3(x, y) = \lim_{\varepsilon, \eta \rightarrow 0+} \int_{\varepsilon}^{\pi} \int_{\eta}^{\pi} \frac{\Delta^2(F; x, y, s, t)}{4 \sin^2 s/2 \sin^2 t/2} ds dt.$$

The question is: under what conditions do the functions $F_i(x, y)$ ($i = 1, 2, 3$) exist, and what is the relation between them?

The functions $\tilde{F}_i(x, y)$ ($i = 1, 2$) exist almost everywhere for every summable 2π -periodic function $f(x, y)$, which may be obtained from the corresponding result of A. Plessner ⁽⁶⁾. For the function $\tilde{F}_3(x, y)$ the analogous assertion is, generally speaking, false, since the following holds.

Theorem 5. There exists a function $f(x, y) \in L(\log^+ L)^\alpha$ for all $\alpha \in [0, 1)$, for which almost everywhere on R

$$\lim_{\varepsilon \rightarrow 0+} \int_{\varepsilon}^{\pi} \int_{\varepsilon}^{\pi} \frac{\Delta^2(F; x, y, s, t)}{4 \sin^2 s/2 \sin^2 t/2} ds dt$$

does not exist.

If, however, $f(x, y) \in L(\log^+ L)^\alpha$, $\alpha \geq 1$, then such an assertion is no longer valid, for the following is true.

Theorem 6. If $f(x, y) \in L \log^+ L$, then $\tilde{F}_3(x, y)$ exists almost everywhere and, for almost all x and y ,

$$\tilde{F}_3(x, y) = \tilde{F}_{i,j}(x, y) = \bar{f}_{i,j}(x, y) \quad (i, j = 1, 2; i \neq j)^*,$$

where

$$\tilde{F}_{i,j}(x, y) = \lim_{\varepsilon \rightarrow 0+} \int_{\varepsilon}^{\pi} \frac{\Delta_x(\tilde{F}_i; x, y, s)}{2 \sin^2 s/2} ds \quad \text{for } i = 2, j = 1.$$

The results given here can be generalized also to the case of functions of n variables. We shall confine ourselves to considering the analogue of Theorem 6.

Let $f(x_1, \dots, x_n) \in L(R')$, where $R' = [-\pi, \pi; \dots; -\pi, \pi]$. Put

$$F(x_1, \dots, x_n) = \int_0^{x_1} \dots \int_0^{x_n} f(s_1, \dots, s_n) ds_1 \dots ds_n,$$

$$\begin{aligned} \Delta_k F &\equiv \Delta_k(F; x_1, \dots, x_n, s_k) = F(x_1, \dots, x_k + s_k, \dots, x_n) + \\ &+ F(x_1, \dots, x_k - s_k, \dots, x_n) - 2F(x_1, \dots, x_n), \end{aligned}$$

$$\Delta^n(F; x_1, \dots, x_n; s_1, \dots, s_n) = \Delta_n(\Delta_{n-1}(\dots(\Delta_1 F)) = \dots = \Delta_1(\Delta_2(\dots(\Delta_n F) \dots)).$$

* The symbol $\bar{f}_{i,j}(x, y)$ means that the conjugation operation is applied to the function $f(x, y)$ first in the i -th

Theorem 7. Let $f(x_1, \dots, x_n) \in L(\log^+ L)^{n-1}$. Then almost everywhere on R' there exists

$$\lim_{(\varepsilon_1, \dots, \varepsilon_n) \rightarrow 0+} \int_{\varepsilon_1}^{\pi} \dots \int_{\varepsilon_n}^{\pi} \frac{\Delta^n(F; x_1, \dots, x_n; s_1, \dots, s_n)}{2^n \sin^2 s_1/2 \dots \sin^2 s_n/2} ds_1 \dots ds_n,$$

and almost everywhere

$$\tilde{F}_{2^n-1}(x_1, \dots, x_n) = \tilde{F}_{i_1, \dots, i_n}(x_1, \dots, x_n)$$

$(i_1, \dots, i_n = 1, 2, \dots, n, \text{ in the order } i_j \neq i_k),$

where

$$\begin{aligned}\tilde{F}_{i_k}(x_1, \dots, x_n) &= \lim_{\varepsilon_{i_k} \rightarrow 0+} \int_{\varepsilon_{i_k}}^{\pi} \frac{\Delta_{i_k}(F; x_1, \dots, x_n, s_{i_k})}{2 \sin^2 s_{i_k}/2} ds_{i_k}, \\ \tilde{F}_{i_k, i_j}(x_1, \dots, x_n) &= \lim_{\varepsilon_{i_j} \rightarrow 0+} \int_{\varepsilon_{i_j}}^{\pi} \frac{\Delta_{i_j}(\tilde{F}_{i_k}; x_1, \dots, x_n, s_{i_j})}{2 \sin^2 s_{i_j}/2} ds_{i_j},\end{aligned}$$

and the assertion loses its force if, in the hypothesis of the theorem, the exponent $n - 1$ is replaced by a smaller one.

I take this opportunity to express my deep gratitude to P. L. Ulyanov for valuable advice in carrying out this work.

Tbilisi
State University

Received
10 VI 1965

REFERENCES

- ¹ N. K. Bari, *Trigonometric Series*, 1961.
- ² P. Turan, Ann. Soc. Polon. Math., **25**, 155 (1952).
- ³ M. Kinukawa, Dissertation Northwestern Univ., 1960.
- ⁴ M. Izumi, S. Izumi, Acta Math. Sci. Hung., **13**, No. 1-2, 133 (1962).
- ⁵ L. V. Zhizhiashvili, DAN, **155**, No. 3, 521 (1964).
- ⁶ A. Plessner, J. Math., **159**, 219 (1927).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.