

ON DISSIPATIVE OPERATORS WITH ABSOLUTELY CONTINUOUS SPECTRUM

L. A. Sakhnovich

1966

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196601.94509>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.432

MATHEMATICS

L. A. SAKHNOVICH

ON DISSIPATIVE OPERATORS WITH ABSOLUTELY CONTINUOUS SPECTRUM

(Presented by Academician M. V. Keldysh on 30 VI 1965)

§ 1. In the present paper we consider a non-self-adjoint operator A of class $i\Omega$ ⁽¹⁾ with continuous spectrum. In addition, we shall assume that the operator A is dissipative, i.e. $(A - A^*)/i \geq 0$. As M. S. Livshits ⁽¹⁾ showed, the characteristic matrix-function of such an operator is represented in the form

$$w(\lambda) = \int_0^l \exp \left[-\frac{i\beta^2(t)}{\alpha(t) - \lambda} dt \right],$$

where the function $\alpha(t)$ increases monotonically and is bounded, while the matrix $\beta(x)$ is nonnegative and $\text{sp } \beta^2(x) \equiv 1$.

Assume additionally that the operator A has absolutely continuous spectrum, i.e. the function $t = \alpha(x)$ has an absolutely continuous inverse function $x = \sigma(t)$. In this case

$$w(\lambda) = \int_0^b \exp \left[-\frac{i\beta_1^2(t)}{t - \lambda} dt \right],$$

where $\beta_1(t) = p(t)\beta(\sigma(t))$, $p(t) = \sqrt{\sigma'(t)}$, $a = \alpha(0)$, $b = \alpha(l)$.

The triangular model of the operator A , as follows from ⁽¹⁾, can be written in the form

$$\vec{A}f = xf(x) + i \int_a^x f(t)\beta_1(t) dt \beta_1(x) \quad (a \leq x \leq b). \quad (1)$$

Theorem 1. *The additional component of the operator $\vec{A}$ consists of those and only those vector-functions $f(x) \in L_r^2[a, b]$ for which, almost everywhere, the equality*

$$f(x)\beta_1(x) \equiv 0, \quad \text{if } \|\beta_1(x)\| \leq M.$$

§ 2. For what follows, the behavior of the multiplicative integral

$$w(b, \lambda) = \int_a^b \exp \left[-\frac{i\beta_1^2(t)}{t-\lambda} dt \right] \quad \left(\int_a^b \|\beta_1^2(t)\| dt < \infty \right) \quad (2)$$

as $\tau \rightarrow 0$ ($\lambda = \sigma + i\tau$) is essential.

Theorem 2. *For almost all $\sigma \in [a, b]$ there exist limiting values*

$$w^\pm(\sigma) = \lim_{\tau \rightarrow \pm 0} w(b, \lambda)$$

and the formulas hold

$$w^\pm(\sigma) = \lim_{\varepsilon \rightarrow 0} \int_a^{\sigma-\varepsilon} \exp \left[-\frac{i\beta_1^2(t)}{t-\gamma} dt \right] \exp[\pm \pi \beta_1^2(\sigma)] \int_{\sigma+\varepsilon}^b \exp \left[-\frac{i\beta_1^2(t)}{t-\gamma} dt \right], \quad (3)$$

where the limits are understood in the sense of strong convergence.

The theorem was previously proved by us under the condition that $\text{sp } \beta_1^2(t)$ is bounded ^(2,3).

From formula (3) it follows that

$$w^\pm(\sigma) = R^{\pm 1}(\sigma) u(b, \sigma), \quad (4)$$

where

$$\begin{aligned} R^{\pm 1}(\sigma) &= \lim_{\varepsilon \rightarrow +0} \int_a^{\sigma-\varepsilon} \exp \left[-\frac{i\beta_1^2(t)}{t-\sigma} dt \right] \exp[\pm \pi \beta_1^2(\sigma)] \left[\int_a^{\sigma-\varepsilon} \exp \left[-\frac{i\beta_1^2(t)}{t-\sigma} dt \right] \right]^{-1} = \\ &= \exp[\pm \pi r^2(\sigma)], \end{aligned} \quad (5)$$

$$U(b, \sigma) = \lim_{\varepsilon \rightarrow +0} \int_a^{\sigma-\varepsilon} \exp \left[-\frac{i\beta_1^2(t)}{t-\sigma} dt \right] \int_{\sigma+\varepsilon}^b \exp \left[-i\frac{\beta_1^2(t)}{t-\sigma} dt \right]. \quad (6)$$

Let H and H_1 denote the closures of the manifolds into which $L_r^2[a, b]$ is mapped upon multiplication of its elements respectively by $\beta_1(x)$ and $R(x) - R^{-1}(x)$.

By Theorem 1, the operator $\vec{A}$ induces its simple part on H .

In what follows we shall consider the operator $\vec{A}$ only on the space H .

In ⁽⁴⁻⁶⁾ we constructed mutually inverse operators B and B^{-1} , defined on dense sets respectively in H_1 and H , by means of the formulas

$$B\varphi = \frac{1}{\sqrt{2\pi}} \frac{d}{dx} \int_a^x \varphi(\sigma) \sqrt{R(\sigma) - R^{-1}(\sigma)} U(x, \sigma) d\sigma \beta_1^{-1}(x), \quad (7)$$

$$B^{-1}f = \frac{1}{\sqrt{2\pi}} \left\{ \int_a^x [f(\sigma) \beta_1^{-1}(\sigma)]' U^*(\sigma, x) d\sigma + f(a) \beta_1^{-1}(a) \right\} \sqrt{R(x) - R^{-1}(x)}. \quad (8)$$

Here

$$U(x, \sigma) = \lim_{\varepsilon \rightarrow +0} \int_a^{\sigma-\varepsilon} \exp \left[-\frac{i\beta_1^2(t)}{t-\sigma} dt \right] \int_{\sigma+\varepsilon}^b \exp \left[-\frac{i\beta_1^2(t)}{t-\sigma} dt \right] d\sigma.$$

Formulas (7)–(8) can be given meaning also in the case when the matrix $\beta_1(x)$ has no inverse on a set of positive measure ⁽⁴⁻⁶⁾.

Item 3. The relation holds

$$\vec{A} = BQB^{-1}, \quad (9)$$

where Q is the operator of multiplication by the independent variable,

$$Qf = xf, \quad f \in H_1.$$

We note that, in deriving relations (7)–(9), in works ⁽⁴⁻⁶⁾ we assumed $\text{sp } \beta_1^2(t)$ to be bounded. However, it is easy to dispense with this condition by using Theorem 2.

If $\vec{A}$ and Q are connected by a relation of the type (9), then they are called **linearly similar** ⁽⁴⁻⁵⁾. If, moreover, B and B^{-1} are bounded, then the operators $\vec{A}$ and Q are called **linearly equivalent**.

Theorem 3. The relations

$$\|e^{-\frac{\pi}{2}r^2} B^{-1}\| = 1, \quad \|Be^{-\frac{\pi}{2}r^2}\| = 1,$$

hold, where the operator $e^{-\frac{\pi}{2}r^2}$ is defined by the formula

$$e^{-\frac{\pi}{2}r^2} f = f(x) e^{-\frac{\pi}{2}r^2(x)}.$$

Theorem 4. In order that the operator $\vec{A}$ be linearly equivalent to a self-adjoint operator, it is necessary and sufficient that

$$\text{vrai sup } \|\beta_1^2(x)\| = M < \infty. \quad (10)$$

Corollary. If condition (10) is satisfied, then the operator $\vec{A}$ is equivalent to Q , and

$$\|B\| = e^{\frac{\pi}{2}M}, \quad \|B^{-1}\| = e^{\frac{\pi}{2}M}.$$

In terms of the characteristic matrix-function, Theorem 4 admits the following reformulation.

Theorem 4'*. In order that a dissipative operator A of class $i\Omega$ with absolutely continuous spectrum be linearly equivalent to a self-adjoint operator, it is necessary and sufficient that

$$\text{vrai sup } \|w^+(\sigma)\| < \infty.$$

Theorem 3 makes it possible to study the operator A also in the case when condition (10) is not satisfied.

Theorem 5. The multiplicity of the spectrum of the operator A is equal to

$$N = \text{vrai sup rang } \beta_1^2(x) = \text{vrai sup rang } [w^+(x) - w^-(x)].$$

Corollary. The operator A decomposes into the sum of N induced operators of first multiplicity.

Theorem 6. There exist invariant subspaces $H^{(n)}$ and $H_1^{(n)}$ of the operators A and Q , respectively, such that the operators $A^{(n)}$ and $Q^{(n)}$ induced on them are linearly equivalent. Moreover, the projection operators onto these subspaces satisfy

$$P^{(n)} \xrightarrow{n \rightarrow \infty} I, \quad P_1^{(n)} \xrightarrow{n \rightarrow \infty} I.$$

All the results remain valid also for operators not belonging to the class $i\Omega$, whose characteristic matrix-function admits the representation (2).

Let us note that the theorems formulated here can be carried over to the case when one or both endpoints of the segment $[a, b]$ are infinite. In this case the operators A and Q are unbounded.

4. Let us study the behavior of e^{iAt} as $t \rightarrow \pm\infty$. We shall compare it with the behavior of e^{iA_1t} as $t \rightarrow \pm\infty$, where A_1 is the real component of A ($A_1 = (A + A^*)/2$). Let G be the subspace corresponding to the absolutely continuous part of the spectrum of A_1 , and let P_G be the projection operator onto G .

* *Note added in proof.* A more general result has been obtained by B. S. Naim and K. Hoffman ⁽¹⁰⁾.

Theorem 7. Let the prime operator A satisfy the conditions of Theorem 4. Then the strong limits exist

$$W_{\pm}(A, A_1) = \lim_{t \rightarrow \pm\infty} e^{iAt} e^{-iA_1 t} P_G,$$

$$W_{\pm}(A_1, A) = \lim_{t \rightarrow \pm\infty} e^{iA_1 t} e^{-iAt}.$$

Moreover, the operators $W_{\pm}(A, A_1)$ and $W_{\pm}(A_1, A)$ are bounded together with their inverses, and the relations hold

$$A = W_{\pm}(A, A_1) A_1 W_{\pm}(A, A_1)^{-1}, \quad A_1 = W_{\pm}(A_1, A) A W_{\pm}(A_1, A)^{-1},$$

$$W_{\pm}(A, A_1) = W_{\pm}^{-1}(A_1, A),$$

where A_1 is considered only on G .

Thus, Theorem 7 extends to non-self-adjoint operators the well-known Rosenblum-Kato theorem ^(7,8). We note that some results in this direction were obtained earlier ⁽⁹⁾.

Introduce the scattering operator S by the formula

$$S = W_{-}^{-1}(A, A_1) W_{+}(A, A_1).$$

The operator S is defined in G and commutes in G with A , $\|S\| \leq 1$.

Theorem 8. The operator S is unitarily equivalent to multiplication by the matrix

$$S(x) = I - \sqrt{R(x) - R^{-1}(x)} R^{1/2}(x) U(b, x) (I + R(x) U(b, x))^{-1} \times \\ \times R^{-1/2}(x) \sqrt{R(x) - R^{-1}(x)}$$

in the space H_1 . Here R and U are related to the characteristic function w of the operator A by relations (4)–(6).

Odessa Electrotechnical
Institute of Communications

Received
21 VI 1965

REFERENCES

- ¹ M. S. Livshits, Mat. sborn., **34(76)**, 1, 175 (1954).
- ² L. A. Sakhnovich, UMN, **12**, issue 3, 205 (1957).
- ³ L. A. Sakhnovich, Ukr. matem. zhurn., vol. 11, issue 3, 275 (1959).
- ⁴ L. A. Sakhnovich, Mat. sborn., **44(86)**, 4, 509 (1958).
- ⁵ L. A. Sakhnovich, DAN, **115**, No. 3, 462 (1957).
- ⁶ L. A. Sakhnovich, On reducing non-self-adjoint operators to diagonal form, Candidate Dissertation, Odessa State Pedagogical Institute named after K. D. Ushinsky, 1956.
- ⁷ M. Rosenblum, Pacif. J. Math., **7**, No. 1, 997 (1957).
- ⁸ T. Kato, Proc. Japan. Acad., **33**, No. 5, 260 (1957).
- ⁹ I. V. Stankevich, DAN, **160**, No. 6 (1965).
- ¹⁰ B. Sz.-Nagy, C. Foias, Acta Sci. Math., **26**, 79 (1965).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.