

CLASSIFICATION OF INDECOMPOSABLE INFINITESIMAL REPRESENTATIONS OF THE LORENTZ GROUP

I. M. Gel'fand, V. A. Ponomarev

1967

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196701.33649>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 519.46

MATHEMATICS

Corresponding Member of the Academy of Sciences of the USSR I. M. GELFAND, V. A. PONOMAREV

CLASSIFICATION OF INDECOMPOSABLE INFINITESIMAL REPRESENTATIONS OF THE LORENTZ GROUP

Let L be the Lie algebra of the proper Lorentz group, and let L_k be its subalgebra corresponding to the maximal compact subgroup, which is isomorphic to the rotation group of three-dimensional space. Recall ⁽²⁾ that a module M is called a **Harish-Chandra module** over the Lie algebra L if, regarded as a module over L_k , it is a direct sum of finite-dimensional submodules: $M = \oplus M_i$. Here each M_i is an irreducible module over L_k , and for every M_{i_0} in the sequence $\{M_i\}$ there exist no more than a finite number equivalent to it.

A module M over the Lie algebra L is defined by specifying six operators $H_+, H_-, H_3, F_+, F_-, F_3$, which satisfy the known commutation relations ⁽¹⁾. By Δ_1 and Δ_2 we shall denote the Laplace operators. They are expressed in terms of the mappings H and F as follows:

$$\begin{aligned}\Delta_1 &= \frac{1}{2}(H_-F_+ + F_-H_+) + H_3F_3 + F_3; \\ \Delta_2 &= H_-H_+ - F_-F_+ + H_3^2 - F_3^2 + 2H_3.\end{aligned}\quad (1)$$

In the paper the explicit form of the operators H and F acting in the space M of an indecomposable module will be written out.

1. The operators H_+, H_-, H_3 determine in the representation the basis of the subalgebra L_k . It is known that each irreducible representation over L_k is specified by one positive number l , integer or half-integer. In accordance with this, in the decomposition $M = \oplus M_i$ we combine those M_i which correspond to representations with the same l . As a result we obtain a decomposition $M = \oplus R_l$ such that the representation in R_l of the subalgebra L_k is a multiple of an irreducible one. This means that R_l is a finite-dimensional subspace invariant with respect to the operators H_+, H_-, H_3 .

By $R_{l,m}$ ($R_{l,m} \subset R_l$) we denote the subspace of eigenvectors of the operator H_3 corresponding to the eigenvalue m :

$$(H_3)_{l,m} = mI_{l,m} \quad (m = -l, -l+1, \dots, l-1, l), \quad (2)$$

where $I_{l,m}$ is the identity operator, and the index l, m by the operator H_3 means that this operator is considered with domain $R_{l,m}$.

Using the explicit form of the representation of the algebra L_k ⁽¹⁾, one can show that all subspaces $R_{l,m}$ with one and the same l have the same dimension and $R_l = \oplus R_{l,m}$. In this case one can introduce operators E_+ and E_- with the following properties:

$$(E_+)_{l,m} : R_{l,m} \rightarrow R_{l,m+1}; \quad (E_-)_{l,m} : R_{l,m} \rightarrow R_{l,m-1}, \quad (3)$$

assuming that $R_{l,l+1} = R_{l,-l-1} = 0$ and

$$(E_-E_+)_{l,m} = I_{l,m} \quad (m < l); \quad (E_+E_-)_{l,m} = I_{l,m} \quad (m > -l). \quad (4)$$

The operators H_+ and H_- are expressed through E_+ and E_- by the formulas:

$$\begin{aligned} (H_+)_{l,m} &= \sqrt{(l+m+1)(l-m)} (E_+)_{l,m}; \\ (H_-)_{l,m} &= \sqrt{(l+m)(l-m+1)} (E_-)_{l,m}. \end{aligned} \quad (5)$$

Relations (3) and (4) show that the mappings E_+ and E_- define compatible isomorphisms between the spaces $R_{l,m}$ and $R_{l,m+1}$.

2. Operators F_+, F_-, F_3 . In order to write expressions for these operators, we introduce the mappings D_+, D_-, Δ

$$(D_+)_{l,m} : R_{l,m} \rightarrow R_{l+1,m}; \quad (D_-)_{l,m} : R_{l,m} \rightarrow R_{l-1,m}; \quad (6)$$

$$(\Delta)_{l,m} : R_{l,m} \rightarrow R_{l,m}.$$

We shall write out these operators later. Now, with their help, we write expressions for the operators F :

$$(F_3)_{l,m} = \sqrt{l^2 - m^2} (D_-)_{l,m} - md_l (\Delta)_{l,m} - \sqrt{(l+1)^2 - m^2} (D_+)_{l,m}; \quad (7)$$

$$\begin{aligned} (F_+)_{l,m} &= \sqrt{(l-m)(l-m-1)} (D_-)_{l,m+1} (E_+)_{l,m} \\ &\quad - d_l \sqrt{(l-m)(l+m+1)} (E_+)_{l,m} (\Delta)_{l,m} \\ &\quad + \sqrt{(l+m+1)(l+m+2)} (E_+)_{l+1,m} (D_+)_{l,m}; \end{aligned} \quad (8)$$

$$\begin{aligned}
(F_-)_{l,m} = & -\sqrt{(l+m)(l+m-1)}(D_-)_{l,m-1}(E_-)_{l,m} \\
& -d_l\sqrt{(l+m)(l-m+1)}(E_-)_{l,m}(\Delta)_{l,m} \\
& -\sqrt{(l-m+1)(l-m+2)}(E_-)_{l+1,m}(D_+)_{l,m},
\end{aligned} \tag{9}$$

where $d_l = -l^{-1}(l+1)^{-1}$.

We shall require of the operators D_+, D_-, Δ that the diagrams be commutative

$$\begin{array}{ccccccc}
R_{l-1,m+1} & \xleftarrow{D_-} & R_{l,m-1} & R_{l,m+1} & \xleftarrow{\Delta} & R_{l,m+1} & \xrightarrow{D_+} & R_{l+1,m+1} & \uparrow E_+ \\
\uparrow E_+ & & \uparrow E_+ & & \uparrow E_+ & & \uparrow E_+ & & \\
R_{l-1,m} & \xleftarrow{D_-} & R_{l,m} & R_{l,m} & \xleftarrow{\Delta} & R_{l,m} & \xrightarrow{D_+} & R_{l+1,m} &
\end{array} \tag{10}$$

$$-l+1 \leq m < l-1 \qquad -l \leq m < l.$$

In addition, the same diagrams with the mapping E_- instead of E_+ must also be commutative. Note that, by definition, the mapping $(E_+)_{l,m}$ is an isomorphism (for $m < l$). Consequently, the commutativity of diagrams (10) means that the mappings D_+, D_-, Δ do not depend on the index m . Therefore, in what follows we shall write $(D_+)_l$ instead of $(D_+)_{l,m}$, etc., wherever this cannot cause misunderstanding.

Next we shall require that the operators D_+, D_-, Δ satisfy the relations

$$(D_-)_l(\Delta)_l = (\Delta)_{l-1}(D_-)_l; \tag{11}$$

$$(D_+)_l(\Delta)_l = (\Delta)_{l+1}(D_+)_l; \tag{12}$$

$$(2l-1)(D_+)_{l-1}(D_-)_l - (2l+3)(D_-)_{l+1}(D_+)_l = I + l^{-2}(l+1)^{-2}(\Delta)_l^2. \tag{13}$$

Let us note that of all the relations which the operators D_+, D_-, Δ must satisfy, essentially all except (13) are trivial.

Assertion 1. *Let M be a Harish-Chandra module over the Lie algebra L of the proper Lorentz group. Then the space M is representable in the form of a direct sum of subspaces $R_{l,m}$. Moreover, one can choose mappings $E_+, E_-, D_+, D_-, \Delta$ such that the operators H and F are expressed through them by formulas (5), (7)–(9). Such mappings D, E, Δ must satisfy relations (3)–(4), (10)–(13).*

The converse is also true.

Assertion 2. *If in the space $M = \bigoplus_{l,m} R_{l,m}$ ($m = -l, -l+1, \dots, l$) operators $D_+, D_-, \Delta, E_+, E_-$ are given which satisfy relations (3), (4), (10)–(13), and through these operators the mappings $H_+, H_-, H_3, F_+, F_-, F_3$ are expressed by formulas (2), (5), (7)–(9), then the operators H and F define a representation of the Lie algebra L of the proper Lorentz group.*

In paper ⁽²⁾ it was shown that in an indecomposable module the Laplace operators Δ_1 and Δ_2 have one eigenvalue each, λ_1 and λ_2 . Moreover, for each indecomposable module one can specify a number l_0 , integer or half-integer ($l_0 \geq 0$), such that the numbers $\lambda_1, \lambda_2, l_0$ are related by

$$l_0^4 + (1 + \lambda_2)l_0^2 - \lambda_1^2 = 0. \quad (14)$$

The pair (λ_1, λ_2) is called special if one can find a real number l_1 satisfying the identities

$$l_1^2 l_0^2 = -\lambda_1^2; \quad l_1^2 + l_0^2 = -1 - \lambda_2 \quad (15)$$

and such that the difference $(|l_1| - l_0)$ is a positive integer. A module M with such a pair (λ_1, λ_2) will be called special. Otherwise we shall say that the pair (λ_1, λ_2) and the module M with such a pair are nonspecial.

3. The operators D_+, D_-, Δ in a nonspecial indecomposable module.

In paper ⁽²⁾ it was shown that to every indecomposable nonspecial module there corresponds, and moreover uniquely, a finite-dimensional space P with a linear nilpotent mapping A , where the matrix $[A]$ in some basis has the form of a single Jordan cell.

Theorem 3. *Let M be a nonspecial indecomposable Harish-Chandra module. Then the spaces $R_{l,m}$ ($l_0 \leq l$) have the same dimension, equal to $\dim P$. At the same time the mappings $(D_-)_{l,m}$ ($l \neq l_0$) and $(D_+)_{l,m}$ are isomorphisms. In the spaces $R_{l,m}$ one can choose a basis such that the matrices of the mappings D_+, D_-, Δ have the form*

$$\begin{aligned} [D_+]_{l,m} &= [I]; & [\Delta]_{l,m} &= [A] + \lambda_1 [I]; \\ [D_-]_{l,m} &= (l^2 - l_0^2)(4l^2 - 1)^{-1}((1 + \lambda_1 l^{-2} l_0^{-2})[I] + \\ &\quad + l^{-2} l_0^{-2}(2[A] + [A]^2)), \end{aligned} \quad (16)$$

where $[I]$ is the identity matrix, and the matrix $[A]$ has the form of a Jordan cell with zeros on the diagonal.

4. The operators D_+, D_-, Δ in a special indecomposable module.

In paper ⁽²⁾ it was shown that in this case the situation is radically different from that just described. A special indecomposable module, in comparison

with an irreducible one, is described by a set of integers, and this set may be arbitrarily long; moreover, in some cases, also by an additional complex number μ . It was shown there that to indecomposable special modules there corresponds, and moreover bijectively, a pair of finite-dimensional spaces P_1 and P_2 with mappings $d_+ : P_1 \rightarrow P_2$; $d_- : P_2 \rightarrow P_1$; $\delta : P_2 \rightarrow P_2$ such that the mappings d_+d_- and δ are nilpotent and $d_-\delta = \delta d_+ = 0$. Knowing the matrices of the mappings d_+, d_-, δ , one can find the matrices of the mappings D_+, D_-, Δ in the module M . It turns out that in a special indecomposable module M the subspaces $R_{l,m}$ ($l_0 \leq l \leq l_1 - 1$) have the same dimension, equal to $\dim P_1$, and the subspaces $R_{l,m}$ ($l_1 \leq l$) also have the same dimension, equal to $\dim P_2$. At the same time, in the subspaces one can choose a basis such that the matrices of the mappings D_+, D_-, Δ have the following form:

$$[D_+]_{l,m} = [I] \quad (l \neq l_1 - 1); \quad [D_+]_{l_1-1,m} = [d_+]; \quad (17)$$

$$[\Delta]_{l,m} = -l_1 l_0 ((4l_1^2 - 1)(l_1^2 - l_0^2)^{-1} [d_-][d_+] + [I])^{1/2} \quad (l \leq l_1 - 1);$$

$$[\Delta]_{l,m} = -il_1 l_0 ((4l_1^2 - 1)(l_1^2 - l_0^2)^{-1} [d_+][d_-] + [\delta] + [I])^{1/2} \quad (l \geq l_1); \quad (18)$$

$$[D_-]_{l,m} = (l^2 - l_0^2)l^{-2}(4l^2 - 1)^{-1}(l_1^2(4l_1^2 - 1)(l_1^2 - l_0^2)^{-1}[d_-][d_+] + (l_1^2 - l^2)[I]) \quad (l_0 \leq l \leq l_1 - 1); \quad (19)$$

$$[D_-]_{l_1,m} = [d_-];$$

$$[D_-]_{l,m} = (l^2 - l_0^2)l^{-2}(4l^2 - 1)^{-1}(l_1^2(4l_1^2 - 1)(l_1^2 - l_0^2)^{-1}[d_+][d_-] + (l_1^2 - l^2)[I] + l_0^2(l^2 - l_1^2)(l^2 - l_0^2)[\delta]) \quad (l > l_1),$$

where $[I]$ is the identity matrix.

Assertion 4. Let M be an indecomposable special Harish-Chandra module over the Lie algebra L . Then the spaces $R_{l,m}$ ($l_0 \leq l \leq l_1 - 1$) have the same dimension, and the spaces $R_{l,m}$ ($l \geq l_1$) also have the same dimension. Moreover, the mappings $(D_+)_{l,m}$ ($l \neq l_1 - 1$) and $(D_-)_{l,m}$ ($l \neq l_0$; $l \neq l_1$) are isomorphisms. The matrices of all mappings D_+, D_-, Δ are expressed in terms of the matrices $[d_+]$, $[d_-]$, $[\delta]$ by formulas (17)–(19). The matrices $[d_+]$, $[d_-]$, $[\delta]$ correspond to operators on the spaces P_1 and P_2 :

$$P_1 \xrightarrow{d_+} P_2 \xrightarrow{\delta} P_2 \xrightarrow{d_-} P_1$$

so that the mappings $(d_+ d_-)$ and (δ) are nilpotent, and $\delta d_+ = d_- \delta = 0$.

The canonical form of an indecomposable object which is a pair of spaces with mappings d_+ , d_- , δ , is described in paper ⁽²⁾. It is of two types. The first type is called an open object, and its invariants are a set of positive integers $[s, n_1, m_1; n_2, m_2; \dots; n_k, m_k]$, where $s = 0$ or 1 ; $n_1 \geq 0$; $n_i > 0$; $m_i > 0$; $m_k \geq -1$. The second type is called a closed object, and its invariants are a set of numbers $[n_1, m_1; n_2, m_2; \dots; n_k, m_k; \mu; N]$, where n_i , m_i , N are positive integers, and μ is an arbitrary complex number. We shall not write out the canonical form of the matrices of the mappings d_+ , d_- , δ . We note only that, in the case when these matrices correspond to an open chain, they consist of zeros and ones. If, however, the matrices correspond to a closed chain, then $[d_+]$ and $[d_-]$ have approximately the same form as in the preceding case, while in the matrix $[\delta]$ the number μ occurs in certain positions, and in the remaining positions there are zeros and ones.

Let us describe what the matrices $[d_+]$, $[d_-]$, $[\delta]$ look like in the simplest closed object, to which the set of numbers $[1, 1; \mu; 1]$ corresponds. In this case the space P_1 has dimension 1, and the space P_2 has dimension 2. In them one can choose a basis l_2 and f_1, f_2 such that

$$d_- f_1 = l_1; \quad d_- f_2 = 0; \quad \delta f_1 = \mu f_2; \quad \delta f_2 = 0; \quad d_+ l_1 = f_2.$$

Thus the matrices $[d_+]$, $[d_-]$, $[\delta]$; $[d_-][d_+]$; $[d_+][d_-]$ have the form

$$[d_+] = \begin{pmatrix} 0 \\ 1 \end{pmatrix}; \quad [d_-] = (1 \ 0); \quad [\delta] = \begin{pmatrix} 0 & 0 \\ \mu & 0 \end{pmatrix}; \quad [d_-][d_+] = (0); \quad [d_+][d_-] = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Received
6 VI 1967

CITED LITERATURE

¹ I. M. Gel'fand, R. A. Minlos, Z. Ya. Shapiro, *Representations of the Rotation Group and the Lorentz Group*, Moscow, 1958. ² I. M. Gel'fand, V. A. Ponomarev, DAN, **176**, No. 2 (1967).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.