

ON THE EXTENSION OF A. N. KOLMOGOROV' S UNIFORM THEOREM TO HOMOGENEOUS MARKOV CHAINS

D. V. Manevich

1967

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196701.43306>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 519.217

MATHEMATICS

D. V. MANEVICH

ON THE EXTENSION OF A. N. KOLMOGOROV'S UNIFORM THEOREM TO HOMOGENEOUS MARKOV CHAINS

(Presented by Academician A. N. Kolmogorov, 9 XII 1966)

In the works ^(1,2) A. N. Kolmogorov gave an estimate for the approximation of the distributions of sums of independent, identically distributed summands by infinitely divisible distributions. In the present note this result is extended to the case of distributions of sums of summands connected in a homogeneous Markov chain.

Let, on a probability space $\{\Omega, \mathcal{F}, P\}$, a sequence of random variables $\{X_k = X_k(\omega)\}$ be given, connected in a homogeneous Markov chain; Ω is an arbitrary set of states ω ; $\mathcal{F}$ is a σ -algebra of its subsets A ; P is a probability measure of the elements of the set $\mathcal{F}$. Denote by $P_{ij}(\omega, A)$ the probabilistic transition function in $j - i$ steps;

$P(A)$ is the probability of A ; $c, c_1, c_2, \dots$ are absolute constants; $s_n = \sum_{i=1}^n X_i$,

$$F_n = P(s_n < x), \quad F = P(X_i < x).$$

We shall assume that the ergodic property of the chain is characterized by the condition adopted in the works of S. V. Nagaev ⁽³⁾:

$$\sup_{\omega, A} |P_{ij}(\omega, A) - P(A)| \leq Cq^\mu, \quad 0 < q < 1, \quad \mu = j - i.$$

We note that the ergodicity condition used in the works of R. L. Dobrushin ⁽⁴⁾ would not change the subsequent reasoning.

Theorem. Let G be the set of all infinitely divisible distribution laws $D(x)$. Then there exists an absolute constant C_1 such that

$$\rho(F_n, G) = \sup_F \inf_{D_n \in G} \sup_x |F_n - D_n| \leq c_1 n^{-1/3} \ln^{\lambda/3} n, \quad (1)$$

where $\lambda > 1$.

We outline the proof. Using the idea of S. N. Bernstein ⁽⁵⁾, we construct sums of independent random variables by whose distributions we shall approximate F_n . Write s_n as follows:

$$s_n = y_1 + u_1 + \dots + y_l + u_l,$$

where

$$y_i = X_{(i-1)(r+k)+1} + \dots + X_{i(r+k)-k}; \quad u_i = X_{i(r+k)-k+1} + \dots + X_{i(r+k)};$$

$$i = 1, \dots, l; \quad (r+k)l \geq n; \quad \Sigma_l = y_1 + \dots + y_l; \quad \sigma_l = u_1 + \dots + u_l;$$

$$s_n = \Sigma_l + \sigma_l.$$

Put $k = r = \ln^\lambda n$, $\lambda > 1$. Then $l \geq n/2 \ln^\lambda n$.

Denote

$$F_\Sigma(x) = P(\Sigma_l < x), \quad F_\sigma(x) = P(\sigma_l < x),$$

and $F_\Sigma(x/A)$, $F_\sigma(x/A)$ the corresponding conditional distribution functions (d.f.) under some condition A ; Φ'_k is some conditional d.f., and Φ_k the unconditional d.f. of y_k .

The rest of the proof rests on three lemmas.

Lemma 1. Let G_l and f be the characteristic functions (c.f.) of Σ_l and y_k , respectively. Then

$$G_l - f^l = \sum_{r=1}^l \delta_r L_r, \quad \text{where} \quad \delta_r = \int e^{itx} d[\Phi'_r - \Phi_r], \quad L_r \equiv G_{r-1} f^{l-r}.$$

Proof is carried out in many respects in the same way as that of Lemma 2 in the author's paper ⁽⁶⁾.

Lemma 2. One can indicate such a continuous d.f. $\Psi_\Sigma(x)$ of the sum of l terms, related to one another in the same way as the y_k in the sum Σ_l , that

$$|F_{\Sigma}(x) - \Psi_{\Sigma}(x)| \leq c_2 n^{-1/3}.$$

For the proof we compose $\Phi(x/\varepsilon)$ —a certain normal distribution—with $F_{\Sigma}(x)$. We show that

$$|F_{\Sigma}(x) - \Psi_{\Sigma}(x)| \leq |\Psi_{\Sigma}(x - 2h\varepsilon) - \Psi_{\Sigma}(x + 2h\varepsilon)| + \frac{2c_3}{h} e^{-h^2/2}, \quad \Psi_{\Sigma} = F_{\Sigma} * \Phi.$$

The estimate

$$\alpha \equiv |\Psi_{\Sigma}(x - 2h\varepsilon) - \Psi_{\Sigma}(x + 2h\varepsilon)|$$

is carried out by S. N. Bernstein's lemma ((7), p.409):

$$\alpha \leq \frac{2\varepsilon}{\pi} \left[A \cdot 2h\varepsilon + \frac{1}{2\sqrt{Nh\varepsilon}} \right],$$

$$A = \frac{1}{2} \left| \int_{-N}^N \varphi_l(t) dt \right|;$$

φ_l is the ch.f. for the d.f. $\Psi_{\Sigma}(x)$. Then, by Lemma 1,

$$A = \frac{1}{2} \left| \int_{-N}^N f^l(t) dt + \sum_{k=1}^l \int_{-N}^N \delta_k L_k dt \right|.$$

The first integral is estimated by known methods, while the second is reduced to the Dirichlet integral. Setting $h = \ln^2 n$, $\varepsilon = \ln^{-2} n \cdot n^{-1/3}$, $\sqrt[3]{n} = n^{2/3}$, and using the ergodicity condition, we verify the validity of the lemma.

Lemma 3. Let $\bar{F}_{\Sigma}(x)$ denote the d.f. of the sum $y_1 + \dots + y_l$ under the assumption of continuity and independence of the terms. Then

$$|\bar{F}_{\Sigma}(x) - \Psi_{\Sigma}(x)| \leq c_4 n^{-1/3}.$$

Proof. For the d.f.'s $\bar{F}_{\Sigma}(x)$ and $\Psi_{\Sigma}(x)$, the corresponding ch.f.'s can be written as follows: $f^l \varphi$; $G_l \varphi$, where φ is the ch.f. for $\Phi(x/\varepsilon)$. Next we use Esseen's theorem (8), for the application of which we consider

$$\Delta = \left| \int_{-T}^T e^{-itx} \frac{(f^l - G_l)\varphi}{-it} h(t) dt \right|$$

instead of

$$\int_{-T}^T \frac{|f^l - G_l| |\varphi|}{|t|} dt,$$

and verify the validity of the lemma.

Further,

$$F_n(x) = M \left\{ \int F_\Sigma \left(\frac{x-t}{A} \right) dF_\sigma \left(\frac{t}{A} \right) \right\}.$$

Since Lemma 2 remains valid also for conditional distributions, i.e.

$$\left| F_\Sigma \left(\frac{x-t}{A} \right) - \Psi_\Sigma \left(\frac{x}{A} \right) \right| \leq c_2 n^{-1/3},$$

the error in replacing $F_\Sigma((x-t)/A)$ by a continuous function will not exceed $c_2 n^{-1/3}$. We shall assume, without making elementary explanations for simplicity, that $F_\Sigma((x-t)/A)$ is continuous. From

$$F_n(x) = M \left[\int_{-\infty}^{\xi} + \int_{\xi}^{\infty} \right]$$

we obtain

$$F_\sigma(\xi) - 1 \leq F_n(x) - F_\Sigma(x-\theta) \leq 0,$$

$-\infty < \theta \leq \xi$. We choose ξ so that $1 - F_\sigma(\xi) \leq n^{-1/3}$. Next we replace $F_\Sigma(x-\theta)$ by the distribution $\bar{F}_\Sigma(x-\theta)$. Then, using Lemma 3, we obtain

$$-c_3 n^{-1/3} \leq F_n(x) - \bar{F}_\Sigma(x-\theta) \leq c_5 n^{-1/3}. \quad (2)$$

According to A. N. Kolmogorov's theorem ⁽²⁾, there exists an infinitely divisible distribution $D_n(x)$ such that

$$|\bar{F}_\Sigma(x-\theta) - D_n(x)| \leq c_6 n^{-1/3} \ln^\lambda n,$$

since the number of terms in Σ_l is equal to $l \geq n/2 \ln^\lambda n$. Combining this with (2), we arrive at the conclusion that (1) holds. The theorem is proved.

Tashkent State Pedagogical Institute
named after Nizami

Received
25 XI 1966

References

1. A. N. Kolmogorov, *Theory of Probability and Its Applications*, 1, § 4, 426 (1956).
2. A. N. Kolmogorov, *Trudy Moskov. matem. obshch.*, 12, 437 (1963).
3. S. V. Nagaev, *Theory of Probability and Its Applications*, 2, No. 4, 389 (1957).
4. R. L. Dobrushin, *Theory of Probability and Its Applications*, 1, No. 1, 72 (1956).
5. S. N. Bernstein, *UMN*, 10 (1944).
6. D. V. Manevich, *Collected Papers: Probability Theory and Mathematical Statistics*, vol. 1, Academy of Sciences of the Uzbek SSR, 1964.
7. S. N. Bernstein, *Probability Theory*, Moscow, 1946.
8. C. G. Esseen, *Acta Math.*, 77, 1 (1945).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.