

APPLICATION OF HAAR SERIES TO ERROR ESTIMATION IN THE COMPUTATION OF INFINITE- DIMENSIONAL INTEGRALS

I. M. Sobol'

1967

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196701.88340>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 518.12+517.392

MATHEMATICS

I. M. SOBOL'

APPLICATION OF HAAR SERIES TO ERROR ESTIMATION IN THE COMPUTATION OF INFINITE-DIMENSIONAL INTEGRALS

(Presented by Academician A. N. Tikhonov on 24 IX 1966)

We estimate the error of the approximation

$$\int_{K_\infty} f(X) dX \approx \frac{1}{N} \sum_{\nu=1}^N f(X_\nu), \quad (*)$$

where K_∞ is the infinite-dimensional unit cube; $X = (x_1, x_2, \dots, x_n, \dots)$ is a point of this cube. The measure on K_∞ is defined by means of the infinite product of Lebesgue measures ($dX = dx_1 dx_2 \dots dx_n \dots$).

In (1) a sequence of points $\{X_\nu^*\}$ was constructed, called a generalized Halton sequence, whose points can be used as nodes in (*), and an error estimate for such an approximation was obtained for certain classes of functions.

In § 2 of the present paper the error of the approximation (*) is estimated for broader classes of functions S_p and H_α (the classes H_α are analogues of the Lipschitz classes $\text{Lip } \alpha$, $0 < \alpha \leq 1$). A new sequence of points $\{X_\nu^{**}\}$, called a generalized τ -sequence, is constructed, for which estimates are obtained that are somewhat better than for $\{X_\nu^*\}$. On the classes S_p the order of these estimates turns out to be the best possible. The concept of a grid $\{X_1, \dots, X_N\}$ with a multiplicative estimate of nonuniformity is introduced. For such grids an estimate of the approximation (*) is obtained in general form.

The paper uses the method of Haar series, which made it possible to obtain error estimates on the classes S_p and H_α in the n -dimensional case (2^{-4}). However, in order to be able to pass to the limit as $n \rightarrow \infty$, it was necessary to improve these estimates somewhat. This is done in § 1.

In what follows, the symbol $K_{i_1 \dots i_s}$ denotes the s -dimensional face of the cube K_∞ on which $x_{i_1}, \dots, x_{i_s}$ range from 0 to 1, while all the remaining $x_i \equiv 1$.

§ 1. The case of n variables.

1.1. Expansion of a function in “different-dimensional” terms.

Consider a function $f(P)$, where $P = (x_1, \dots, x_n)$, defined in the n -dimensional unit cube $K \equiv K_{12\dots n}$. Write its expansion in the Fourier-Haar series, explicitly separating the first function of the Haar system $\chi_1(x) \equiv 1$. Let

$$C_{k_1\dots k_s}^{i_1\dots i_s} = \int_K f(P) \chi_{k_1}(x_{i_1}) \cdots \chi_{k_s}(x_{i_s}) dP,$$

$$(dP = dx_1 \cdots dx_n),$$

where $1 \leq i_1 < i_2 < \dots < i_s \leq n$, $1 \leq s \leq n$. Then

$$f(P) = c_0 + \sum_{k_1, \dots, k_s}^{\wedge} C_{k_1\dots k_s}^{i_1\dots i_s} \chi_{k_1}(x_{i_1}) \cdots \chi_{k_s}(x_{i_s}), \quad (1)$$

where the symbol $\sum^{\wedge}$ denotes summation over all different s -index quantities for $s = 1, 2, \dots, n$:

$$\sum^{\wedge} T_{i_1\dots i_s} = \sum_{i_1=1}^n T_{i_1} + \sum_{1 \leq i_1 < i_2 \leq n} T_{i_1 i_2} + \dots + T_{12\dots n}.$$

In (1), as throughout what follows, the indices k_σ vary from 2 to ∞ . Each of the sums following the sign $\sum^{\wedge}$ depends only on the s variables $x_{i_1}, \dots, x_{i_s}$.

1.2. Estimation of the integration error. Choose an integration net Σ , consisting of arbitrary points $P_1, \dots, P_N$ of the cube K , and estimate the error

$$\delta(f; \Sigma) = \frac{1}{N} \sum_{\nu=1}^N f(P_\nu) - \int_K f(P) dP. \quad (2)$$

Suppose that the series (1) converges uniformly, and substitute it into (2). After transformations entirely analogous to the transformations of Sec. 1 from ⁽³⁾, we obtain the estimate

$$|\delta(f; \Sigma)| \leq \frac{1}{N} \sum^{\wedge} A_p^{i_1\dots i_s}(f) \Phi_q^{i_1\dots i_s}(\Sigma), \quad (3)$$

in which $1/p + 1/q = 1$;

$$A_p^{i_1\dots i_s}(f) = \sum_m 2^{(m_1-1)/2 + \dots + (m_s-1)/2} \left\{ \sum_j |C_k^i|^p \right\}^{1/p},$$

and $\Phi_q^{i_1 \dots i_s}(\Sigma)$ is the s -dimensional discrepancy of the projections of the points $P_1, \dots, P_N$ onto $K_{i_1 \dots i_s}$ ⁽³⁾. Here $i = (i_1, \dots, i_s)$, $k = (k_1, \dots, k_s)$, $m = (m_1, \dots, m_s)$, $j = (j_1, \dots, j_s)$; $1 \leq m_\sigma < \infty$, $1 \leq j_\sigma \leq 2^{m_\sigma - 1}$.

In ⁽³⁾ the norm $\|f\|_p = \sum \hat{A}_p^{i_1 \dots i_s}(f)$ was used. Therefore the estimate (3) was coarsened and written in the form

$$|\delta(f; \Sigma)| \leq N^{-1} \|f\|_p \varphi_q(\Sigma), \quad \text{where } \varphi_q(\Sigma) = \max \Phi_q^{i_1 \dots i_s}(\Sigma).$$

Below only the simplest quantities $\Phi_\infty^{i_1 \dots i_s}(\Sigma)$ are used; we shall call them the discrepancies of the net Σ . It is easy to prove that for any net

$$\Phi_q^{i_1 \dots i_s}(\Sigma) \leq N^{1/q} \Phi_\infty^{i_1 \dots i_s}(\Sigma). \quad (4)$$

1.3. Classes of functions S_p and H_α .

Definition. A function $f(x_1, \dots, x_n)$ belongs to the class $S_p(L_{i_1 \dots i_s})$, $1 \leq p < \infty$, if for any $1 \leq i_1 < i_2 < \dots < i_s \leq n$ and $1 \leq s \leq n$

$$A_p^{i_1 \dots i_s}(f) \leq L_{i_1 \dots i_s}. \quad (5)$$

The constants $\{L_{i_1 \dots i_s}\}$ are called the defining constants of the class.

As in ^(2, 3), it is proved that if $f \in S_p$, then the series (1) converges uniformly in K , and estimate (3) with $L_{i_1 \dots i_s}$ in place of $A_p^{i_1 \dots i_s}(f)$ is an exact estimate on S_p (for any prescribed net Σ).

Definition. A function $f(x_1, \dots, x_n)$ belongs to the class $H_\alpha(L_{i_1 \dots i_s})$, $0 < \alpha \leq 1$, if for any $1 \leq i_1 < i_2 < \dots < i_s \leq n$ and $1 \leq$

$\leq s \leq n$ in the cube K

$$\left| \Delta_{\xi_{i_1}} \dots \Delta_{\xi_{i_s}} f(P) \right| \leq L_{i_1 \dots i_s} \left(\frac{\alpha + 1}{2} \right)^s |\xi_{i_1} \dots \xi_{i_s}|^\alpha. \quad (6)$$

Just as in ⁽²⁾, it is proved that if $f \in H_\alpha(L_{i_1 \dots i_s})$ and $\alpha p > 1$, then

$$A_p^{i_1 \dots i_s}(f) \leq L_{i_1 \dots i_s} (2^{1+\alpha} - 2^{1+1/p})^{-s}. \quad (7)$$

From (3), (4), and (7) one can obtain an estimate of $|\delta(f; \Sigma)|$ on the classes H_α , quite analogous to estimate (8) from ⁽³⁾.

§ 2. The case of infinitely many variables

2.1. Classes of functions

The classes $S_p(L_{i_1 \dots i_s})$ and $H_\alpha(L_{i_1 \dots i_s})$ are defined in the same way as in the n -dimensional case: the corresponding inequalities (5) or (6) must be satisfied in K_∞ for arbitrary $1 \leq i_1 < i_2 < \dots < i_s$, $1 \leq s < \infty$. In addition, it is assumed that at each point $f(X) = \lim_{n \rightarrow \infty} f(x_1, \dots, x_n, 1, 1, 1, \dots)$, and passage to the limit is possible:

$$\int_{K_\infty} f(X) dX = \lim_{n \rightarrow \infty} \int_K f(x_1, \dots, x_n, 1, 1, \dots) dP.$$

2.2. Estimate of the error of integration

Choose a grid consisting of points $X_1, \dots, X_N$ in K_∞ . The projections of these points onto $K_{i_1 \dots i_s}$ form an s -dimensional grid, whose discrepancy shall be denoted by $\Phi_\infty^{i_1, \dots, i_s}$.

Suppose that there exist constants B and $\{h_i\}$, $1 \leq i < \infty$, such that for any $i_1 < \dots < i_s$

$$\Phi_\infty^{i_1 \dots i_s} \leq B h_{i_1} \dots h_{i_s}. \quad (8)$$

In this case we shall say that the grid admits a multiplicative estimate of discrepancies.

Theorem. For a grid $X_1, \dots, X_N$ admitting the multiplicative estimate of discrepancies (8), on classes of functions S_p and H_α whose defining constants decrease uniformly with respect to $\{h_i\}$, the estimate holds

$$\left| \frac{1}{N} \sum_{\nu=1}^N f(X_\nu) - \int_{K_\infty} f(X) dX \right| \leq AB N^{-1/p} \exp \sum_{i=1}^{\infty} v_i, \quad (9)$$

where $v_i = \varepsilon_i h_i$ in the case of the class S_p , while in the case of the class H_α the parameter $p > 1/\alpha$ and

$$v_i = \varepsilon_i h_i (2^{1+\alpha} - 2^{1+1/p})^{-1}.$$

Proof scheme. Use inequality (3) for the function $f(x_1, \dots, x_n, 1, 1, \dots)$, then inequalities (5) or (7), (4), and (8). The right-hand side is estimated by means of Lemma 2 from ⁽¹⁾. Then pass to the limit as $n \rightarrow \infty$.

2.3. Use of the generalized Halton sequence

As the grid $X_1, \dots, X_N$ we choose the points $X_1^*, \dots, X_N^*$. Since the projections of the points $\{X_\nu^*\}$ onto $K_{i_1 \dots i_s}$ form an s -dimensional Halton sequence, it is easy to show (cf. (1, 4)) that estimate (8) will be satisfied with $B = 1$, $h_i = \bar{\beta}_i \ln N + \bar{\gamma}_i$. The asymptotics of the coefficients as $i \rightarrow \infty$ is known: $\bar{\beta}_i \sim 4i$, $\bar{\gamma}_i \sim 8i \ln i$. Hence, for the applicability of the preceding theorem it is sufficient to require that the defining constants of the class decrease uniformly with respect to $\{i \ln i\}$.

Choose an arbitrary $\varepsilon > 0$. One can choose A and $\{\varepsilon_i\}$ so that $\sum \varepsilon_i \bar{\beta}_i = \varepsilon$; then the order of estimate (9) on the classes S_p will be $N^{-1/p+\varepsilon}$.

* According to (1), this means that $L_{i_1 \dots i_s} \leq A \varepsilon_{i_1} \dots \varepsilon_{i_s}$ and, in addition, the series $\sum_{i=1}^{\infty} \varepsilon_i h_i$ converges.

If p is fixed so that $0 < \alpha - 1/p < \varepsilon$, and then A and $\{\varepsilon_i\}$ are chosen so that $\sum \varepsilon_i \beta_i = (2^{1+\alpha} - 2^{1+1/p})(\varepsilon - \alpha + 1/p)$, then the order of the estimate (9) on the classes H_α will be $N^{-\alpha+\varepsilon}$.

We note that the orders of the estimates on the classes S_p and H_α , even in the n -dimensional case, cannot be better than $N^{-1/p}$ and, respectively, $N^{-\alpha}$.

2.4. Use of a generalized LP_τ -sequence

Definition. By a generalized LP_τ -sequence we shall mean a sequence of points $\{X_\nu^{**}\}$ with coordinates

$$X_{\nu+1}^{**} = (p^{(1)}(\nu), p^{(2)}(\nu), \dots, p^{(n)}(\nu), \dots),$$

where $\{p^{(n)}(\nu)\}$ is a sequence of type DR corresponding to the n -th monocyclic operator (5) (the operators are numbered so that their orders $\bar{m}_n$ do not decrease).

As the grid $X_1, \dots, X_N$ we choose the points $X_1^{**}, \dots, X_N^{**}$. Since the projections of the points $\{X_\nu^{**}\}$ onto $K_{i_1, \dots, i_s}$ form an s -dimensional LP_τ -sequence, it follows from (5) that estimate (8) holds with $B = 1/2$, $h_i = 2^{\bar{m}_i}$, and also the asymptotic estimate for $\bar{m}_i$ as $i \rightarrow \infty$,

$$\bar{m}_i \leq \log_2 i + \log_2 (\log_2 i \log_2 \log_2 i) + O(1). \quad (10)$$

Thus, $h_i \leq Ci \ln i \ln \ln i$, and for the applicability of our theorem it is sufficient to require that the defining constants of the class decrease uniformly with respect to $\{i \ln i \ln \ln i\}$.

The order of the estimate (9) on the classes S_p turns out to be $N^{-1/p}$. This is the best possible order.

Let us now consider the classes H_α . Let $0 < \varkappa < \varepsilon$. Fix $1/p = \alpha - \varkappa/\sqrt{\ln N}$. After a suitable choice of A and $\{\varepsilon_i\}$, the order of the estimate (9) will be equal to $N^{-\alpha+\varepsilon/\sqrt{\ln N}}$.

Remark. In this subsection we had to require a more rapid decrease of the defining constants in comparison with Sec. 2.3. Apparently this is caused by the crudeness of estimate (10).

Received
20 IX 1966

References

1. I. M. Sobol' , *Zhurn. vychislit. matem. i matem. fiz.*, **1**, No. 5, 917 (1961).
2. I. M. Sobol' , *DAN*, **132**, No. 4, 773 (1960).
3. I. M. Sobol' , *DAN*, **132**, No. 5, 1041 (1960).
4. I. M. Sobol' , *DAN*, **139**, No. 4, 821 (1961).
5. I. M. Sobol' , *UMN*, **21**, No. 5, 271 (1966).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.