

INTEGRAL INEQUALITIES WITH OPERATORS OF VOLTERRA TYPE

Kh. M. Salpagarov

1967

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196701.89854>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.43

MATHEMATICS

Kh. M. Salpagarov

INTEGRAL INEQUALITIES WITH OPERATORS OF VOLTERRA TYPE

(Presented by Academician A. A. Dorodnitsyn, February 1, 1967)

By E^+ we shall denote a linear partially ordered set of nonnegative elements $u, v, \psi, \dots$, and by K^+ the set of operators L, l, L_i ($i = 1, \dots, n$) of Volterra type that map the set E^+ into itself, for which the series

$$R_i(\lambda_i) = (I - \lambda_i L_i)^{-1} = \sum_{k=0}^{\infty} \lambda_i^k L_i^k$$

converge for every value of the complex parameter λ_i . We shall assume that the set K^+ , together with any two operators, contains their sum, product, and the operator $T_i(\lambda) = L_i R_i(\lambda)$, if $L_i \in K^+$ ($\lambda \geq 0$).

We shall be interested in the following properties of the operators. Multiplying the equality

$$L_i = (\lambda - \alpha)^{-1}[(I - \alpha L_i) - (I - \lambda L_i)] \quad (\lambda \neq \alpha)$$

by $R_i(\alpha)R_i(\lambda)$, we easily arrive at the relation

$$T_i(\alpha)R_i(\lambda) = (\lambda - \alpha)^{-1}[R_i(\lambda) - R_i(\alpha)],$$

whence, for $\lambda > \alpha \geq 0$, in turn follows the inequality

$$T_i(\alpha)R_i(\lambda) \leq (\lambda - \alpha)^{-1}[R_i(\lambda) - I] < (\lambda - \alpha)^{-1}R_i(\lambda).$$

Using this inequality, by the method of complete mathematical induction one can show that

$$T_i^m(\alpha)R_i(\lambda) < (\lambda - \alpha)^{-m}[R_i(\lambda) - I]. \quad (1)$$

Let now the operators L_i satisfy the relation $L_i L_k \leq L_k L_i$ for $i > k$.

Then, if $\alpha_i \geq 0$, $\alpha_k \geq 0$, $i > k$, then

$$T_i(\alpha_i)R_k(\alpha_k) \leq R_k(\alpha_k)T_i(\alpha_i). \quad (2)$$

Consider the operator $R = R_1(\lambda_1) \cdots R_n(\lambda_n)$. If $\lambda_i > \alpha_i \geq 0$, using inequalities (1) and (2), one can obtain that

$$T_i^m R \leq (\lambda_i - \alpha_i)^{-m} (R - I). \quad (3)$$

In what follows we shall assume that the nonnegative real numbers λ_i and α_{ij} ($i = 1, \dots, n$; $j = 1, \dots, p$) satisfy the inequality $\lambda_i > \alpha_{ij}$, and m_{ij} will denote integers.

For simplicity put $T_{ij} = T_i(\alpha_{ij})$. Starting from inequality (3), by the method of complete mathematical induction one can finally prove the inequality

$$\prod_{i=1}^n \prod_{j=1}^p T_{ij}^{m_{ij}} R \leq \prod_{i=1}^n \prod_{j=1}^p (\lambda_i - \alpha_{ij})^{-m_{ij}} (R - I), \quad (4)$$

Let us introduce into consideration the operator

$$L = \sum_{m=1}^N a(m_{11}, \dots, m_{ij}, \dots, m_{np}) \prod_{i=1}^n \prod_{j=1}^p T_{ij}^{m_{ij}},$$

where, respectively, $m = \sum_{i=1}^n \sum_{j=1}^p m_{ij}$ are natural numbers, and the coefficients a are nonnegative numbers. The equation

$$F(\lambda) \equiv \sum_{m=1}^N a(m_{11}, \dots, m_{np}) \prod_{i=1}^n \prod_{j=1}^p (\lambda_i - \alpha_{ij})^{-m_{ij}} = 1, \quad (5)$$

where $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$, will be called characteristic for the equation $u = \psi + Lu$. Obviously, $L \in K^+$.

Let us formulate the main inequality.

Theorem 1. If an element u satisfies the inequality $u \leq \psi + Lu$ ($u, \psi \in E^+$), then

$$u \leq R\psi = R_1(\lambda_1) \cdots R_n(\lambda_n)\psi \equiv v(\lambda), \quad (6)$$

where $\lambda_1, \dots, \lambda_n$ satisfy the characteristic equation (5).

To prove the assertion of the theorem, consider the expression $\psi + Lv$. From inequality (4), taking into account that λ satisfy relation (5), it is easy to obtain the inequalities

$$\psi + Lv = \psi + LR\psi \leq \psi + F(\lambda)(R - I)\psi = \psi + (R - I)\psi = R\psi = v,$$

i.e., $v \geq \psi + Lv$. Now the assertion of the theorem follows from the following proposition.

Lemma. If u, v satisfy the inequalities $u \leq \psi + Lu$, $v \geq \psi + Lv$, where $L \in K^+$, then $u \leq v$.

The assertion of the lemma is proved as follows: subtracting the first inequality from the second, we obtain $v - u \geq L(u - v)$. Hence $v - u = \psi_1 + L(v - u)$ ($\psi_1 \in E^+$), and $v - u = (I - L)^{-1}\psi_1 \geq 0$.

Remark 1. The characteristic equation (5) for $n > 1$ has an infinite set of solutions. To each solution there corresponds its own estimate of the form (6). Denote $v_* = \inf_{F(\lambda)=1} v(\lambda)$. It can be proved: if u satisfies the condition of Theorem 1, then $u \leq v_*$.

For operators of a special form one can also obtain the reverse inequality.

Theorem 2. If v satisfies the inequality $v \geq \psi + \sum_{k=1}^n a_k L_k v$, then

$$v \geq R_n(a_n) \cdots R_1(a_1)\psi.$$

Corollary. From Theorems 1 and 2 there follows the assertion: if the operator l is defined by the rule

$$lu = \sum_{k=1}^n a_k L_k u,$$

then

$$R_n(a_n) \cdots R_1(a_1) \leq (I - l)^{-1} \leq R_1(\lambda_1) \cdots R_n(\lambda_n),$$

where the nonnegative numbers $\lambda_1, \dots, \lambda_n$ satisfy the relation

$$\lambda_1^{-1}a_1 + \cdots + \lambda_n^{-1}a_n = 1. \quad (7)$$

Examples. If

$$L_k u = \int_0^{t_k} \alpha_k(s_k) u(t_1, \dots, t_{k-1}, \tau_k, t_{k+1}, \dots, t_n) d\tau_k, \quad (8)$$

then

$$T_k(\lambda_k)\psi = \int_0^{t_k} \alpha_k \exp \left\{ \lambda_k \int_{s_k}^{t_k} \alpha_k d\tau_k \right\} \psi(\dots, t_{k-1}, s_k, t_{k+1}, \dots) ds_k. \quad (9)$$

and $R_k(\lambda_k) = I + \lambda_k T_k(\lambda_k)$, where the operators L_k ($k = 1, \dots, n$) commute pairwise.

From Theorems 1 and 2 one can derive the following inequalities.

1. If the function $u(t)$ satisfies the inequality

$$u(t) \leq \psi(t) \int_0^t + \alpha u d\tau,$$

then

$$u(t) \leq \psi(t) + \int_0^t \alpha \exp \left\{ \int_s^t \alpha d\tau \right\} \psi d\tau.$$

This is the well-known Gronwall–Bellman inequality ⁽¹⁾.

2. If the function $u = u(t_1, \dots, t_n)$ satisfies the inequality

$$u \leq \psi + \sum_{k=1}^n L_k u,$$

then

$$u \leq \prod_{k=1}^n (I + \lambda_k T_k) \psi = \psi + \sum_{k=1}^n \lambda_k T_k \psi + \dots + \lambda_1 \dots \lambda_n T_1 \dots T_n \psi;$$

in particular, if $\psi = C > 0$ is constant, then

$$u \leq C \exp \left\{ \sum_{k=1}^n \lambda_k \int_0^{t_k} \alpha_k d\tau_k \right\}; \quad (10)$$

if

$$\psi = \int_0^{t_1} \dots \int_0^{t_n} \varphi d\tau_n \dots d\tau_1,$$

then

$$u \leq \int_0^{t_1} \dots \int_0^{t_n} \exp \left\{ \sum_{k=1}^n \lambda_k \int_{s_k}^{t_k} \alpha_k d\tau_k \right\} \varphi d\tau_n \dots d\tau_1, \quad (11)$$

where the operators $L_k, T_k = T_k(\lambda_k)$ are determined from relations (8) and (9), $\psi \geq 0$, $\lambda_1^{-1} + \dots + \lambda_n^{-1} = 1$.

These inequalities improve some inequalities of Wendroff ⁽¹⁾ and inequalities from ⁽²⁾.

Remark. By varying the numbers λ_k , instead of inequalities (10) and (11) one can obtain the sharper inequalities

$$u \leq C \exp \left\{ \sum_{k=1}^n \left[\int_0^{t_k} \alpha_k d\tau_k \right]^{1/2} \right\}^2,$$

$$u \leq \int_0^{t_1} \cdots \int_0^{t_n} \exp \left\{ \sum_{k=1}^n \left[\int_{s_k}^{t_k} \alpha_k d\tau_k \right]^{1/2} \right\}^2 \varphi ds_n \dots ds_1.$$

3. Let now

$$L_k u = \int_0^t e^{a_k(t-\tau)} u d\tau \quad (k = 1, \dots, n).$$

Then the operators L_k commute among themselves and

$$R_k(\lambda_k)\psi = \psi + \lambda_k \int_0^t e^{(\lambda_k + a_k)(t-\tau)} \psi(\tau) d\tau.$$

One can also find constants C_k ($k = 1, \dots, n$) such that

$$\prod_{k=1}^n R_k(\lambda_k)\psi \leq \psi + \sum_{k=1}^n C_k \int_0^t e^{(\lambda_k + a_k)(t-\tau)} \psi(\tau) d\tau. \quad (12)$$

Therefore, if the function $u(t)$ satisfies the inequality

$$u \leq \psi + \sum_{k=1}^n a_k \int_0^t e^{a_k(t-\tau)} u d\tau,$$

where $a_k \geq 0$, $\psi \geq 0$, then the function u will be less than the right-hand side of relation (12) for a corresponding choice of the constants C_k . ($\lambda_k > 0$ are determined from relation (7).)

4. From Theorems 1 and 2 one can obtain various inequalities also with matrix coefficients; for example, if

$$u(t) \leq \psi(t) + A \int_0^t e^{P(t-\tau)} u d\tau + B \int_0^t e^{Q(t-\tau)} u d\tau,$$

then

$$u(t) \leq \psi(t) \lambda A \int_0^t e^{(\lambda A + P)(t-\tau)} \psi(\tau) d\tau + \mu B \int_0^t e^{(\mu B + Q)(t-\tau)} \psi(\tau) d\tau$$

$$+ \lambda \mu A B \int_0^t e^{(\lambda A + P)(t-s)} \int_0^s e^{(\mu B + Q)(s-\tau)} \psi(\tau) d\tau,$$

where u, ψ are nonnegative vector functions; A, B, P, Q are square matrices with nonnegative components that are pairwise permutable with one another; $\lambda > 1$, $\mu > 1$ satisfy the equality $\lambda^{-1} + \mu^{-1} = 1$.

These inequalities can be used to prove theorems on existence, uniqueness, continuous dependence, and stability of solutions of differential and integral equations in many variables.

Karachay-Cherkess State Pedagogical Institute

Received
30 I 1967

REFERENCES

¹ E. Beckenbach, R. Bellman, *Inequalities*, Moscow, 1965. ² Kh. M. Salpagarov, Proceedings of the XII Scientific Conference of Kyrgyz University, Frunze, 1964.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.