

ON ZOLOTAREV STABILITY OF FUNCTIONALS

E. V. Voronovskaya

1967

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196701.94322>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.946

MATHEMATICS

E. V. VORONOVSKAYA

ON ZOLOTAREV STABILITY OF FUNCTIONALS

(Presented by Academician S. N. Bernstein on 29 IV 1966)

Let $Z_n(x, \theta)$ denote the polynomials determined by the segment-functional $0_0, 0_1, \dots, 0_{n-2}, 1, \theta$ for $(n-1)/2 \leq \theta \leq (n+1)/2$ (the critical interval). They all belong to the passport $[n, n, 0]$, and the nature of their deformation with respect to θ has been studied in detail in ⁽¹⁾. Recall that, under a monotone increase of θ , all interior nodes (extremum points) shift to the right. The polynomials $\pm Z_n(x, \theta)$ exhaust all polynomials of the indicated passport.

Definition. The segment-functional $\mu_0, \dots, \mu_{n-1}, \theta$ possesses **Zolotarev stability** if, in the critical interval $\mu'_n \leq \theta \leq \mu''_n$, the segment belongs to the passport $[n, n, 0]$, and moreover is serviced either by all $+Z_n(x, \theta)$, or by all $-Z_n(x, \theta)$ (and only by them).

Theorem 1. *Whatever the prescribed basis $\mu_0, \mu_1, \dots, \mu_{n-2}$, there always exist two numbers $A'_0 \leq A''_0$ such that a segment of the form $\mu_0, \dots, \mu_{n-2}, A, \bar{\theta}$ is stable for $A \geq A''_0$ and for $A \leq A'_0$. In the first case it is serviced by $Z_n(x, \theta)$, and in the second by $-Z_n(x, \theta)$.*

Expand $\mu_0, \mu_1, \dots, \mu_{n-2}, A$ with respect to the nodes $[\sigma_i(\theta)]_1^n$ of any of the polynomials $\pm Z_n(x, \theta)$ ⁽¹⁾; we have

$$\delta_j = R_{n-1}^{(j)}(\bar{\mu}, \theta) / R'_n(\sigma_j) \quad (j = 1, 2, \dots, n), \quad (1)$$

where

$$R_n(x) = \prod_1^n (x - \sigma_i); \quad R_{n-1}^{(j)}(x) = R_n(x) / (x - \sigma_j).$$

The signs of the denominator in (1) alternate with j ; to obtain $\text{Sgn } \delta_j = \pm Z_n(\sigma_j, \theta)$ it is necessary and sufficient that either $R_{n-1}^{(j)}(\bar{\mu}) \geq 0$ ($j = 1, \dots, n$), or $R_{n-1}^{(j)}(\bar{\mu}) \leq 0$ ($j = 1, \dots, n$) (zeros are also possible). If

$$R_{n-1}^{(j)}(x, \theta) = x^n - s_1^{(j)}(\theta)x^{n-1} + s_2^{(j)}(\theta)x - \dots,$$

we have

$$R_{n-1}^{(j)}(\bar{\mu}, \theta) = A - s_1^{(j)}(\theta)\mu_{n-2} + s_2^{(j)}(\theta)\mu_{n-3} - \dots + (-1)^{n-2}s_{n-1}^{(j)}(\theta)\mu_0 = A - M_j(\theta).$$

The family of continuous functions $[M_j(\theta)]_1^n$ on $(n-1)/2 \leq \theta \leq (n+1)/2$ is bounded. Let

$$A_0'' = \max_{(j, \theta)} M_j(\theta) \quad \text{and} \quad A_0' = \min_{(j, \theta)} M_j(\theta);$$

then for fixed $A \geq A_0''$ or $A \leq A_0'$ the (δ_i) have alternating signs. The parameter $\bar{\theta}$ for each θ is determined uniquely from the condition $R_n(\bar{\mu}) = 0$, i.e.

$$\bar{\theta} - s_1(\theta)A + s_2(\theta)\mu_{n-2} - \dots + (-1)^n s_n(\theta)\mu_0 = 0$$

for all $(n-1)/2 \leq \theta \leq (n+1)/2$. This proves the theorem.

Corollary 1. For $A_0' < A < A_0''$, the sign-alternation conditions for (δ_j) are not satisfied for the whole family $+Z_n(x, \theta)$ (or $-Z_n(x, \theta)$); that is, the segment does not possess Zolotarev stability.

Thus, for every basis $\mu_0, \mu_1, \dots, \mu_{n-2}$ there exists a completely determined Zolotarev “critical” interval (A_0', A_0'') ; then and only then is the segment $\mu_0, \dots, \mu_{n-2}, \mu_{n-1}, \bar{\theta}$ stable, if μ_{n-1} is chosen outside or on the boundary of the interval (A_0', A_0'') .

Remark 1. $A_0' = A_0''$ if and only if the basis $(\mu_k)_0^{n-2} \equiv 0$, since here the simultaneous possibility (for $A = A_0' = A_0''$) of two expansions with opposite signs is required.

Corollary 2. Introduce the following notation: $A''(\theta) = \max_{(j)} M_j(\theta)$ and $A'(\theta) = \min_{(j)} M_j(\theta)$. Then the necessary and sufficient condition that the segment $\mu_0, \dots, \mu_{n-2}, \mu_{n-1}, \bar{\theta}$ should under no $\bar{\theta}$ belong to the passport $[n, n, 0]$, i.e. should be completely “detached” from this passport, is the following: $\mu_{n-1} - A''(\theta) < 0$, $\mu_{n-1} - A'(\theta) > 0$ for all θ in the interval $[(n-1)/2, (n+1)/2]$.

Remark 2. If the segment $\mu_0, \dots, \mu_{n-2}, \mu_{n-1}, \theta_0$ is served by some $Z_n(x, \theta_0)$, then the segment $\mu_0, \dots, \mu_{n-2}, \mu_{n-1} + A, \theta + A\theta_0$ is served by the same polynomial for any $A > 0$.

Indeed, the segment $0_0, \dots, 0_{n-2}, 1, \theta_0$ is served by the polynomial $Z_n(x, \theta_0)$; consequently, the segment $0_0, \dots, 0_{n-2}, A, A\theta_0$ is also served, and then so is the termwise sum $\mu_0, \dots, \mu_{n-2}, \mu_{n-1} + A, \theta + A\theta_0$. Thus, serving by one polynomial is extended in two parameters.

Theorem 2. The Zolotarev interval (A'_0, A''_0) of any basis $\mu_0, \mu_1, \dots, \mu_{n-2}$ contains the critical (Chebyshev) interval of the parameter μ_{n-1} .

Thus, if μ'_{n-1}, μ''_{n-1} are the endpoints of the critical interval for the variable parameter μ_{n-1} , then

$$A'_0 \leq \mu'_{n-1} < \mu''_{n-1} \leq A''_0. \quad (2)$$

Indeed, the segment

$$\mu_0, \dots, \mu_{n-2}, \mu''_{n-1} + h \quad (3)$$

for $h > 0$ is served by the polynomial $T_{n-1}(x) \equiv Z_n(k, n/2)$ with all $\delta_i \neq 0$. By the theorem on continuous deformation (1), if $\theta^*(h)$ is the best continuation of (3) to the n -th place, then the segment $\mu_0, \dots, \mu_{n-2}, \mu''_{n-1} + h, \theta^*(h) \pm \varepsilon$ is served by some $Z_n(x, n/2 \pm \varepsilon)$. But if we take $\mu_0, \dots, \mu_{n-2}, \mu''_{n-1} - h$ ($h > 0$), then $T_{n-1}(x)$ is no longer suitable; consequently, some $Z_n(x, n/2 \pm \varepsilon)$ also will not be suitable. Hence $\mu''_{n-1} - h = A < A'_0$ for every $h > 0$. Therefore, $A'_0 \geq \mu''_{n-1}$.

The left-hand part of inequality (2) is proved in the same way.

Corollary. A segment of the form $\mu_0, \dots, \mu_{n-2}, \mu_n, \bar{\theta}$ has “partial Zolotarev stability in the interval $\bar{\theta}^* \leq \bar{\theta} \leq \bar{\theta}^*$.”

Let us give some applications of the results obtained.

Example 1. The segment-functional

$$1, \rho, \dots, \rho^{n-1}, \rho^n + \bar{\theta} \quad (4)$$

for $\rho > 1$ is Zolotarev-stable, since here $A - M_j(\theta) = R_{n-1}^{(j)}(\rho, \theta) > 0$, i.e. the segment determines all $Z_n(x, \theta)$. For $0 < \rho < 1$ the segment $(\rho^i)_0^{n-1}$, when expanded with respect to arbitrary nodes $(\sigma_i)_1^n$ ($0 \leq \sigma_i \leq 1$ and $\sigma_i \neq q$), always gives one repetition of sign for the loads (δ_i) . Consequently, the segment (4) is completely detached from the passport $[n, n, 0]$. For $\rho = 1$ the segment (4) is served over the whole critical interval only by Chebyshev transforms $T_n(ax)$, i.e. although (4) belongs only to the passport $[n, n, 0]$, the requirement of completeness is not fulfilled—the serving is partial, but without involving extraneous passports.

Example 2. The segment-functional $(\mu_i)_0^n = 0_0, \dots, 0_{n-3}, 1_{n-2}, n/2, \bar{\theta}$ gives an example of partial serving by Zolotarev polynomials, with the involvement of polynomials of another passport. Indeed, since here $n/2 = \mu''_{n-1}$, the basis $(\mu_i)_0^{n-1}$ is served by the polynomial $T_{n-1}(x)$

with one unloaded end node $\sigma_1 = 0$. Let us find the boundaries of the Zolotarev critical interval from the conditions of Theorem 1, which here take the form

$$A_0'' = \max_{j, \theta} [\theta - \sigma_j(\theta)]$$

and

$$A_0' = \min_{(j, \theta)} [\theta - \sigma_j(\theta)].$$

For all θ in the interval $[(n-1)/2, (n+1)/2]$ one has

$$\theta - \sigma_n(\theta) \leq \theta - \sigma_{n-1}(\theta) \leq \dots \leq \theta - \sigma_1(\theta).$$

Thus, the max is attained at $\theta - \sigma_1(\theta)$, and the min at $\theta - \sigma_n(\theta)$. Finally we have

$$A_0'' = (n+1)/2 - \tilde{\sigma}_1 (> n/2);$$

$$A_0' = (n-1)/2 - \tilde{\sigma}_{n-1} (< n/2 - 1),$$

where $\tilde{\sigma}_1$ and $\tilde{\sigma}_{n-1}$ are the corresponding nodes of

$$T_n(x) = \cos n \arccos(2x - 1).$$

Thus, the segment $(\mu_i)_0^n$ is served only by part of the polynomials $Z_n(x, \theta)$, and only in part of the critical interval $\theta^* \leq \theta \leq \theta''$. In the remaining part, service belongs to another passport.

Example 3. The segment $(\mu_i)_0^n = \alpha_0, \alpha_1, \dots, \alpha_{n-1}, \bar{\theta}$ with the amorphous basis $(\alpha_i)_0^{n-1}$ cannot, for any $\bar{\theta}$, be served by primitive Zolotarev polynomials, i.e. by those among whose nodes $(\sigma_i)_1^n$ there are both $\sigma_1 = 0$ and $\sigma_n = 1$. Indeed, if for some $\bar{\theta}_0$ one has

$$\alpha_k = \sum_1^{s_1} \delta_i' \sigma_i^k - \sum_1^{s_2} \delta_i'' \sigma_i^k \quad (k = 0, 1, \dots, n-1),$$

then the resulting equalities

$$\alpha_k + \sum_1^{s_1} \delta_i'' \sigma_i^k = \sum_1^{s_1} \delta_i' \sigma_i^k \quad (k = 0, 1, \dots, n-1)$$

are impossible, since on the left we have an amorphous segment, and on the right a nodal one ⁽²⁾.

In this note the question of Zolotarev stability is studied from a somewhat different standpoint than in the article ⁽³⁾.

Leningrad Electrotechnical Institute of Communications
named after M. A. Bonch-Bruевич

Received
6 III 1966

CITED LITERATURE

¹ E. V. Voronovskaya, *The Method of Functionals and Its Applications*, L., 1963.

² E. V. Voronovskaya, DAN, **166**, No. 6 (1966).

³ E. V. Voronovskaya, DAN, **161**, No. 2 (1965).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.