

ON ONE ANALOGUE OF THE HARDY- LITTLEWOOD EQUATION

A. A. Polyanskii

1968

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196801.01585>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 511

MATHEMATICS

A. A. POLYANSKII

ON ONE ANALOGUE OF THE HARDY-LITTLEWOOD EQUATION

(Presented by Academician Yu. V. Linnik on 19 VI 1967)

In the monograph of Yu. V. Linnik ⁽¹⁾, among analogues of the Hardy-Littlewood equation, the equation

$$p_1 p_2 + \xi^2 + \eta^2 = n, \quad (1)$$

is considered, where p_1 and p_2 independently run through the prime numbers; ξ and η run through the integers; n is a sufficiently large natural number (the main parameter).

For the number $S(n)$ of solutions of equation (1), in ⁽¹⁾ an asymptotic formula is derived under the additional restrictions

$$p_1, p_2 > \exp(\ln \ln n)^2.$$

The aim of the present note is to remove these restrictions and to obtain $S(n)$ with a better remainder term than in ⁽¹⁾.

Theorem. As $n \rightarrow \infty$,

$$S(n) = \pi A_0 A(n) \frac{n}{\ln n} \prod_{p|n} \frac{(p-1)(p-\chi(p))}{p^2 - p + \chi(p)} + O\left(\frac{n}{(\ln n)^{1.042}}\right), \quad (2)$$

where

$$A_0 = \prod_{p>2} \left(1 + \frac{\chi(p)}{p(p-1)}\right); \quad \chi(m) \text{ is the nonprincipal character mod } 4; \quad A(n) \text{ is}$$

some arithmetical factor, whose structure is clear from the proof.

We precede the proof of the theorem with two lemmas. Consider the equation

$$ap + b(\xi^2 + \eta^2) = n, \quad (3)$$

where a, b are integers satisfying the conditions: $a = O(\exp(\ln n)^\alpha)$, $b = O(\ln^C n)$, $C > 0$ is a sufficiently large constant, and p runs through the prime numbers. We shall assume that $(a, b) = 1$, $(ab, 2n) = 1$. The latter conditions are not essential.

Let $Q(n)$ be the number of solutions of equation (3).

Lemma 1. As $n \rightarrow \infty$,

$$Q(n) = \pi A_0 \frac{n}{ab \ln n} \prod_{p|an} \frac{(p-1)(p-\chi(p))}{p^2 - p + \chi(p)} \prod_{p|b} \frac{p^2}{p^2 - p + \chi(p)} + O\left(\frac{n}{ab(\ln n)^{1.042}}\right). \quad (4)$$

For the proof of the lemma, following C. Hooley ⁽⁴⁾ and B. M. Bredikhin ⁽⁴⁾, we represent $Q(n)$ in the following form:

$$\begin{aligned} Q(n) &= \sum_{ap+b(\xi^2+\eta^2)=n} 1 = 4 \sum_{ap+2^\lambda bxy=n} \chi(x) = 8 \sum_{\substack{ap+2^\lambda bxy=n \\ x \leq \sqrt{n} n_1^{-1}}} \chi(x) \\ &\quad - 4 \sum_{\substack{ap+2^\lambda bxy=n \\ \sqrt{n} n_1^{-1} < x < \sqrt{n} n_1 \\ y < \sqrt{n} n_1}} \chi(x) + 4 \sum_{\substack{ap+2^\lambda bxy=n \\ \sqrt{n} n_1^{-1} < x < \sqrt{n} n_1}} \chi(x) + O\left(\frac{n}{ab \ln^2 n}\right) \\ &= \Sigma_A - \Sigma_{B_1} + \Sigma_{B_2} + O\left(\frac{n}{ab \ln^2 n}\right), \end{aligned}$$

where $n_1 = \exp(\ln n)^{\alpha_1}$; $\alpha_0, \alpha_1 > 0$ are sufficiently small constants.

Σ_A is evaluated with the aid of E. Bombieri' s lemma ⁽⁵⁾, while Σ_{B_1} and Σ_{B_2} are estimated by S. Hooley' s method ⁽⁴⁾ according to the scheme developed in ⁽⁸⁾.

Equation (3) may be regarded as a special case of the linear equation $ax+by = n$, when x runs through the primes and y assumes values of the quadratic form $\varphi(\xi, \eta) = \xi^2 + \eta^2$. This equation is a natural generalization of the classical Hardy-Littlewood equation and is of interest from the point of view of possible applications (see, for example, below and ^(7, 8)).

Next, let $S_1(n)$ be the number of solutions of equation (1) when

$$p_1, p_2 > P_0 = \exp(\ln n)^{\alpha_0}.$$

Lemma 2. As $n \rightarrow \infty$,

$$S_1(n) = \pi A_0 \prod_{p|n} \frac{(p-1)(p-\chi(p))}{p^2-p+\chi(p)} \sum_{\substack{p_1 p_2 < n \\ p_1 p_2 > P_0}} 1 + O\left(\frac{n}{(\ln n)^{1.042}}\right). \quad (5)$$

Proof. We write $S_1(n)$ in the form

$$\begin{aligned} S_1(n) &= 2 \sum_{\substack{p_1 p_2 + \xi^2 + \eta^2 = n \\ P_0 < p_1 < P}} 1 + \sum_{p_1 p_2 + \xi^2 + \eta^2 = n} 1 + O\left(\frac{n}{\ln^2 n}\right) = \\ &= 2S_2(n) + S_3(n) + O\left(\frac{n}{\ln^2 n}\right), \end{aligned} \quad (6)$$

where $P = \exp \ln n \frac{\ln \ln \ln n}{K \ln \ln n}$, and K is a sufficiently large constant.

With the aid of the dispersion method ⁽¹⁾ (see ^(6,13,10) and p. 131) we obtain

$$S_2(n) = \pi A_0 \prod_{p|n} \frac{(p-1)(p-\chi(p))}{p^2-p+\chi(p)} \sum_{\substack{p_1 p_2 < n \\ P_0 < p_1 < P}} 1 + O\left(\frac{n}{\ln^2 n}\right). \quad (7)$$

Refining the proof of Theorem 2 of ⁽²⁾, we obtain

$$S_3(n) = \pi A_0 \prod_{p|n} \frac{(p-1)(p-\chi(p))}{p^2-p+\chi(p)} \sum_{\substack{p_1 p_2 < n \\ p_1, p_2 > P}} 1 + O\left(\frac{n}{(\ln n)^{1.042}}\right). \quad (8)$$

From (6)–(8), (5) follows.

Proof of the theorem. Decompose $S(n)$ as follows:

$$S(n) = 2 \sum_{\substack{p_1 p_2 + \xi^2 + \eta^2 = n \\ p_1 < P_0}} 1 + S_1(n) + O\left(\frac{n}{\ln^2 n}\right) = 2S_4(n) + S_1(n) + O\left(\frac{n}{\ln^2 n}\right). \quad (9)$$

We find $S_4(n)$:

$$S_4(n) = \sum_{\substack{(p_1, n)=1 \\ p_1 < P_0}} \sum_{p_1 p_2 + \xi^2 + \eta^2 = n} 1 + \sum_{\substack{p_1 | n \\ p_1 < P_0}} \sum_{p_1 p_2 + \xi^2 + \eta^2 = n} 1 = \Sigma_1 + \Sigma_2. \quad (10)$$

The inner sum in Σ_1 is evaluated with the aid of Lemma 1. We obtain

$$\Sigma_1 = \sum_{\substack{(p_1, n)=1 \\ p_1 < P_0}} \left(\pi A_0 \frac{n}{p_1 \ln n} \prod_{p|p_1 n} \frac{(p-1)(p-\chi(p))}{p^2 - p + \chi(p)} + O\left(\frac{n}{p_1 (\ln n)^{1.042}}\right) \right). \quad (11)$$

Further, by virtue of

$$\sum_{\substack{d|ml \\ (m, l)=1}} \chi(d) = \sum_{d|m} \chi(d) \sum_{d|l} \chi(d)$$

and Lemma 1, we have (with ad-
with admissible error)

$$\begin{aligned} \Sigma_2 &= 4 \sum_{\substack{p_1|n \\ p_1 < P_0}} \sum_{\lambda} \sum_{s=0, 1, \dots} \sum_{d|p_1^{1+s}} \chi(d) \sum_{p_2 + 2^\lambda p_1^{sxy} = n_0} \chi(x) \\ &= \sum_{\substack{p_1|n \\ p_1 < P_0}} \sum_{s=0, 1, \dots} \sum_{d|p_1^{1+s}} \chi(d) \left(\pi A_0 \frac{n}{p_1^{1+s} \ln n} \prod_{p|n_0} \frac{(p-1)(p-\chi(p))}{p^2 - p + \chi(p)} \frac{p_1^2}{p_1^2 - p_1 + \chi(p_1)} + \right. \\ &\quad \left. + O\left(\frac{n}{p_1^{1+s} (\ln n)^{1.042}}\right) \right), \end{aligned} \quad (12)$$

where $n_0 = n/p_1$.

Now (2) follows from (5), (9)-(12).

By the methods of the papers ^(6, 8, 9) one can derive geometric and ergodic properties of the solutions of equation (1).

Equation (1) can be generalized. By applying analogous means one can treat the equation

$$p_1 p_2 \dots p_k + \varphi(\xi, \eta) = n,$$

where $k > 2$, and $\varphi(\xi, \eta)$ is some prescribed quadratic form.

Kuibyshev Pedagogical Institute
named after V. V. Kuibyshev

Received
13 VI 1967

REFERENCES

- ¹ Yu. V. **Linnik**, *The dispersion method in binary additive problems*, L., 1961.
- ² Yu. V. **Linnik**, *Matem. sborn.*, 52 (94), 2, 661 (1960).
- ³ B. M. **Bredikhin**, *UMN*, 20, No. 2, 89 (1965).
- ⁴ C. **Hooley**, *Acta Math.*, 97, 189 (1957).
- ⁵ E. **Bombieri**, *On the Large Sieve*, Milano, 1965.
- ⁶ A. A. **Polyanskii**, *DAN*, 168, No. 1 (1966).
- ⁷ A. A. **Polyanskii**, *The solution of certain binary equations of Hardy-Littlewood type*, Dissertation, Kuibyshev, 1966.
- ⁸ B. M. **Bredikhin**, Yu. V. **Linnik**, *DAN*, 166, No. 6, 1267 (1966).
- ⁹ A. I. **Vinogradov**, *Matem. zametki*, 1, No. 2, 189 (1967).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.