

ON A CLASS OF OPERATORS WITH SPECTRUM CONCENTRATED AT ZERO

L. E. Isaev

1968

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196801.32729>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 513.88+517.948.35

MATHEMATICS

L. E. ISAEV

ON A CLASS OF OPERATORS WITH SPECTRUM CONCENTRATED AT ZERO

(Presented by Academician I. M. Vinogradov, April 7, 1967)

Let A be a bounded linear operator acting in a separable Hilbert space $\mathfrak{H}$. The operator A is called **completely non-selfadjoint** if there is no subspace invariant with respect to A and A^* in which it induces a selfadjoint operator.

We shall assign a completely non-selfadjoint operator A to the class $\Lambda^{(\text{exp})}$ if it satisfies the following conditions: 1) A has no spectral points distinct from zero; 2) $\left(A - \frac{1}{\mu}E\right)^{-1}$ is a function of exponential type; 3) A is dissipative, i.e. $A_I = (A - A^*)/2i \geq 0$. If, moreover, $\text{sp } A_I < \infty$ ($\text{sp } A_I = \infty$), then we shall assign A to the class $\Lambda^{[\text{exp}]}$ ($\Lambda_\infty^{(\text{exp})}$).

The class $\Lambda^{[\text{exp}]}$ was studied by M. S. Livshits, M. S. Brodskii, M. G. Krein, I. Ts. Gokhberg, and G. E. Kisilevskii in papers ⁽¹⁻⁸⁾. The present article is devoted to extending some results of M. S. Brodskii and G. E. Kisilevskii to the class $\Lambda^{(\text{exp})}$.

We note that all operators of the class $\Lambda^{[\text{exp}]}$ are completely continuous, which cannot be said, in general, of operators of the class $\Lambda^{(\text{exp})}$.

An operator $A \in \Lambda^{(\text{exp})}$ can be included in the node

$$\theta = \begin{pmatrix} A & K \\ \mathfrak{H} & \mathfrak{G} \end{pmatrix},$$

where $\mathfrak{G}$ is some separable space and K is a bounded linear operator acting from $\mathfrak{G}$ into $\mathfrak{H}$, such that $KK^* = A_I$. The function

$$W(\lambda) = E - 2iK^*(A - \lambda E)^{-1}K$$

is called the **characteristic** function of the node θ . It has the following properties: 1) its values are bounded linear operators acting in $\mathfrak{G}$; 2) $W(1/\mu)$ is an entire function of exponential type; 3) $\lim_{\lambda \rightarrow \infty} \|W(\lambda) - E\| = 0$; 4) $W^*(\lambda)W(\lambda) - E \geq 0$ ($\text{Im } \lambda > 0$); 5) $W^*(\lambda)W(\lambda) - E = 0$ ($\text{Im } \lambda = 0, \lambda \neq 0$).

We shall denote the types of growth of the functions $\left(A - \frac{1}{\mu}E\right)^{-1}$ and $W\left(\frac{1}{\mu}\right)$, respectively, by $\sigma[A]$ and $\sigma[W]$. Obviously,

$$\sigma[W] \leq \sigma[A]. \quad (1)$$

By $\Omega_{\mathfrak{G}}^{(\text{exp})}$ we denote the class of all functions possessing properties 1)–5).

Consider the Hilbert space $\widetilde{\mathfrak{L}}_2(0, l)$ ($l < \infty$) of matrices of the form

$$f(x) = \|f_1(x) \ f_2(x) \dots\|, \quad (2)$$

where $f_j(x)$ ($j = 1, 2, \dots$) are functions measurable on $(0, l)$ satisfying the condition

$$\sum_{j=1}^{\infty} \int_0^l |f_j(x)|^2 dx < \infty.$$

The scalar product in $\widetilde{\mathfrak{L}}_2(0, l)$ is defined by the formula

$$(f, g) = \sum_{j=1}^{\infty} \int_0^l f_j(x) \overline{g_j(x)} dx \quad \left(g(x) = \|g_1(x) \ g_2(x) \dots\| \in \widetilde{\mathfrak{L}}_2(0, l)\right).$$

Define in $\mathcal{L}_2(0, l)$ the operator $\tilde{I}_l$, which assigns to the matrix (2) the matrix

$$\tilde{I}_l f(x) = \|I_l f_1(x) \ I_l f_2(x) \dots\| \quad \left(I_l f_j(x) = 2i \int_x^l f_j(t) dt\right).$$

Theorem 1*. *If the operator T is induced by the operator $\tilde{I}_l$ in one of its invariant subspaces, then $T \in \Lambda^{(\text{exp})}$ and $\sigma[T] \leq 2l$. Conversely, if $A \in \Lambda^{(\text{exp})}$ and $\sigma[A] \leq 2l$, then there exists an invariant subspace of the operator $\tilde{I}_l$ in which an operator T , unitarily equivalent to A , is induced.*

Proof. The first assertion of the theorem is trivial. To prove the second, embed A in the simple dissipative node

$$\theta = \begin{pmatrix} A & K \\ \mathfrak{H} & \mathfrak{G} \end{pmatrix},$$

and let $W(\lambda)$ be the characteristic operator-function of this node. For all $\lambda \neq 0$ there exists the operator $W^{-1}(\lambda)$, and the estimate

$$\|W(\lambda)\| \leq e^{2l |\operatorname{Im} \frac{1}{\lambda}|}$$

holds, from which it follows that $W^{-1}(\lambda)e^{2il/\lambda} \in \Omega_{\mathfrak{H}}^{(\text{exp})}$. To complete the proof it remains to refer to Theorem 2 of paper ⁽²⁾.

Theorem 2. *Every function $W(\lambda) \in \Omega_{\mathfrak{H}}^{(\text{exp})}$ is characteristic for some node*

$$\theta = \begin{pmatrix} A & K \\ \mathfrak{H} & \mathfrak{G} \end{pmatrix},$$

where $A \in \Lambda^{(\text{exp})}$ and $\sigma[A] = \sigma[W]$.

It is known that every completely noncontinuous operator has a nontrivial invariant subspace ⁽¹⁰⁾ (see also ⁽¹¹⁾, § 65). It turns out that operators of the class $\Lambda_{\infty}^{(\text{exp})}$ also possess this property. Moreover, the following holds.

Theorem 3. *For every operator $A \in \Lambda_{\infty}^{(\text{exp})}$ there exists a continuous strictly increasing function $P(x)$ ($0 \leq x \leq \infty$) satisfying the following conditions: 1) the values of the function $P(x)$ are orthoprojectors in $\mathfrak{H}$, with $P(0) = 0$, $P(\infty) = E$; 2) all subspaces $P(x)\mathfrak{H}$ are invariant with respect to A ;*

$$3) \quad \text{sp}(P(x)A_I P(x)) = x \quad (0 \leq x \leq \infty). \quad (3)$$

Proof. Consider the operator $B = -A^*$. By Theorem 1, the generality of the argument will not be violated if we assume that B is induced by the operator $\tilde{I}_l$ in some invariant subspace $\mathfrak{H} \subset \tilde{\mathcal{L}}_2(0, l)$ ($l = \frac{1}{2}\sigma[A]$). In $\tilde{\mathcal{L}}_2(0, l)$ there exists a subspace $\mathfrak{H}_0$ possessing the following properties: 1) $\mathfrak{H}$ and $\mathfrak{H}_0^\perp = \tilde{\mathcal{L}}_2(0, l) \ominus \mathfrak{H}_0$ are invariant with respect to $\tilde{I}_l$; 2) in $\mathfrak{H}_0$ the operator $\tilde{I}_l$ induces a Volterra operator with a nuclear imaginary component; 3) $\mathfrak{H} \cap \mathfrak{H}_0^\perp \neq \mathfrak{H}$. From Lemma 4 of paper ⁽⁴⁾ it follows that $\mathfrak{H} \cap \mathfrak{H}_0^\perp \neq 0$ and that A induces in its invariant subspace $\mathfrak{H} \ominus (\mathfrak{H} \cap \mathfrak{H}_0^\perp)$ a Volterra operator with nuclear imaginary component. The subsequent arguments differ only insignificantly from those given in Sec. 1 of paper ⁽³⁾.

An operator is called **unicellular** if the set of all its invariant subspaces is ordered by inclusion.

Theorem 4. *The class $\Lambda_{\infty}^{(\text{exp})}$ contains no unicellular operators. In other words, every unicellular operator of the class $\Lambda^{(\text{exp})}$ is Volterra and has a nuclear imaginary component.*

The proof follows from Theorem 3 and the criterion for unicellularity of Volterra operators with nuclear imaginary components, established by M. S. Brodskii and G. E. Kisilevskii in the work ⁽⁴⁾.

* For the case $A \in \Lambda^{[\text{exp}]}$ Theorem 1 was proved by G. E. Kisilevskii.

Theorem 5. Let A be a completely non-self-adjoint operator whose spectrum contains no points different from zero. In order that A belong to the class $\Lambda^{(\text{exp})}$,

it is necessary and sufficient that there exist a sequence of subspaces invariant with respect to A , $\mathfrak{H}_1 \subset \mathfrak{H}_2 \subset \dots$, satisfying the following conditions:

- 1) $\bigcup_{j=1}^{\infty} \mathfrak{H}_j = \mathfrak{H}$;
- 2) the operators A_j , induced in $\mathfrak{H}_j$, belong to the class $\Lambda^{[\text{exp}]}$;
- 3) the set of numbers $\sigma[A_j]$ is bounded.

Proof. Necessity follows from Theorem 3. To prove sufficiency, include A and A_j in simple dissipative nodes

$$= \begin{pmatrix} A & K \\ \mathfrak{H} & \mathfrak{G} \end{pmatrix} \quad \text{and} \quad \theta_j = \begin{pmatrix} A_j & P_{jK} \\ \mathfrak{H}_j & \mathfrak{G} \end{pmatrix},$$

where P_j is the orthoprojector onto $\mathfrak{H}_j$. Let $W(\lambda)$ and $W_j(\lambda)$ be the characteristic functions of these nodes. Since $\lim_{j \rightarrow \infty} W_j(\lambda)f = W(\lambda)f$ ($f \in \mathfrak{G}$) and the set of numbers $\sigma[A_j]$ is bounded, $W(\lambda)$ has finite type of growth. The required result follows from Theorem 2.

The function $P(x)$ ($0 \leq x \leq \infty$), introduced in Theorem 3, makes it possible to obtain the following generalization of the known theorem on the triangular representation of Volterra operators ⁽³⁾ (see also ⁽¹¹⁾, §115).

Theorem 6. Every operator $A \in \Lambda^{(\text{exp})}$ can be represented in the form

$$A = 2i \lim_{x \rightarrow \infty} \int_0^x P(t) A_I dP(t),$$

where all the integrals exist in the sense of the operator norm, and the passage to the limit is carried out in the sense of strong convergence.

The author expresses deep gratitude to M. S. Brodskii for his guidance of this work.

Odessa State Pedagogical Institute named after K. D. Ushinsky

Received
5 IV 1967

REFERENCES

- ¹ M. S. Livshits, *Matem. sborn.*, **34** (76) (1954).
- ² M. S. Brodskii, M. S. Livshits, *UMN*, **13**, no. 1 (79), 3 (1958).
- ³ M. S. Brodskii, *UMN*, **16**, no. 1 (97), 135 (1961).
- ⁴ M. S. Brodskii, G. E. Kisilevskii, *Izv. AN SSSR, ser. matem.*, **30**, no. 6 (1966).
- ⁵ G. E. Kisilevskii, *DAN*, **159**, no. 3 (1964).
- ⁶ G. E. Kisilevskii, *Tr. 1 republican conference of young researchers*, Kiev, 1964.
- ⁷ G. E. Kisilevskii, *DAN*, **159**, no. 4 (1964).

⁸ I. Ts. Gohberg, M. G. Krein, *Introduction to the Theory of Linear Non-Self-Adjoint Operators*, Moscow, 1965.

⁹ M. S. Brodskii, Yu. L. Shmul' yan, *UMN*, **19**, no. 1 (115), 143 (1964).

¹⁰ N. Agonszajn, R. J. Smith, *Ann. Math.*, **60**, 316 (1954).

¹¹ N. I. Akhiezer, I. M. Glazman, *Theory of Linear Operators in Hilbert Space*, Moscow, 1966.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.