

ANALYTIC REPRESENTATIONS OF THE FULL LINEAR GROUP OVER A FINITE FIELD

S. I. Gel'fand

1968

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196801.72532>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 519.46

MATHEMATICS

S. I. GELFAND

ANALYTIC REPRESENTATIONS OF THE FULL LINEAR GROUP OVER A FINITE FIELD

(Presented by Academician I. G. Petrovsky, 29 I 1968)

This note studies one important class of irreducible representations of the group $G_n = GL(n, K_q)$, where K_q is the field of order q .

Let $Z_n \subset G_n$ be the subgroup of upper triangular matrices with ones on the diagonal.

Definition 1. A one-dimensional representation χ of the subgroup Z_n is called **degenerate** if there exists another one-dimensional representation χ' of Z_n for which $\text{Ker } \chi' \subsetneq \text{Ker } \chi$.

It is not hard to see that any one-dimensional representation χ of the group Z_n has the form

$$\chi(\xi) = \chi_0(a_1\xi_{12} + a_2\xi_{23} + \cdots + a_{n-1}\xi_{n-1,n}),$$

where χ_0 is a nontrivial additive character of the field K_q , $a_i \in K_q$, and the representation χ is nondegenerate if and only if all $a_i \neq 0$.

It is known ⁽²⁾ that the restriction of any irreducible representation T of the group G_n to Z_n contains at least one one-dimensional representation.

Definition 2. A representation T of the group G_n is called an **analytic representation** (briefly, an A -representation) if the restriction of T to Z_n contains no degenerate one-dimensional representations.

Introduce two subgroups of the group G_n : H_n is the subgroup of matrices $\|a_{ij}\|$ for which

$$a_{n1} = a_{n2} = \cdots = a_{n,n-1} = 0, \quad a_{nn} = 1;$$

$B_n \subset H_n$ is the subgroup of matrices $\|a_{ij}\|$ for which

$$a_{11} = a_{22} = \cdots = a_{nn} = 1$$

and all off-diagonal elements, except the elements of the last column, are equal to zero.

Proposition 1. *With the exception of the special case $q = n = 2$, the group H_n has one irreducible representation T_0 of dimension*

$$(q-1)(q^2-1)\cdots(q^{n-1}-1) = k_n(q);$$

the dimensions of the remaining irreducible representations of H_n are smaller than $k_n(q)$.

Proof. H_n is the semidirect product of the commutative normal divisor B_n and the subgroup G_{n-1} . Therefore ⁽⁴⁾ any irreducible representation T of the group H_n is induced by a representation μL of the group $B_n A_\mu$, where μ is a one-dimensional representation of B_n , $A_\mu \subset G_{n-1}$ is the subgroup of those $g \in G_{n-1}$ such that $\mu(gbg^{-1}) = \mu(b)$, $b \in B_n$, and L is an irreducible representation of A_μ . To finish the proof one must apply a simple induction on n .

Proposition 2. *The restriction of an irreducible A -representation of the group G_n to H_n coincides with T_0 .*

Proof. We first show that in the restriction of any irreducible representation of the group H_n different from T_0 to Z_n there is a degenerate one-dimensional representation. Suppose this has already been proved for the group H_{n-1} .

Any irreducible representation T of the group H_n is induced by a representation μL of the group $B_n A_\mu$. If μ is the identity representation, then for any one-dimensional representation χ of the group Z_n contained in the restriction of T to Z_n , $a_{n-1} = 0$, and hence χ is degenerate. Suppose now that μ is a nonidentity representation. Then A_μ is isomorphic to H_{n-1} . If $\dim T < k_n(q)$, then $\dim L < k_{n-1}(q)$, and hence the restriction of L to Z_{n-1} contains a degenerate one-dimensional representation. It follows at once that the restriction of T to Z_n also contains a degenerate one-dimensional representation. Therefore the restriction of any A -representation of the group G_n to H_n is a multiple of T_0 .

On the other hand, every nondegenerate one-dimensional representation χ of the group Z_n enters into the restriction of T to Z_n no more than once ⁽²⁾. Therefore the restriction of T to H_n coincides with T_0 , and the proposition is proved.

The representation T_0 of the group H_n can be realized in the following way. Fix a nondegenerate character

$$\chi(\xi) = \chi_0(\xi_{12} + \cdots \xi_{n-1,n})$$

of the group Z_n . The representation T_0 is defined in the space E_χ of functions on G_{n-1} for which

$$f(\zeta g) = \chi(\zeta)f(g), \quad \zeta \in Z_{n-1}, \quad g \in G_{n-1}^*,$$

with scalar product

$$(\varphi_1, \varphi_2)_{n-1} = |G_{n-1}|^{-1} \sum_{g \in G_{n-1}} \varphi_1(g) \overline{\varphi_2(g)}$$

($|G|$ is the number of elements in the finite group G). The operator $T_0(h)$, where $h \in H_n$, $h = ba$, $b \in B_n$, $a \in G_{n-1}$, acts by the formula

$$T_0(h)f(g) = \chi(gbg^{-1})f(ga). \quad (1)$$

This representation is clearly unitary. From (1) it follows that the restriction of T_0 to $G_{n-1} \subset H_n$ coincides with the representation T_χ of the group G_{n-1} induced by the one-dimensional representation χ of the group Z_{n-1} . From (2) follows

Proposition 3. *The restriction of an irreducible A -representation T of the group G_n to G_{n-1} contains each irreducible component no more than once.*

In what follows, an essential role is played by the so-called Bessel functions, introduced in (1,2). They are defined as follows.

Consider the restriction of the regular representation of the group G_n to those functions $f(g)$ on G_n for which

$$f(\zeta g) = \chi(\zeta)f(g), \quad \zeta \in Z_n, \quad g \in G_n.$$

According to (2), in the space of such functions there exists a unique invariant subspace E_T , the representation in which is equivalent to T . In E_T there exists, up to a factor, a unique vector $J(g)$ for which

$$J(g\zeta) = J(g)\chi(\zeta), \quad \zeta \in Z_n.$$

This vector is called the Bessel function. From the definition it follows easily that

1.

$$J(\zeta_1 g \zeta_2) = \chi(\zeta_1 \zeta_2)J(g).$$

2.

$$\overline{J(g)} = J(g^{-1}).$$

3.

$$J(e) = 1.$$

4.

$$J * J = cJ; \quad * \text{ is convolution on the group } G_n,$$

$$f_1 * f_2(g) = |G_n|^{-1} \sum_{g' \in G_n} f_1(g')f_2(gg'^{-1}).$$

In what follows we normalize J so that $c = k_n(g)$. It is easy to establish the following expression for $J(g)$ in terms of the character ψ of the representation T :

$$J(g) = |Z_n|^{-1} \sum_{\zeta \in Z_n} \chi(\zeta)\psi(g\zeta^{-1}).$$

Let us now determine where the function $J(g)$ is concentrated. It is known ^(3,5) that any element $g \in G_n$ can be written in the form

$$g = \zeta_1 \delta s \zeta_2,$$

where $\zeta_1, \zeta_2 \in Z_n$, $\delta \in D_n$ is a diagonal matrix, and $s \in W_n$ is a permutation matrix. Hence, in view of property 1, it suffices to determine for which δs one has $J(\delta s) \neq 0$. Denote by s_k the matrix of order k in which the second diagonal consists of ones, and all other entries are zero. By

$$s_{i_1, \dots, i_l}(\delta_1, \dots, \delta_l)$$

we denote the matrix

$$s_{i_1, \dots, i_l}(\delta_1, \dots, \delta_l) = \begin{pmatrix} \delta_1 s_{i_1} & 0 \\ & \ddots \\ 0 & \delta_l s_{i_l} \end{pmatrix} s_n,$$

where $\delta_i \in K_q^*$, $i_1, \dots, i_l$ are integers, and

$$i_1 + \dots + i_l = n.$$

* We assume that G_{n-1} is embedded in G_n so that if $g \in G_{n-1}$, $\begin{pmatrix} g & 0 \\ 0 & 1 \end{pmatrix} \in G_n$.

Proposition 4. If $J(\delta s) \neq 0$, then $\delta s = s_{i_1, \dots, i_l}(\delta_1, \dots, \delta_l)$ for some i_k and δ_k .

The proof follows easily from property 1.

Let φ_i be an orthonormal basis in the space E_χ , and let $k_{ij}(h)$ be the corresponding matrix elements of the representation T_0 of the group H_n . Let $f_i(g)$ be the image of φ_i in E_T under the isometric mapping $\tau: E_\chi \rightarrow E_T$, intertwining the operators of the representation T_0 .

Lemma. For any i and j the equality

$$|H_n|^{-1} \sum_{h \in H_n} J(gh) \overline{k_{ij}(h)} = \gamma_i f_j(g), \quad (2)$$

holds, where γ_j are the coefficients in the expansion

$$J(g) = \sum_i \gamma_i f_i(g).$$

Proof.

$$J(gh) = \sum_i \gamma_i f_i(gh) = \sum_{i,j} \gamma_i k_{ij}(h) f_j(g),$$

$$\sum_{h \in H_n} J(gh) \overline{k_{i_3 j_3}(h)} = \sum_{i,j} \gamma_i f_j(g) \sum_{h \in H_n} k_{ij}(h) \overline{k_{i_3 j_3}(h)} = |H_n| \gamma_i f_j(g).$$

Theorem. The matrix elements $K_{ij}(g)$ of the representation T in the basis $\{f_i\}$ are given by the formula

$$K_{ij}(g) = c \sum_{a_1, a_2 \in G_{n-1}} J(a_2 g a_1^{-1}) \varphi_i(a_1) \overline{\varphi_j(a_2)},$$

where $c = |G_{n-1}|^{-3} |Z_{n-1}|^{-1}$.

Proof. Transform formula (2):

$$\begin{aligned} \gamma_i f_j(g) &= |H_n|^{-1} \sum_{h \in H_n} J(gh) \overline{k_{ij}(h)} = \\ &= |H_n|^{-1} |G_{n-1}|^{-1} \sum_{\substack{a, a_0 \in G_{n-1} \\ b \in B_n}} J(g b a_0) \overline{\chi(aba^{-1})} \varphi_i(a a_0) \overline{\varphi_j(a)} = \\ &= |H_n|^{-1} |G_{n-1}|^{-1} \sum_{\substack{a, a_0 \in G_{n-1} \\ b \in B_n}} J(g a_0) \overline{\varphi_i(a a_0)} \varphi_j(a) \chi(a_0^{-1} b a_0) \overline{\chi(aba^{-1})}. \end{aligned}$$

Make the substitution $aa_0 = a'$, $aba^{-1} = b_1$. Then

$$\gamma_i f_j(g) = |G_{n-1}|^{-1} |H_n|^{-1} \sum_{a, a' \in G_{n-1}} J(g a^{-1} a') \overline{\varphi_i(a')} \varphi_j(a) \sum_{b_1 \in B_n} \overline{\chi(b_1)} \chi(a'^{-1} b_1 a').$$

If $\varphi^0 \in E_\chi$ is such a function that $\varphi^0(a) = 0$ for $a \in Z_{n-1}$, $\varphi^0(e) = 1$, and β_i are its coordinates in the basis φ_i , then

$$\sum_{a \in G_{n-1}} J(g a^{-1}) \varphi_j(a) = c_0 f_j(g), \quad c_0 = |G_{n-1}|^2 \sum_i \gamma_i \overline{\beta_i}. \quad (3)$$

The matrix elements $K_{ij}(g)$ of the representation T are computed by the formula

$$K_{ij}(g_0) = |G_n|^{-1} \sum_{g \in G_n} f_i(g g_0) \overline{f_j(g)}.$$

Substituting (3), we obtain

$$\begin{aligned} K_{ij}(g_0) &= |G_n|^{-1} c_0^{-2} \sum_{\substack{g \in G_n \\ a_1, a_2 \in G_{n-1}}} J(g g_0 a_1^{-1}) \overline{J(g a_2^{-1})} \varphi_i(a_1) \overline{\varphi_j(a_2)} = \\ &= c_0^{-2} \sum'_{a_1, a_2 \in G_{n-1}} \varphi_i(a_1) \overline{\varphi_j(a_2)} |G_n|^{-1} \sum'_{g \in G_n} J(g a_2 g_0 a_1^{-1}) \overline{J(g)}. \end{aligned}$$

In view of properties 2 and 4 of the Bessel function, the inner sum is equal to

$$|G_n|^{-1} \sum_{g \in G_n} J(ga_2g_0a_1^{-1}) \overline{J(g)} = J * J(a_2g_0a_1^{-1}) = \bar{k}_n(q) J(a_2g_0a_1^{-1}).$$

Finally we obtain that

$$K_{ij}(g_0) = \bar{k}_n(q) c_0^{-2} \sum_{a_1, a_2 \in G_{n-1}} J(a_2g_0a_1^{-1}) \varphi_i(a_1) \overline{\varphi_j(a_2)}.$$

To find c_0 , observe that $\sum_i \bar{\gamma}_i \bar{\beta}_i = (\tau^{-1}J, \varphi^0)_{n-1}$. But $\tau^{-1}J$, as is not difficult to see, is a multiple of φ^0 , $|\varphi^0| = \sqrt{|\bar{Z}_{n-1}| |G_{n-1}|^{-1}}$ and

$$|J| = \left(|G_n|^{-1} \times \sum_g J(g) \overline{J(g)} \right)^{1/2} = \sqrt{\bar{k}_n(q)} \sqrt{J(e)} = \sqrt{\bar{k}_n(q)}.$$

Therefore

$$c_0 = \sqrt{|\bar{Z}_{n-1}| \bar{k}_n(q) |G_{n-1}|^3},$$

and the theorem is proved.

The theorem just proved shows the importance of the Bessel function for the study of representations of the group G_n . We shall now give an expression for the Bessel function of an irreducible A -representation of the group G_3 .*

Consider the extension K_{q^3}/K_q . Each irreducible A -representation of the group G_3 is determined by a multiplicative character π of the field K_{q^3} , not identically equal to one on K_q . For $\sigma \in K_{q^3}$ introduce the symmetric functions

$$P_1(\sigma) = \sigma + \sigma^q + \sigma^{q^2}; \quad P_2(\sigma) = \sigma^{q+1} + \sigma^{q^2+1} + \sigma^{q^2+q}; \quad P_3(\sigma) = \sigma^{q^2+q+1}.$$

It is clear that for any $\sigma \in K_{q^3}$, $P_i(\sigma) \in K_q$ ($i = 1, 2, 3$), and the equation

$$Q_\sigma(\lambda) = \lambda^3 - P_1(\sigma)\lambda^2 + P_2(\sigma)\lambda - P_3(\sigma) = 0$$

has roots $\sigma, \sigma^q, \sigma^{q^2}$.

Using the formulas for the characters of irreducible representations of the group G_3 , given in ⁽⁶⁾ (see also ⁽⁷⁾), one can obtain the following expressions for the function $J(g)$:

$$J(s_3(\lambda)) = \pi(\lambda),$$

$$J(s_{1,2}(\lambda_1, \lambda_2)) = q^{-2} \sum_{\sigma^{q^2+q+1}=\lambda_1 \lambda_2^2} \chi_0(\lambda_2^{-1} P_1(\sigma)) \pi(\sigma),$$

$$J(s_{2,1}(\lambda_1, \lambda_2)) = q^{-2} \sum_{\sigma^{q^2+q+1}=\lambda_1^2 \lambda_2} \chi_0(-\lambda_1^{-1} \lambda_2^{-1} P_2(\sigma)) \pi(\sigma),$$

$$J(s_{1,1,1}(\lambda_1, \lambda_2, \lambda_3)) = q^{-3} \sum_{\sigma^{q^2+q+1}=-\lambda_1 \lambda_2 \lambda_3} u(\sigma) \pi(\sigma) - q^{-2} \pi(\lambda_2) \delta(\lambda_1 \lambda_3 + \lambda_2^2),$$

where

$$u(\sigma) = \sum_{\substack{\zeta_1, \zeta_2 \in K_q \\ \zeta_1 \zeta_2 = -\lambda_3^{-1} \lambda_2^{-2} Q_\sigma(\lambda_2)}} \chi_0(\zeta_1 + \zeta_2),$$

$\delta(x)$ is the Kronecker symbol.

According to Proposition 4, $J(\delta s) = 0$ for the remaining δs . In view of property 1, the formulas written down make it possible to find $J(g)$ for all $g \in G_3$.

In conclusion, the author expresses his gratitude to M. I. Graev for his constant attention to the work and to A. A. Kirillov for valuable advice and suggestions.

Moscow State University
named after M. V. Lomonosov

Received
25 I 1968

CITED LITERATURE

1. I. M. Gelfand, M. I. Graev, DAN, **146**, No. 4 (1962).
2. I. M. Gelfand, M. I. Graev, DAN, **147**, No. 3 (1962).
3. I. M. Gelfand, M. A. Naimark, Tr. Mat. Inst. im. V. A. Steklova, AN SSSR, **36** (1950).
4. G. Mackey, Sborn. per. Matematika, **6**, No. 6 (1962).
5. K. Chevalley, Sborn. per. Matematika, **2**, No. 1 (1958).

6. R. Steinberg, *Canad. J. Math.*, **3**, No. 2, 225 (1951).

7. J. A. Green, *Trans. Am. Math. Soc.*, **80**, 402 (1955).

* In ⁽¹⁾ the Bessel function was computed for the group G_2 .

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.