

REGULARIZATION ALGORITHMS FOR LINEAR EQUATIONS WITH UNBOUNDED OPERATORS

A. B. Bakushinskii

1968

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196801.81006>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.948.35

MATHEMATICS

A. B. BAKUSHINSKII

REGULARIZATION ALGORITHMS FOR LINEAR EQUATIONS WITH UNBOUNDED OPERATORS

(Presented by Academician A. N. Tikhonov on 4 III 1968)

Consider the operator equation:

$$Ax = f, \quad (1)$$

where A is a linear closed operator with an everywhere dense domain of definition in some Hilbert space H_1 , acting into the Hilbert space H_2 , having no bounded inverse, and $f \in A(D_A)$. The spaces H_1 and H_2 are assumed to be complete and real.

Interest in the approximate solution of such equations has recently increased thanks to the works of A. N. Tikhonov (for example ^(1,2)). In these works the problem of solving (1) was reduced to the problem of constructing a regularizing algorithm. This reduction proved convenient, in particular, for the numerical solution of equation (1).

In what follows, by a regularizing algorithm (r.a.) for equation (1) we shall mean a family of linear bounded operators $\{R\}$, acting from H_2 to H_1 , such that for any $f \in A(D_A)$ one can indicate $x \in A^{-1}f$ such that for any $\varepsilon > 0$ there exists $\delta = \delta(f, \varepsilon, A)$ and an operator R from our family such that $\|R\tilde{f} - x\| \leq \varepsilon$, provided only that $\|\tilde{f} - f\| \leq \delta$.

In ⁽³⁾ an abstract analogue of the r.a. considered in ⁽²⁾ was extended to operator equations (1), under the assumption that $H_1 = H_2$ and $f \in D(A^*)$. The aim of the present work is to generalize the scheme for obtaining an r.a. proposed in our work ⁽⁴⁾ for bounded operators A , to equations with unbounded operators. The r.a. studied in ⁽¹⁻³⁾ enters into this scheme as a special case.

First of all, let us note that, by the theorem of J. von Neumann ⁽⁵⁾, there exists a self-adjoint operator A^*A with dense domain of definition $D(A^*A)$, acting from H_1 to H_1 . Denote by $\{E_\lambda\}$ its resolution of the identity.

Let the real function $\psi(\lambda, \alpha)$ possess the following properties: it is defined for $\lambda \in S(A^*A)$ and $\alpha > 0$ as follows:

$$\psi(\lambda, \alpha) = \begin{cases} \varphi(\lambda, \alpha)/\lambda, & \lambda \neq 0, \\ K, & |K| < \infty, \quad \lambda = 0. \end{cases} \quad (2)$$

$\varphi(\lambda, \alpha)$ is bounded with respect to λ and α , measurable in λ for each α with respect to $\{E_\lambda\}$; moreover,

- a) $\lim_{\alpha \rightarrow 0} \varphi(\lambda, \alpha) = 1, \quad \varphi(0, \alpha) = 0;$
- b) $\sup_{\substack{\lambda \in S(A^*A) \\ \lambda \neq 0}} |\varphi(\lambda, \alpha)|/\sqrt{\lambda} = K_\alpha < \infty \quad (\alpha > 0),$

where in a) the convergence to the limit is uniform on the set $S(A^*A) \cap (c, \infty)$, where c is any positive number.

Denote

$$\Omega_k = \{\lambda : k-1 \leq \psi(\lambda, \alpha) < k\}, \quad k = \dots, -2, -1, 0, 1, \dots$$

By our assumptions, every set Ω_k is measurable with respect to $\{E_\lambda\}$.

Let $E(\Omega_k)$ be its spectral measure. Define the operator $B_{k\alpha}$ by the formula

$$B_{k\alpha} = \overline{\int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda E(\Omega_k) A^*}. \quad (3)$$

(The bar denotes the closure of an operator. The possibility of taking the closure will be shown below.)

The operator $B_{k\alpha}$ is bounded and $\|B_{k\alpha}\| \leq K_\alpha$. Indeed, the operator

$$\int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda E(\Omega_k) A^*$$

is defined on an everywhere dense set in H_2 and has an adjoint given by the formula

$$AE(\Omega_k) \int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda = A \int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda E(\Omega_k). \quad (4)$$

Moreover, on a dense set in H_1 the equality

$$\left(AE(\Omega_k) \int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda x, AE(\Omega_k) \int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda x \right) = \left(\int_{\Omega_k} \lambda \psi^2(\lambda, \alpha) dE_\lambda x, x \right) \quad (5)$$

holds.

Indeed, if $x \in \overline{H}_{1k} = E(\Omega_k)H_1$, then equality (5) is valid (both sides are equal to 0). If $x \in H_{1k}$, then the operator

$$\int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda$$

is invertible in the subspace H_{1k} ($k \neq 0, 1$). In the case $k = 0$ (or 1) the space H_{1k} itself can be decomposed into a direct sum of its invariant subspaces with respect to

$$\int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda,$$

in each of which the operator

$$\int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda$$

is invertible. Consequently, since $D(A^*A)$ is dense in H_{1k} , there exists a dense set in H_{1k} (respectively, in each of the subspaces forming H_{1k}) on which the operator

$$\int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda E(\Omega_k) A^* A E(\Omega_k) \int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda = \int_{\Omega_k} \lambda \psi^2(\lambda, \alpha) dE_\lambda$$

is defined.

From (5) and the closedness of the operator (4) it follows that (4) is bounded; moreover, by condition b) its norm is $\leq K_\alpha$. The adjoint of (4) is equal to $B_{k\alpha}$. Therefore, $\|B_{k\alpha}\| \leq K_\alpha$. Finally, note that the ranges of the operators $B_{k\alpha}$ for different k are orthogonal. Define the operator

$$B_\alpha = \sum_{k=-\infty}^{\infty} \int_{\Omega_k} \psi(\lambda, \alpha) dE_\lambda E(\Omega_k) A^*. \quad (6)$$

By what was said above, the operator (6) is bounded. It is easy to show that the exact equality $\|B_\alpha\| = K_\alpha$ holds.

We shall now prove that the family $\{B_\alpha\}$ forms a r.a. for equation (1). Let $f \in A(D_A)$. Take such a solution $\hat{x}$ of (1) for which the condition $\hat{x} \perp \ker A^*A$

is satisfied. This means that the function $(E_\lambda \hat{x}, \hat{x})$ is continuous at $\lambda = 0$. Note that

$$B_\alpha A \hat{x} = \int_0^\infty \varphi(\lambda, \alpha) dE_\lambda \hat{x}.$$

Consider $B_\alpha \tilde{f} - \hat{x}$. We have

$$\begin{aligned} \|B_\alpha \tilde{f} - \hat{x}\| &\leq \|B_\alpha A \hat{x} - \hat{x}\| + K_\alpha \|\tilde{f} - f\| = \\ &= \left(\int_0^\infty (\varphi(\lambda, \alpha) - 1)^2 d(E_\lambda \hat{x}, \hat{x}) \right)^{1/2} + K_\alpha \|\tilde{f} - f\|. \end{aligned} \quad (7)$$

By virtue of condition a) and the continuity of $(E_\lambda \hat{x}, \hat{x})$ at $\lambda = 0$, for any $\varepsilon > 0$ one can choose such an α (and consequently also B_α) and such a δ that $\|B_\alpha \tilde{f} - \hat{x}\| \leq \varepsilon$, provided only that $\|\tilde{f} - f\| \leq \delta$. All the specific functions $\psi(\lambda, \alpha)$ (with the exception of $\psi(\lambda, \alpha) = [1 - (1 - \mu\lambda)^{1/\alpha}]/\lambda$ for $\lambda \neq 0$; $\psi(\lambda, \alpha) = \mu/\lambda$ for $\lambda = 0$), considered in ⁽⁴⁾, are suitable for constructing r.a. in our case. In particular, for (1) one can also construct an "optimal" r.a. ⁽⁴⁾. In the case when $H_1 = H_2$ and A is self-adjoint, the scheme for obtaining r.a. can be substantially simplified. The results of ⁽⁴⁾ carry over to equation (1) in this case without any changes.

An analogous scheme for obtaining r.a. in this special case was also proposed in ⁽⁶⁾.

The r.a. obtained above may be applied, for example, to constructing an approximate solution of unstable boundary-value problems for differential equations under the condition that such a problem is solvable, and in other analogous cases.

Moscow State University
named after M. V. Lomonosov

Received
1 III 1968

REFERENCES

- ¹ A. N. Tikhonov, DAN, 151, No. 3, 501 (1963).
- ² A. N. Tikhonov, DAN, 153, No. 1, 49 (1963).
- ³ V. P. Maslov, *Perturbation Theory and Asymptotic Methods*, Moscow, 1965.
- ⁴ A. B. Bakushinskii, *Zh. Vychisl. Mat. i Mat. Fiz.*, 7, No. 3, 672 (1967).
- ⁵ M. A. Naimark, *Normed Rings*, Moscow, 1966.
- ⁶ Yu. T. Antokhin, *Differential Equations*, 3, No. 7, 1135 (1967).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.