

ON THE QUESTION OF POINTS OF NONANALYTICITY OF THE GREEN FUNCTIONS OF A SYSTEM OF INTERACTING PARTICLES

L. Ya. Kobelev

1968

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196801.99433>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 536.75

PHYSICS

L. Ya. KOBELEV

ON THE QUESTION OF POINTS OF NONANALYTICITY OF THE GREEN FUNCTIONS OF A SYSTEM OF INTERACTING PARTICLES

(Presented by Academician N. N. Bogolyubov, 10 VI 1967)

1. Phase transitions (including those of the second kind) in nonrelativistic systems of interacting particles (see, for example, ⁽¹⁻⁵⁾) testify to a nonanalytic dependence of the specific heat and the thermodynamic potential of the indicated systems on the temperature difference $\Delta T = T_0 - T$ (T_0 is the temperature of the phase transition). The nonanalytic dependence on ΔT may be a consequence of the nonanalyticity of the exact one-particle (two-particle, etc.) thermodynamic Green functions of the system, defined by the corresponding equations with variational derivatives (Schwinger equations). It is shown below that such a dependence does indeed exist and can be extracted from the exact Green functions of the system, written with the aid of functional integrals over "external fields" ⁽⁶⁾ and "classical" trajectories ⁽⁷⁾, using a variant of perturbation theory consisting in expanding the functional integrals (without restrictions on the interaction parameter or the density of the number of particles) in a series in their deviation from Gaussian integrals over "external fields." It is also shown that near the points of nonanalyticity, i.e., as $\Delta T \rightarrow 0$, the "non-Gaussian" corrections to the Green function are small.

2. Consider a system of N interacting nonrelativistic particles (Bose or Fermi particles), described by the creation and annihilation operators from the ground state of particles and field quanta of the interaction $\Psi(x)$, $\Psi^*(x)$, and $A(x)$, respectively, and by the Hamiltonian $H = \Delta/2m + A(x)$ ($\hbar = c = 1$). We define the interaction energy of two particles of the system by the relation (g is the interaction constant)

$$D_0(x) = g^2 \sum_n \int dp \frac{e^{ipx}}{L(p, w_n)},$$

where $L(p, w_n)$ characterizes the type of interaction and satisfies the condition $L(0) < \infty$. Following ^(6,8,9), it is not difficult to show that the exact solution of

the thermodynamic Schwinger equation ($t = -i\beta$; $\beta = 1/kT$; $x = \mathbf{x}, t$)

$$\left[-i\frac{\partial}{\partial t} - \frac{\Delta}{2m} - \langle A(x) \rangle + ig\frac{\delta}{\delta I_0(x)} \right] G(x, x') = \delta(x - x'),$$

$$L(\Delta, \partial/\partial t) \langle A(x) \rangle = I_0(x) + igG(x, x) \quad (1)$$

for the thermodynamic one-particle causal Green function $G(x, x') = \langle T(\Psi(x)\Psi^*(x')) \rangle$ ($\langle \dots \rangle$ denotes averaging over the statistical ensemble), which describes the behavior of the system under consideration, has the form

$$G(x, x) = \frac{1}{\langle S \rangle} \int dA(x) \exp \left\{ -\frac{i}{2} \int A(x) D_0^{-1}(x - y) A(y) dx dy - \int_0^1 d\lambda \int G(x, x | \lambda A) A(x) dx \right\} G(x, x' | A); \quad (2)$$

$$\langle S \rangle = \int dA(x) \exp \left\{ -\frac{i}{2} \int A(x) D_0^{-1}(x - y) A(y) dx dy - \int_0^1 d\lambda \int G(x, x | \lambda A) A(x) dx \right\}; \quad (3)$$

$$[-i\partial/\partial t - \Delta/2m - \lambda A(x)] G(x, x' | \lambda A) = \delta(x - x'); \quad (4)$$

$$[-i\partial/\partial t - (\Delta - \Delta')/2m - \lambda(A(x) - A^*(x'))] G(x, x') = 0, \quad x' \rightarrow x. \quad (5)$$

In (2)–(3) the continual integral over $A(x)$ of the functional $\varphi(x|A)$ (the functional dependence is separated by a vertical bar) is defined as the limit of an n -fold integral of a function of n variables $\varphi(x; A_1, \dots, A_n)$ as $n \rightarrow \infty$, $A_{i+1} - A_i \rightarrow 0$, $\Delta x_i \rightarrow 0$.

Integration over x_4 and variational derivatives with respect to $I_0(x)$, $\langle A(x) \rangle$, etc., differ from zero only for $0 \leq ix_4 \leq 1/T$; the function $\delta(x_4)$ is defined at discrete “time” points ⁽¹⁰⁾. For retarded, advanced, commutator, etc., thermodynamic Green functions, the representation (2)–(4) remains valid when equation (4) for $G(x, x' | A)$ is replaced by the equation for the retarded, advanced, etc., function $G(x, x' | A)$. The “external source” $I_0(x)$ in (2)–(3) is equal to zero.

3. Let us now represent $G(x, x' | A)$ and $G(x, x | A)$ through continual integrals over all continuous trajectories $\mathbf{x}(\nu)$, beginning at “time” $\nu = t'$ at the point $\mathbf{x}'$ and ending at “time” t at the point $\mathbf{x}$ ^(11,12):

$$\begin{aligned}
G(x, x'|A) &= \int_{G_{x, x'}} d_G \mathbf{x}(\nu) \exp \left\{ i \int_{t'}^t A(x(\nu)) d\nu \right\} \equiv \\
&\equiv \lim_{\substack{\Delta \nu = \nu_{i+1} - \nu_i \rightarrow 0 \\ \Delta A = A(x_{i+1}) - A(x_i) \rightarrow 0 \\ \Delta x = x_{i+1} - x_i \rightarrow 0 \\ n \rightarrow \infty}} \int dx_1 \dots \int dx_n G_0(\mathbf{x} - \mathbf{x}_1; t - \nu_1) \dots G_0(\mathbf{x}_n - \mathbf{x}'; \nu_n - t') \times \\
&\times \exp \left\{ i \sum_{i=1}^n A(\mathbf{x}_i)(\nu_{i+1} - \nu_i) \right\}, \quad it' < it, \quad t = -i\beta;
\end{aligned} \tag{6}$$

$$(-i\partial/\partial t - \Delta/2m)G_0(x, x') = \delta(x - x'); \tag{7}$$

$$G(x, x) = \lim_{x' \rightarrow x} \int_{G_{x, x'}} d_G \mathbf{x}(\nu) \exp \left\{ i \int_{x_4^0}^{x_4} [dA_1(x(\nu)) + 2iA_2(x(\nu))] d\nu \right\}. \tag{8}$$

Here x_4^0 is an arbitrary parameter, which in what follows, for convenience, is taken equal to x_4 ; $dA_1(x) = A_1(x + \Delta x) - A_1(x) + d_{A_1}A_1(x)$ ($d_{A_1}A_1(x)$ is the variation of $A_1(x)$ at the points x); $A = \text{Re } A + i \text{Im } A = A_1 + iA_2$. In the last integrals in (6)–(8) one should replace the solutions (7) $G_0(p)$ by $G_0(p)n(p)$. Taking (6) into account, $G(x, x')$ takes the form ($G(x, x') = 0$ for $ix'_4 > ix_4$)

$$\begin{aligned}
G(x, x') &= \frac{1}{\langle S \rangle} \int dA(x) \int_{G_{x, x'}} d_G \mathbf{x}(\nu) \exp \left\{ -\frac{i}{2} \int A(x) D_0^{-1}(x - y) A(y) dx dy \right. \\
&\quad \left. - \int_0^1 d\lambda \int G(x, x|\lambda A) A(x) dx + i \int_{x_4^0}^{x_4} A(x(\nu)) d\nu \right\}, \quad ix'_4 < ix_4.
\end{aligned} \tag{9}$$

4. Let us isolate in (9) the “Gaussian” part in $A(x)$ by means of a nonlinear transformation of $A(x)$ of the form

$$A(x) = A'(x) + \int D_0(x - y) \tilde{F}(y) dy,$$

where

$$\tilde{F}(y) = F(y) + \varepsilon(x(\nu)) = i \int_0^1 d\lambda G(y|\lambda A) + \int_{x_4^0}^{x_4} d\nu \delta(x(\nu) - y) \delta(\nu - y^0).$$

This transformation eliminates from the exponent (9) the second and third terms and brings (9) to the form

$$G(x, x') = \frac{1}{\langle S \rangle} \int dA'(x) \int_{G_{x, x'}} d_G \mathbf{x}(\nu) D(|A|) \exp \left\{ -\frac{i}{2} \int A'(x) D_0^{-1}(x-y) \times \right. \\ \left. \times A'(y) dx dy + \frac{i}{2} \int \tilde{F}(x) D_0(x-y) \tilde{F}(y) dx dy \right\}, \quad (10)$$

where $D(|A|)$ is the Jacobian of the transformation, coinciding with the Fredholm determinant of the integral equation (see (13)), $A(x) = A'(x) + \int K(x, y) A(y) dy$, for $K(x, y) = \int D_0(x-y_1) \frac{\delta F(y_1)}{\delta A(y)} dy_1$.

We shall show that $D(|A|)$ contains a nonanalytic factor due to the dependence of $G(x, x|A)$ on the real part A . Considering the transformation only for A_1 and using the relations $G(x, y) - G(y, x) = \delta_{x, y}$, $\delta F(x)/\delta A_1(y) = \delta_{y, x} \cdot \tilde{\Pi}_\lambda(0) = \delta_{y, x} \cdot \Pi_\lambda(0)(1 - \Pi_\lambda(0)D_0(0))^{-1}$, where

$$\Pi_\lambda(0) = -\frac{T}{(2\pi)^2} \sum_{w_n} \int dp \int_0^1 \lambda d\lambda G^2(p, w_n | \lambda A), \quad D_0(0) = g^2 L^{-1}(0) \quad (11)$$

and $G(p, w_n | A)$ satisfies (5), and restricting ourselves to the study of the case $F(x)$ independent of x (the analogue of a Bose-particle condensate), we find, separating from $\Pi_\lambda(0|A)$ the term with $d_{A_1} A_1(x)$ in the linear approximation ($\tilde{D}_0(0) = D_0(0)(1 - \Pi_\lambda(0)D_0(0))^{-1}$):

$$D(|A) = D(|A_2)(1 - \Pi_\lambda(0)\tilde{D}_0(0))^{-1} \exp \left\{ \frac{\Pi_\lambda(0)D_0(0)}{1 - 2\Pi_\lambda(0)D_0(0)} d_A A_1 dv \right\}, \\ \eta = \pm 1. \quad (12)$$

The appearance of the nonanalytic term is due to the retention in $G(x, x|A)$ of first-order terms with respect to the variation of A_1 . The ambiguity of the integral over A_1 is removed by the choice of the sign of $d_A A_1$ (determined by η) from the normalization condition of $G(x, x')$.

5. To eliminate the remaining linear dependence on $dA_1(x)$ (and $A_2(x)$), we separate in $D(|A)$ and in one of the functionals $F(x)$ in the expression $\int \tilde{F}(x) D_0 F(y) dx dy$ the terms linear in A_1 and A_2 , and eliminate them by means of the transformation

$$A'(x) = A''(x) + \int D_1(x-y)F_1(y)dy, \quad (*)$$

where $F_1(x)$ is a functional of A_1 (and A_2), chosen from the condition of compensation of the linear terms (F_1 has the form $F_1 = \int F D_0 \Pi_\lambda dx dy - \int_{x_0^4}^{x_4'} D_0(x(y)-y) \Pi_\lambda(y-x) dy dx dy$). In this case, in (10) there will appear the Jacobian $D'_1(|A'')$ and the factor $\exp \left\{ \frac{i}{2} \int F_1(x) D_1(x-y) F_1(y) dx dy \right\}$. Omitting the intermediate calculations, we give the expression for $G(x, x')$ after application of an infinite number of transformations of the form (*) and complete elimination from the subintegral expression (10) of the terms linear in $A_1(x)$ (it is not difficult to show that elimination of the terms linear in $A_2(x)$ does not lead to the appearance of new nonanalyticities):

$$G(x, x') = \langle S \rangle^{-1} \int dA^\infty(x) \int_{G_{x,x'}} d_G \mathbf{x}(y) D(|A|) \prod_{i \geq 1}^\infty D_i(|A^\infty|) \times \\ \times \exp \left\{ -\frac{i}{2} \int A^\infty D_\infty^{-1}(x-y) A^\infty dx dy + \frac{i}{2} \int \left[F(x) D_0(x-y) F(y) + \sum_{i=1}^\infty F_i(x) D_i(x-y) F_i(y) \right] dx dy \right\}, \quad (13)$$

where $A^\infty = A^i|_{i=\infty}$; $D_i(x)$ is the function $D_0(x)$ renormalized by the inclusion of terms $\sim AA$ at each of the transformations; F_i is a functional of A^i , eliminating the linear terms in the $(i+1)$ -st transformation; $D_i(|A^i|)$ are the Jacobians of the $(i+1)$ -st transformation. For $i = \infty$ the quantities F

and F_i , and also the variational derivatives of F and F_i with respect to A^i , do not contain terms linear and quadratic in $A(x)$. The parts $D_i(|A|)$ that depend on A_1^i have the symbolic form

$$D_i(|A_1^i|) = \exp \left\{ \ln \left[1 + D_i(0) \frac{\delta F_i}{\delta A^i} (1 + d_{A_1} A_1 dv \eta) \right] \right\}, \quad (14)$$

$$D_i(0) \frac{\delta F_i}{\delta A^i} = \frac{a_i}{1 - b_i} = D_i(0) \frac{\delta A F_i}{\delta A^i} \left[1 - \int_{x_0^4}^{x_4} D_i(x(\nu) - y') \frac{\delta F_i(y')}{\delta F_i(y)} dy' dy d\nu \right]^{-1},$$

where $\delta F_i / \delta A^i$ and $\delta A F_i / \delta A^i$ are the total and “partial” (with respect to the explicit dependence on A) variational derivatives. In computing F_i and D_i one must take into account that $\delta_F F_i / \delta F_i$ no longer depends on A_1 , and the dependence on A_1 remains only in $\int F_i D_i F_i dx dy$ and in $\delta A F_i / \delta A^i$. Each of the Jacobians $D_i(|A_1^i|)$ has a point of nonanalyticity determined by the relation $1 - b_{iT} = 0$.

6. The perturbation-theory method used below consists in separating out in (13) (when representing $\langle S \rangle$ by means of analytic transformations A' , A'' , $(*)$ etc.) the Gaussian quadratic form in $A_1(x)$ (i.e., in separating out in the exponent the terms proportional to A_1^∞) and expanding the remaining terms in powers of A_1^∞ . Taking account of the normalization

$$\lim_{j \rightarrow \infty, dA_1 \rightarrow 0, \rho x \rightarrow 0} \sum d_{A_1} A_1(x_j) d\nu = 1,$$

we then obtain ($G(x, x') = G_\Gamma(x, x') + G_{nG}(x, x')$)

$$G_\Gamma(x, x') = \prod_{i \geq 0} \left(\frac{D_\infty}{D_\infty^s} \right)^{1/2} \left(1 + \frac{a_i}{1 - b_i} \right) \left(1 + \frac{a_i^s}{1 - b_i^s} \right)^{-1} \times \\ \times \exp \left\{ \frac{a_i}{1 - b_i} - \frac{a_i^s}{1 - b_i^s} + \frac{i}{2} \int (F_i^0 D_i F_i^0 - F_i^{0s} D_i^s F_i^{0s}) dx dy \right\} G_\Gamma^0(x, x'). \quad (15)$$

In (15), $a_0 = \widetilde{D}_0(0)\Pi_\lambda(0)$; F_i^0 does not depend on $x(\nu)$; $b_0 = \widetilde{D}_0(0)\Pi_\lambda(0)$; the index s means that the quantity was obtained under the transformation $\langle S \rangle$;

$$D_i^{-1} = \widetilde{D}_i^{-1}(0) + \sum_{n=0}^i \frac{\delta^2 a_n}{\delta A^i \delta A^i} (1 - b_n)^{-1} \simeq D_0^{-1}(0) + \sum_{n=0}^i \frac{\tilde{a}_n}{1 - b_n};$$

$$F_j \simeq \frac{i}{2} \int F_{j-1} D_j \frac{\delta F_{j-1}}{\delta A^j} dx dy - \frac{\delta a_j}{\delta A^j} (1 - b_j)^{-1}; \quad F_0 \equiv \widetilde{F} \quad (i, j = 1, 2, \dots, \infty).$$

The function G_Γ^0 satisfies an approximate Dyson equation. The non-Gaussian part of the Green function G_{nG} near $1 - b_i \simeq \Delta T_i/T \rightarrow 0$ has order $(1 - b_i)^2 \tilde{a}_i^{-2} G_\Gamma$, since $D_\infty^{-1} \sim \tilde{a}_{iT}/\Delta T_i$; therefore, near the points of nonanalyticity, $G \simeq G_\Gamma + O(G_\Gamma \Delta T_i^2/T^2)$. It can be shown that the presence of nonanalytic singularities in ΔT_i in the thermodynamic potential Ω of the particle system, continuous at the point T_i , leads to $\partial^2 \Omega / \partial T^2 \sim \ln(T/\Delta T_i)$, which makes it possible to regard the smallest of the T_i as the temperature of a second-order phase transition. In this case the dependence of $\partial \Omega / \partial \mu$ (μ is the chemical potential) on ΔT contains both an analytic and a nonanalytic term.

Ural State University
named after A. M. Gorky

Received
25 IV 1967

REFERENCES

1. M. I. Buckingham, W. H. Fairbank, *Progress in Low Temperature Physics*, **3**, 1963, p. 80.
2. V. L. Ginzburg, *FTT*, **11**, 2031 (1961).
3. A. B. Voronel, Yu. R. Chashkin, *ZhETF*, **51**, 394 (1966).
4. L. Onsager, *Phys. Rev.*, **65**, 117 (1944).
5. G. Domb, *Adv. Phys.*, **9**, 149 (1960).
6. N. N. Bogolyubov, *DAN*, **99**, 225 (1954).
7. L. Ya. Kobelev, *Izv. vyssh. uchebn. zaved., Fizika*, **4**, 206 (1960).
8. E. S. Fradkin, *DAN*, **98**, 47 (1954).
9. S. E. Edwards, R. E. Pais, *Proc. Roy. Soc.*, **224**, 24 (1954).
10. E. S. Fradkin, *ZhETF*, **36**, 957, 1286 (1954); *DAN*, **125**, 311 (1959); *Tr. Fiz. inst. im. P. N. Lebedeva AN SSSR*, **59**, "Nauka" (1965).
11. L. Ya. Kobelev, *Tr. Vsesoyuzn. konf. po funktsionalnomu analizu*, Baku, 1960.
12. L. Ya. Kobelev, *Tr. Mezhdunarodn. simpoziuma*, Novosibirsk, 1964; *Uch. zap. Uralsk. gos. univ. im. A. M. Gorkogo, ser. fiz.*, **1**, 166 (1965).
13. S. E. Edwards, *Proc. Roy. Soc.*, **228**, 411 (1955).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.