

ON CONTINUOUS PARTITIONS OF COLLECTIVELY NORMAL SPACES

N. S. Lashnev

1969

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-196901.07194>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 513.83

MATHEMATICS

N. S. LASHNEV

ON CONTINUOUS PARTITIONS OF COLLECTIVELY NORMAL SPACES

(Presented by Academician P. S. Aleksandrov on 17 I 1969)

Definition 1. Let there be a set L of cardinality τ , and let to each element λ of L there be assigned a discrete system of sets γ_λ . Let

$$\gamma = \bigcup_{\lambda \in L} \gamma_\lambda.$$

We shall say that the system γ is τ -discrete.

Definition 2. Let X be a completely regular space. We shall say that its **uniform weight is equal** to τ if there exists a uniform structure with a base of τ covers, compatible with the topology of the space X , and for any cardinal number $\sigma < \tau$ no such uniform structure exists.

Theorem. *Let X be a collectively normal space of uniform weight τ , and let $\{A\}$ be its continuous partition. Then in all elements of the partition, except for the elements of some τ -discrete system, every discrete set has cardinality $< \tau$.*

Proof. 1. Denote by $\{\alpha_\lambda\}_{\lambda \in L}$ a base of the uniform structure in X , consisting of τ covers α_λ . The system $\{\alpha_\lambda\}$ has the following properties: a) for any two covers α_{λ_1} and α_{λ_2} there exists a third, which is inscribed in their intersection $\alpha_{\lambda_3} < \alpha_{\lambda_1} \wedge \alpha_{\lambda_2}$; b) if x is a point in the space X , then the system of all stars of this point with respect to the covers α_λ forms a base at the point x .

2. To each λ from the set L we assign a discrete system of elements of the partition $\{A\}$. For this purpose denote by $V_\lambda(A)$ the star of the element A with respect to the cover α_λ . Into the system γ_λ we select those and only those elements of the partition which have the property: the star of the element of the partition with respect to the system α_λ contains wholly no element of the partition distinct from it itself,

$$A_0 \in \alpha_\lambda \iff V_\lambda(A_0) \cap A_1 \neq A_1$$

for $A_1 \neq A_0$.

The discreteness of the system α_λ follows from the continuity of the partition $\{A\}$: if $A_0 \notin \gamma_\lambda$, then for any $x \in A_0$ there exists a neighborhood Ox which does not meet the bodies G_λ of the system α_λ . Otherwise one would have $[G_x] \cap A_0 \neq \Lambda$, and by virtue of the continuity of the partition the neighborhood $V_\lambda(A_0)$ of the set A_0 would contain wholly some element $A_1 \in \alpha_\lambda$, which contradicts the construction of the system α_λ .

3. We shall prove that a discrete subset of any element of the partition which has not entered into any system α_λ has cardinality $< \tau$. Suppose this is not so: $A_0 \notin \bigcup_{\lambda \in L} \gamma_\lambda$ and $A_0 \supset P$, where P is a discrete set of cardinality τ . Since L has cardinality τ , one can establish a one-to-one correspondence between P and L ; then each point $x \in P$ acquires an index λ : $P = \{x_\lambda\}_{\lambda \in L}$.
4. Since X is collectively normal, the discrete system of points $\{x_\lambda\}$ has a discrete system of neighborhoods $\{Ox_\lambda\}$ in the space X .
5. For each point $x_\lambda \in P$ one can choose a cover $a(x_\lambda) \in \{\alpha_\lambda\}$ so that the star of this cover with respect to the point x_λ is contained in the set Ox . It may happen that the same cover α_λ corresponds to different points x_1 and x_2 of P . We have constructed a mapping φ of the set $P = \{x_\lambda\}$ into the set of covers $\{\alpha_\lambda\}_{\lambda \in L}$.
6. In item 3 a one-to-one correspondence between the sets L and P was established. It can be turned into a one-to-one mapping $\psi: \{x_\lambda\} \rightarrow \{\alpha_\lambda\}$, by assigning to each point x_λ the cover with the same index.
7. Let us construct a third mapping ω of the set P into the set of covers α_λ . Using property b) of item 1, to each point $x \in P$ we assign some cover α_x belonging to the system $\{\alpha_\lambda\}$, in such a way that, for any λ , one has $\alpha_x^\lambda < \alpha(x_\lambda) \cap \alpha_\lambda$, or $\omega(x) < \varphi(x) \cap \psi(x)$ ($x \in P$).
8. For any $x \in P$ we have:

$$St(\alpha_x, x) \subset Ox; \quad (1)$$

there exists an element A_x of the decomposition $\{A\}$ such that

$$St(\alpha_x, A_x) \supset A. \quad (2)$$

9. Since $\{\alpha_x\}_{x \in P}$ forms a base of the uniform structure, we have $\left[\bigcup_{x \in P} A_x\right] \supset A$.
10. Consider an arbitrary A_x . In view of (2), there exists an element H of the cover α_x which intersects A_x and contains the point x . From (1) it follows that $A_x \cap Ox \neq \Lambda$. Choose in the set $A_x \cap Ox$ a point $y(x)$. Obviously, the set Q of all points $y(x)$ is discrete.

11. Consider the open set $X \setminus Q$. Since the decomposition $\{A\}$ is continuous, there exists an open saturated set $U \subset X \setminus Q$ which contains A_0 . It is easy to see that $X \setminus Q$ does not contain entirely any one of the sets A_x ($x \in P$); hence, for any $x \in P$, $A_x \cap U = \Lambda$ and $(\bigcup A_x) \cap U = \Lambda$, which contradicts the fact that $[\bigcup_{x \in P} A_x] \supset A_0$. The theorem is proved.

Faculty of Mechanics and Mathematics
M. V. Lomonosov Moscow State University

Received
8 I 1969

REFERENCES

¹ A. V. Arkhangel'skii, DAN, 166, No. 6 (1966). ² N. S. Lašnev, DAN, 165, No. 4 (1965). ³ V. V. Filippov, DAN, 178, No. 3 (1968).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.