

ON THE THEORY OF LINEAR PASSIVE SYSTEMS

V. S. Vladimirov

1969

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196901.44525>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.9.004.14

MATHEMATICS

Corresponding Member of the Academy of Sciences of the USSR V. S. VLADIMIROV

ON THE THEORY OF LINEAR PASSIVE SYSTEMS

1. Introduction. By a **linear passive system** we shall mean a system of convolution equations

$$Z * u = f, \quad (1)$$

where $f = (f_1, \dots, f_N)$ is a given vector distribution from $(\mathcal{D}')^{\times N} = \mathcal{D}' \times \dots \times \mathcal{D}'$, $u = (u_1, \dots, u_N)$ is an unknown element of $(\mathcal{D}')^{\times N}$, and $Z = \|Z_{kj}\|$ is an $N \times N$ matrix with material elements $Z_{kj}(\xi)$ from $\mathcal{D}' = \mathcal{D}'(\mathbb{R}^n)$, satisfying the conditions:

- a) **causality** with respect to some convex closed acute cone $\Gamma \subset \mathbb{R}^n$ with vertex at O : $Z(\xi) = 0$, $\xi \notin \Gamma$;
- b) **passivity**: for all complex $a \in \mathbb{C}^N$, $\varphi \in \mathcal{D}$, and $q \in C = \text{int } \Gamma^*$, the inequality

$$\text{Re} \int [(Ze^{-(q, \xi)} a, a) * \varphi](x) \overline{\varphi}(x) \geq 0; \quad (2)$$

holds; here Γ^* is the cone conjugate to the cone Γ .

The operator $Z*$, defining the linear passive system (1), will be called a **passive operator** (with respect to the cone Γ), and the corresponding matrix $Z(\xi)$ the (generalized) **impedance** of the system.

A number of problems of mathematical physics reduce to linear passive systems: linear thermodynamic systems, the theory of electric circuits, scattering of nuclear particles, scattering of electromagnetic waves, etc. ⁽¹⁻⁸⁾. Inequality (2) reflects the ability of a physical system to absorb energy, but not to generate it.

Linear passive systems have been studied in especially great detail in the case of one independent variable ⁽¹⁻⁴⁾. The principal apparatus of investigation in this case is the classical integral representation of functions holomorphic in the upper half-plane and having there nonnegative real part ⁽⁹⁾. Using the results of

(¹⁰) on the representation of holomorphic functions of many complex variables with nonnegative real part in a tubular domain over a cone, we have investigated linear passive systems in the multidimensional case. Here we present the most important results of these investigations.

2. Characteristic of a passive operator. From conditions a) and b) it follows that $Z_{kj} \in \mathcal{S}'(\Gamma)$, so that the matrix $Z(\xi)$ has a Fourier-Laplace transform (¹¹).

Theorem 1. In order that an $N \times N$ matrix $Z(\xi)$ define a passive operator with respect to the cone Γ , it is necessary and sufficient that its Fourier-Laplace transform $\tilde{Z}(\zeta)$, $\zeta = p + iq$, be a holomorphic matrix-function in the tubular domain $T^C = \mathbb{R}^n + iC$, $C = \text{int } \Gamma^*$, satisfying the relation

$$\tilde{Z}(p + iq) = \overline{\tilde{Z}(-p + iq)}, \quad \zeta \in T^C, \quad (3)$$

and the inequality

$$\text{Re}(\tilde{Z}(\zeta)a, a) \geq 0, \quad a \in \mathbb{C}^N, \quad \zeta \in T^C. \quad (4)$$

Remark. The sufficient conditions in Theorem 1 admit a weakening: if the matrix $\tilde{Z}(\zeta)$ is holomorphic in the tube domain T^C (the cone C may also be nonconvex), satisfies inequality (4) in T^C , and the matrix $\tilde{Z}(iq)$ is real for all $q \in C$, then the matrix $Z(\xi)$ defines a passive operator with respect to the cone C^* .

Theorem 2. In order that an $N \times N$ -matrix $Z(\xi)$ define a passive operator with respect to the cone Γ , it is necessary and sufficient that, for any cone

$$C' = [q : (e_j, q) > 0, j = 1, \dots, n] \subset C = \text{int } \Gamma^*$$

it be representable (uniquely) in the form *

$$Z(\xi) = (e_1, D)^2 \dots (e_n, D)^2 Z_{C'}(\xi) + \sum_{j=1}^n Z_{C'}^{(j)} D_j \delta(\xi), \quad (5)$$

where $Z_{C'}(\xi)$ is a real continuous matrix of polynomial growth in $\mathbb{R}^n$ with support in the cone $(C')^*$, and such that for any $a \in \mathbb{C}^N$ the generalized function **

$$(e_1, D)^2 \dots (e_n, D)^2 ([Z_{C'}(\xi) + Z_{C'}^T(-\xi)]a, a)$$

is positive-definite in the sense of Bochner-Schwartz; the matrices $Z_{C'}^{(j)}$, $j = 1, \dots, n$, are real symmetric and such that the matrix

$$\sum_{j=1}^n q_j Z_{C'}^{(j)} \geq 0$$

for all $q \in C'$.

Remark. For $n = 1$, Theorem 2 was proved by König and Zemanian ⁽⁴⁾.

Corollary. In order that a function $f(\zeta)$ be holomorphic in T^C , $\text{Im } f(\zeta) \geq 0$, $\zeta \in T^C$, and

$$\lim_{q \rightarrow 0} \text{Im } f(p + iq) = 0, \quad q \in C \text{ in } \mathcal{S}',$$

it is necessary and sufficient that it be representable in the form $f(\zeta) = (a, \zeta)$, where $a \in C^*$ (a theorem of the type of the Liouville and Phragmén-Lindelöf theorems).

3. Multidimensional dispersion relations

Theorem 3. In order that an $N \times N$ -matrix $Z(\xi)$ define a passive operator with respect to the cone Γ , it is necessary and sufficient that its Fourier transform $\tilde{Z}(p)$, for any cone $C' \subset C = \text{int } \Gamma^*$ (see Theorem 2), satisfy the dispersion relation

$$\tilde{Z}(p) = \frac{2(-1)^n}{(2\pi)^n} (e_1, p)^2 \dots (e_n, p)^2 (\tilde{M}_{C'} * K_{C'}) - i \sum_{j=1}^n Z_{C'}^{(j)} p_j; \quad (6)$$

where

$$K_{C'}(p) = \det \|e_{jk}\| \left\{ -i\pi\delta[(e_1, p)] + P \frac{1}{(e_1, p)} \right\} \times \dots \\ \dots \times \left\{ -i\pi\delta[(e_n, p)] + P \frac{1}{(e_n, p)} \right\};$$

$\tilde{M}_{C'}(p) = F[M_{C'}(\xi)]$ is a Hermitian matrix with the properties: I. The matrix $M_{C'}(\xi)$ is real continuous of polynomial growth with support in the cone $(C')^* \cup (-C')^*$. II. For any $a \in \mathbb{C}^N$ the generalized function

$$\frac{(-1)^n (e_1, p)^2 \dots (e_n, p)^2}{[1 + (e_1, p)^2] \dots [1 + (e_n, p)^2]} (\tilde{M}_{C'}(p)a, a)$$

is a nonnegative finite measure on $\mathbb{R}^n$; $Z_{C'}^{(j)}$, $j = 1, \dots, n$, are real symmetric matrices such that for all $q \in C'$

* $D = (D_1, \dots, D_n) = (\partial/\partial x_1, \dots, \partial/\partial x_n)$.

** Z^T is the matrix transposed to the matrix Z .

$$\sum_{j=1}^n q_j Z_C^{(j)} \geq 0.$$

In this case the equality holds

$$M_{C'}(\xi) = \frac{1}{2} [Z_{C'}(\xi) + Z_{C'}^T(-\xi)]. \quad (7)$$

Remark. For $n = 1$, Theorem 3 was proved by Beltrami and Wohlers ⁽³⁾.

4. Existence of a fundamental solution. A **fundamental solution** of the passive operator $Z*$ is any matrix $A(\xi)$, $A_{kj} \in \mathcal{D}'$, satisfying the matrix convolution equation

$$Z * A = I\delta(\xi). \quad (8)$$

The operator $A*$ is also called the **inverse operator** to the operator $Z*$, and the matrix $A(\xi)$ the (generalized) **admittance** of the system. A passive operator $Z*$ is called **nondegenerate** (**completely nondegenerate**) if $\det Z(\zeta) \neq 0$, $\zeta \in T^C$ (respectively, for every $a \in C^N$, $a \neq 0$, there exists a point $\zeta_0 \in T^C$ such that $\operatorname{Re}(Z(\zeta_0)a, a) > 0$).

In order that the passive operator $Z*$ be completely nondegenerate, it is necessary and sufficient that

$$(Z(\xi)a, a) \neq ig^0(\xi), \quad a \in C^N, \quad a \neq 0, \quad \operatorname{Im} g = 0. \quad (9)$$

The following is the main result.

Theorem 4. *In order that the operator $Z*$, $Z_{kj} \in \mathcal{D}'(\Gamma)$, be a (completely) nondegenerate passive operator with respect to the cone Γ , it is necessary and sufficient that there exist a unique inverse operator $A*$ in the class of (completely) nondegenerate passive operators with respect to Γ .*

5. Examples of passive systems. We first note a sufficient condition for passivity. Let a real $N \times N$ matrix $Z(\xi)$ satisfy the causality condition a), item 1, and the condition

$$\text{b')} \quad \operatorname{Re} \int_{-\Gamma} [((Za, a) * \varphi)(x) \bar{\varphi}(x)] dx \geq 0, \quad \varphi \in \mathcal{D}, \quad a \in C^N. \quad (10)$$

(We assume that the cone Γ satisfies the conditions of item 1 and, moreover, is solid.) Then the matrix $Z(\xi)$ defines a passive operator with respect to the cone Γ , i.e., it satisfies condition b), item 1.

- 1) If a system of differential equations with constant coefficients is passive, then it is of first order.
- 2) In order that the differential operator $[(l, D) + c]\delta(\xi)_*$ with constant coefficients be passive with respect to the cone Γ , it is necessary and sufficient that $l \in \Gamma$ and $c \geq 0$.
- 3) In order that a system of linear differential equations with real constant coefficients

$$\sum_{k=1}^n A_k \frac{\partial u}{\partial x_k} = f, \quad (11)$$

be passive and completely nondegenerate, it is necessary and sufficient that the $N \times N$ matrices A_k , $k = 1, \dots, n$, be symmetric and that there exist such a vector l that

$$\sum_{k=1}^n l_k A_k > 0.$$

In this case the passivity of the system (11) holds with respect to the convex hull of the cone

$$[\xi : \xi_1 = (A_1 a, a), \dots, \xi_n = (A_n a, a), \quad a \in R^N].$$

Remark. Systems (11) satisfying the conditions of assertion 3) are the principal parts of systems symmetric in the sense of Friedrichs ⁽¹³⁾ with constant coefficients.

- 4) In order that the difference operator

$$B_0 \delta(\xi)_* + \sum_{\nu=1}^m B_\nu \delta(\xi - h_\nu)_*,$$

where B_k , $k = 0, 1, \dots, m$, are real $N \times N$ matrices, be passive, it is necessary and sufficient that: I. The smallest closed convex cone Γ containing the points $\{0, h_\nu, \nu = 1, \dots, m\}$ be acute. II. For all $\zeta = p + iq \in T^C$, $C = \text{int } \Gamma^*$, the matrix

$$B_0 + B_0^T + \sum_{\nu=1}^m e^{-(q, h_\nu)} \left[\cos(p, h_\nu) (B_\nu + B_\nu^T) - \sin(p, h_\nu) \frac{B_\nu - B_\nu^T}{i} \right] \geq 0.$$

In this case passivity holds with respect to the cone Γ .

Correction note. It can be proved that every matrix $Z(\xi)$ defining a passive operator with respect to a solid cone Γ also satisfies condition b' .

Mathematical Institute named after V. A. Steklov
Academy of Sciences of the USSR

Received
13 III 1969

REFERENCES

1. H. König, A. H. Zemanian, SIAM, J. Appl. Math., **13**, 1036 (1965).
2. H. König, T. Meixner, Math. Nachr., **19**, 265 (1958).
3. E. T. Beltrami, M. R. Wohlers, *Distributions and the Boundary Values of Analytic Functions*, N.Y.—London, 1966.
4. A. H. Zemanian, *Distribution Theory and Transform Analysis*, 1966.
5. W. Güttinger, SIAM, J. Appl. Math., **15**, 964 (1967).
6. V. P. Silin, A. A. Rukhadze, *Electromagnetic Properties of Plasma and Plasma-like Media*, 1961.
7. M. A. Leontovich, ZhETF, **40**, 907 (1961).
8. T. T. Wu, J. Mathem. Phys., **3**, 262 (1962).
9. R. Nevanlinna, *Eindeutige Analytische Funktionen*, 1936.
10. V. S. Vladimirov, Mat. Sb., **79** (121), No. 1 (5), 131 (1969).
11. V. S. Vladimirov, *Methods of the Theory of Functions of Several Complex Variables*, "Nauka," 1964.
12. L. Schwartz, *Théorie des distributions*, 1–2, Paris, 1950–51.
13. K. Friedrichs, Comm. Pure and Appl. Math., **7**, 345 (1954).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.