

ON THE REPRESENTATION OF SOLVABLE GROUPS BY MATRICES OVER A CERTAIN FIELD OF CHARACTERISTIC ZERO

E. M. Levich

1969

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196901.57058>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 519.44

MATHEMATICS

E. M. LEVICH

ON THE REPRESENTATION OF SOLVABLE GROUPS BY MATRICES OVER A CERTAIN FIELD OF CHARACTERISTIC ZERO

(Presented by Academician V. M. Glushkov, 18 II 1969)

One of the basic results in the theory of finite-dimensional representations of solvable groups is the theorem of A. I. Mal' tsev ⁽¹⁾: a solvable matrix group has a normal divisor of finite index whose commutant is nilpotent. In connection with this theorem the question arises: what abstract properties must a solvable group possess in order that it can be represented isomorphically by matrices over some field? As an example one may cite the following assertion: every polycyclic group is representable by matrices over the field of rational numbers ^(2, 3).

We give the definitions needed in what follows. Let G be an arbitrary locally nilpotent torsion-free group; Ω an arbitrary field of characteristic zero, and x^λ a single-valued function which assigns to any elements $x \in G$ and $\lambda \in \Omega$ a certain element of G .

G is called an Ω -powered group ⁽⁴⁾ if the following conditions are satisfied:

- 1) $x^1 = x$, $x^{\lambda+\mu} = x^\lambda x^\mu$, $x^{\lambda\mu} = (x^\lambda)^\mu$;
- 2) $y^{-1} x^\lambda y = (y^{-1} xy)^\lambda$,
- 3)

$$x_1^\lambda x_2^\lambda \dots x_n^\lambda = t_1^\lambda t_2^{\binom{\lambda}{2}} \dots t_k^{\binom{\lambda}{c}},$$

where c is the nilpotency class of the group generated by the elements $x_1, x_2, \dots, x_n$, and $t_1, t_2, \dots, t_c$ are Petresco words ⁽⁴⁾, p. 21).

In condition 1), 1 is the identity of the field Ω ; the elements x_i, x, y are arbitrary in G , and λ, μ are arbitrary elements of Ω ; finally, condition 3) is assumed to hold for every finite n .

An Ω -powered locally nilpotent group G is called an ΩR -powered group if from the equality $x^\lambda = y^\lambda$ for some $\lambda \in \Omega$ it follows that $x = y$. If n is the least number which bounds from above the minimal number of Ω -generators of each finitely generated Ω -powered subgroup of the Ω -powered group G , then we shall

say that G has Ω -rank n . The notion of an Ω -homomorphism of one Ω -powered group into another is defined in the natural way.

Theorem 1. In order that a finitely generated solvable torsion-free group Γ have a faithful matrix representation over a field Ω of characteristic zero, it is necessary and sufficient that the group Γ have the structure

$$\Gamma \supset \Gamma_1 \supset \Gamma_2 \supset \{e\},$$

where Γ/Γ_1 is a finite group, Γ_1/Γ_2 is a finitely generated abelian group, and the group Γ_2 can be embedded isomorphically in an ΩR -powered nilpotent group H of finite Ω -rank, moreover the restriction of each inner automorphism of the group Γ_1 to the subgroup Γ_2 induces an Ω -automorphism of the group H .

Corollary. In order that a finitely generated solvable torsion-free group Γ have a faithful matrix representation over the field of ra

rational numbers, it is necessary and sufficient that the group Γ have a series

$$\Gamma \supset \Gamma_1 \supset \Gamma_2 \supset \{e\},$$

where Γ/Γ_1 is a finite group, Γ_1/Γ_2 is a finitely generated abelian group, and Γ_2 is a torsion-free nilpotent group of finite rational rank.

This corollary gives an answer to the question posed in ⁽⁵⁾.

Theorem 2. In order that a torsion-free solvable group Γ with trivial center be isomorphically represented by matrices over some field Ω of characteristic zero, it is necessary and sufficient that the following conditions be satisfied:

- 1) in Γ the minimal condition holds for the centralizers of an ascending sequence of subgroups of Γ ;
- 2) in Γ there is a normal series

$$\Gamma \supset \Gamma_1 \supset \Gamma_2 \supset \{e\},$$

where Γ/Γ_1 is a finite group, Γ_1/Γ_2 is an abelian group, and the group Γ_2 can be isomorphically embedded in the ΩR -completion of a nilpotent group H of finite Ω -rank, with the restriction of each inner automorphism of the group Γ_1 to the subgroup Γ_2 inducing an Ω -automorphism of the group H .

Using the theorem of A. I. Mal'cev cited above, D. M. Smirnov showed in ⁽⁶⁾ that a finitely generated free solvable group of derived length 3 has no faithful matrix representation over any field. In connection with this there arises the question of the representability of two-step solvable groups by matrices over some field of characteristic zero. It is quite easy to construct an example of a two-step solvable torsion-free group which cannot be faithfully represented by matrices over any field of characteristic zero. In particular, such a group will be

the group Γ which is the semidirect product of the direct sum H of a countable number of rational groups H_n ($n = 1, 2, \dots$) and the infinite cyclic group $\{z\}$, where $z^{-1}h_n z = nh_n$ ($h_n \in H_n$).

Theorem 3. Every finitely generated two-step solvable torsion-free group Γ has a faithful matrix representation over some field of characteristic zero.

Theorem 4. The discrete wreath product $G = \Gamma \wr H$, where Γ and H are torsion-free abelian groups, is isomorphically embeddable in the group of matrices of the second order over some field of characteristic zero.

Theorem 5*. A free nilpotent group of nilpotency class n admits a faithful matrix representation over some field of characteristic zero.

Latvian State University
named after P. Stučka
Riga

Received
10 II 1969

CITED LITERATURE

- ¹ A. I. Mal'cev, *Matem. sborn.*, **28**, 567 (1951).
- ² L. Auslander, *Ann. Math.*, No. 7 (1967).
- ³ R. Swan, *Proc. Am. Math. Soc.*, **18**, 385 (1967).
- ⁴ F. Hall, *Collection of Translations. Mathematics*, **12**, 1, 3 (1968).
- ⁵ M. I. Kargapolov, *Algebra and Logic, Seminar*, **6**, 5, 17 (1967).
- ⁶ D. M. Smirnov, *DAN*, **155**, No. 3, 535 (1964).

* This theorem was obtained jointly with V. G. Vilyatser.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.