

ASYMPTOTICS OF CERTAIN SUBSEQUENCES OF EIGENVALUES OF BOUNDARY-VALUE PROBLEMS FOR THE HELMHOLTZ EQUATION IN THE MULTIDIMENSIONAL CASE

M. M. Popov

1969

SovietRxiv

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 534.21

MATHEMATICAL PHYSICS

M. M. POPOV

ASYMPTOTICS OF CERTAIN SUBSEQUENCES OF EIGENVALUES OF BOUNDARY-VALUE PROBLEMS FOR THE HELMHOLTZ EQUATION IN THE MULTIDIMENSIONAL CASE

(Presented by Academician V. I. Smirnov on 10 VI 1968)

1. Let Ω be a domain of Euclidean space R^{n+1} , bounded by a sufficiently smooth closed surface Σ . We consider problems for the Helmholtz equation

$$[\Delta + \omega^2 C^{-2}(M)]U(M) = 0, \quad M \in \Omega; \quad (1)$$

$$\text{I. } U|_{\Sigma} = 0 \quad \left(\text{or II. } \frac{\partial}{\partial n} U|_{\Sigma} = 0 \right), \quad (2)$$

where Δ is the Laplace operator in R^{n+1} ; $C(M) > 0$ is a sufficiently smooth function in Ω ; ω is a real parameter.

The present note is devoted to constructing, by the ray method, asymptotics of such subsequences of eigenvalues of problem (1), (2) to which there correspond eigenfunctions concentrated in a neighborhood of certain one-dimensional cycles in Ω . Asymptotically, as $\omega \rightarrow \infty$, within the framework of the ray method the solution of equation (1) is sought in the form $U(M) \sim [u + O(1/\omega)] \exp(i\omega\Phi)$, where u and Φ satisfy the equations

$$(\nabla\Phi)^2 = C^{-2}(M), \quad (3)$$

$$2(\nabla u, \nabla\Phi) + u\Delta\Phi = 0. \quad (4)$$

2. Consider in Ω a closed curvilinear N -gon L_N , whose sides are formed by smooth curves and whose vertices belong to Σ . We shall call the polygon L_N an extremal one-dimensional cycle in Ω if the first variation $\delta I = 0$ on L_N , where $I = \int C^{-1}(M) d\sigma$ is the functional of geometrical optics ($d\sigma$ is an element of length in R^{n+1}).

In what follows it is assumed that, if the extremal cycle under consideration in Ω is stable in the first approximation ^(1,2), then there exists the desired subsequence of eigenvalues of problem (1), (2), to which there correspond eigenfunctions concentrated in a neighborhood of this cycle, i.e. essentially different from zero in some neighborhood of the cycle and decreasing sufficiently rapidly when moving away from it.

3. Let $\mathbf{r} = \mathbf{r}(s)$, $\mathbf{r} \in R^{n+1}$, be an extremal cycle in Ω with vertices $\mathbf{r}(s_1), \dots, \mathbf{r}(s_N) = \mathbf{r}(0)$; s is arc length along the cycle. In each plane normal to the cycle, introduce the radius vector $\mathbf{q}(s) \in R^n$ so that along the cycle $\mathbf{q}(s) \equiv 0$. Denote the components of $\mathbf{q}(s)$ by q_j , $j = 1, \dots, n$. The conjugate momenta are defined by the formula

$$p_j = C^{-1}(s, q_j) \frac{\partial}{\partial \dot{q}_j} \frac{d\sigma}{ds},$$

where $\dot{q}_j = \frac{d}{ds} q_j$. We expand the Hamiltonian function $\mathcal{H}(s; q_j; p_j)$, corresponding to the functional I , in a neighborhood of the cycle into the series

$$\mathcal{H}(s; q_j; p_j) = -\frac{1}{C(s, 0)} + \mathcal{H}_2(s; q_j; p_j) + \mathcal{H}_3(s; q_j; p_j) + \dots, \quad (5)$$

where $\mathcal{H}_k(s; q_j; p_j)$ is a homogeneous polynomial of degree k in the canonical variables with coefficients depending on s . In the first approximation, discarding in the expansion (5) the terms of order $\mathcal{H}_k$, $k \geq 3$, we replace I by the functional I_0 :

$$I_0 = \int \left\{ \sum_{j=1}^n p_j dq_j + \left[\frac{1}{C(s, 0)} - \mathcal{H}_2(s; q_j; p_j) \right] ds \right\}, \quad (6)$$

for which the canonical system of equations has the form

$$\frac{d}{ds} q_j = \frac{\partial \mathcal{H}_2}{\partial p_j}, \quad \frac{d}{ds} p_j = -\frac{\partial \mathcal{H}_2}{\partial q_j}, \quad j = 1, 2, \dots, n. \quad (7)$$

A solution of the system (7)

$$\vec{\chi}^{(k)}(s) = (q_1^{(k)}(s), \dots, q_n^{(k)}(s), p_1^{(k)}(s), \dots, p_n^{(k)}(s)),$$

defined on the k -th ($k = 1, 2, \dots, N$) side L_N , i.e. for $s_{k-1} \leq s \leq s_k$, will be called a ray. Let the $2n$ vectors $Z_t^{(k)}(s)$, $t = 1, \dots, 2n$, form on the k -th side L_N a fundamental system of solutions of the equations (7). With the aid of the matrix

$$W^{(k)}(s) = \|Z_1^{(k)}(s), \dots, Z_{2n}^{(k)}(s)\|$$

the ray is defined by the formula

$$\vec{\chi}^{(k)}(s) = W^{(k)}(s) \mathbf{A}^{(k)}$$

through the specification of a $2n$ -dimensional vector $\mathbf{A}^{(k)}$ independent of s .

Upon reflection of the ray from the boundary Σ in a neighborhood of the k -th vertex L_N we have:

$$\mathbf{A}^{(k+1)} = [W^{(k+1)}(s_k)]^{-1} \cdot \gamma_k \cdot W^{(k)}(s_k) \cdot \mathbf{A}^{(k)} + O(\|\vec{\chi}\|^2), \quad (8)$$

where the reflection matrices γ_k of order $2n$ are determined by the local properties of Σ at the point $\mathbf{r}(s_k)$, $\det \gamma_k = 1$. Let $\mathbf{A}_0^{(k)}$ specify some initial ray, and let $\mathbf{A}_1^{(k)}$ be the ray arising after a single "traversal" of the initial ray along the cycle; then, in the first approximation, discarding the correction terms in formula (8), we obtain

$$\mathbf{A}_1^{(k)} = \Gamma_k \mathbf{A}_0^{(k)},$$

and after m traversals

$$\mathbf{A}_m^{(k)} = \Gamma_k^m \mathbf{A}_0^{(k)},$$

with $\det \Gamma_k = 1$ by virtue of the existence of the integral invariant

$$\int \delta q_1 \dots \delta q_n \delta p_1 \dots \delta p_n$$

for the canonical system (7).

An extremal cycle is called stable in the first approximation (1) if

$$\|\mathbf{A}_m^{(k)}\| < M < \infty$$

as $m \rightarrow \infty$. Let λ_t , $t = 1, \dots, 2n$, be the eigenvalues of the matrix Γ_k^* . The conditions for stability of the cycle in the first approximation are that $|\lambda_t| = 1$, and the elementary divisors of Γ_k are simple.

4. For cycles stable in the first approximation there exist $2n$ linearly independent Floquet solutions of the equations (7), $X_t(s)$, such that

$$X_j(s+s_N) = e^{i\varphi_j} X_j(s), \quad X_{n+j}(s+s_N) = e^{-i\varphi_j} X_{n+j}(s),^{**} \quad j = 1, \dots, n.$$

We divide the Floquet solutions into two groups of n each so that, for the solutions of the first group,

$$W(X_j, \bar{X}_t) \equiv \sum_{k=1}^n (q_k^{(j)}(s) \cdot \bar{p}_k^{(t)}(s) - p_k^{(j)}(s) \cdot \bar{q}_k^{(t)}(s)) = -i\delta_{jt}, \quad j, t = 1, \dots, n,$$

where δ_{jt} is the Kronecker symbol; $\bar{X}_t$ is the vector complex-conjugate to the vector X_t . At the same time, for the Floquet solutions of the second group

$$W(X_{n+j}, \bar{X}_{n+t}) = +i\delta_{jt}.$$

Introduce normal coordinates ⁽⁴⁾ Q_j, P_j by the formula

$$\vec{\chi} = \sum_{j=1}^n (\mathbf{B}_j(s)Q_j + \mathbf{B}_{n+j}(s)P_j).$$

The functions $\mathbf{B}_t(s)$, periodic on the cycle, have the form

$$\mathbf{B}_j(s) = e^{-i\mu_j s} X_j(s), \quad \mathbf{B}_{n+j}(s) = e^{i\mu_j s} X_{n+j}(s), \quad \mu_j = \varphi_j/s_N.$$

In nor-

* The characteristic equation of the matrix Γ_k is reciprocal (see, for example, (3)) and therefore reduces to an equation of degree n .

** By these formulas the phases φ_j of the eigenvalues of the matrix Γ_k are determined uniquely as the increment of $\arg X_j$ on the cycle.

normal coordinates

$$I_0 = \int \left\{ -i \sum_{j=1}^n P_j dQ_j + \left[\frac{1}{C(s, 0)} - \sum_{j=1}^n \mu_j P_j Q_j \right] ds \right\},$$

and the ray is described by the functions $Q_j(s) = Q_j(0)e^{i\mu_j s}$, $P_j(s) = P_j(0)e^{-i\mu_j s}$, continuous on the cycle. Here $Q_j(0) = \bar{P}_j(0)$ are constants of integration.

5. The set of rays whose constants of integration satisfy the conditions $Q_j(0)P_j(0) = \varepsilon_j^2$, $j = 1, \dots, n$, where ε_j are fixed real parameters, forms in the space $R^{2n} \times S^{1*}$ an invariant manifold T_ε , since for all s

$$Q_j(s)P_j(s) = Q_j(0)P_j(0) = \varepsilon_j^2, \quad j = 1, \dots, n. \quad (9)$$

The section of T_ε by the Euclidean plane R^{2n} , $s = \text{const}$, is the direct product of n circles (9); therefore topologically T_ε is described as the direct product of $(n+1)$ circles $S^1 \times S^1 \times \dots \times S^1$. The manifolds T_ε are Lagrangian; therefore the solution of the eikonal equation (3) is found by integrating over T_ε the total differential

$$d\Phi = -i \sum_{j=1}^n P_j dQ_j + \left[\frac{1}{C(s, 0)} - \sum_{j=1}^n \mu_j P_j Q_j \right] ds.$$

6. The asymptotics of the eigenvalues of the problem (1), (2) within the framework of the ray method is found from the “quantization conditions”^(5,6) following from the requirement that the function $ue^{i\omega\Phi}$ be single-valued on the invariant manifold T_ε . The one-dimensional homology group $H_1(T_\varepsilon)$ of the manifold T_ε is the direct product of $(n+1)$ free cyclic groups and has $(n+1)$ generators l_k , $k = 0, 1, \dots, n$. In the case of boundary-value problem I, the “quantization conditions,” which constitute a system of $(n+1)$ equations with respect to ω and the n parameters ε_j , have the form

$$\omega \oint_{l_k} d\Phi = 2\pi m_k + \pi m'_k + \frac{\pi}{2} m''_k, \quad k = 0, 1, \dots, n, \quad (10)$$

where m_k is an integer; m'_k and m''_k are, respectively, the “reflection” and “caustic” intersection indices, i.e. the Kronecker intersection indices (see also ^(6,10)) of the oriented basis cycles l_k , respectively, with the boundary Σ and the caustic. As n basis cycles l_j , $j = 1, \dots, n$, we take the circles (9) in the plane $s = \text{const}$; for them $m'_j = 0$, $m''_j = 2$. The cycle l_0 is fixed by the conditions $Q_j = \text{const}$, $0 \leq s \leq s_N$; for it $m'_0 = N$, $m''_0 = 0$.

From equations (10) there follows a formula for the eigenvalues $\omega_{[m]}$

$$\omega_{[m]} \int_0^{s_N} \frac{ds}{C(s, 0)} = 2\pi(m_0 + 1/2 N) + \sum_{j=1}^n (m_j + 1/2) \varphi_j, \quad (11)$$

where $m_c \gg 1$; φ_j are the Floquet exponents of the first group of Floquet solutions.

In the case of boundary-value problem II, in formula (11) the term $1/2 N$ should be omitted, since in this case the term $\pi m'_k$ is absent in equations (10).

Formula (11), obtained on the basis of the linear system (7), coincides with the formula found by the parabolic-equation method ^(7,8).

* The one-dimensional cycles under consideration are homeomorphic to the circle S^1 , since the points of self-intersection L_N , if there are any, should, in view of the linearity of the problem (1), (2), be regarded as lying on different copies of the domain Ω .

7. Using “Birkhoff series”⁹, one can, within the framework of the ray method, successively take into account further terms in the expansion (5), provided that φ_j and 2π are linearly independent over the ring of integers. In this case, at each step, in a neighborhood of a cycle stable in the first approximation, we shall have a continuous family of invariant Lagrangian manifolds with the same topology as in the first approximation. The correction terms to formula (11) that arise in this way are homogeneous polynomials in $(m_j + 1/2)$ of the second, etc., degrees. There are grounds,

however, for believing (see ¹¹, where, in a special case, the next terms in the formula for the eigenvalues were obtained) that the corrections obtained by the ray method are meaningful only for $m_0 \gg m_j \gg 1$.

The proposed scheme admits a generalization to the case of more general elliptic operators.

The author expresses his deep gratitude to his scientific adviser V. S. Buldyrev for his help in this work.

Moscow State University
named after M. V. Lomonosov

Received
2 VI 1968

REFERENCES

- ¹ V. S. Buldyrev, *Vestn. LGU*, No. 22, issue 4, 38 (1965).
- ² V. S. Buldyrev, M. M. Popov, *Optics and Spectroscopy*, 20, 905 (1966).
- ³ A. M. Lyapunov, *The General Problem of the Stability of Motion*, Ch. III, Moscow—Leningrad, 1950.
- ⁴ E. T. Whittaker, *Analytical Dynamics*, Moscow—Leningrad, 1937, § 192.
- ⁵ J. Keller, S. Rubinow, *Ann. Phys.*, 9, No. 1, 24 (1960).
- ⁶ V. P. Maslov, *Perturbation Theory and Asymptotic Methods*, Moscow, 1965.
- ⁷ V. M. Babich, *Proceedings of the LOMI Seminars* (in press).
- ⁸ M. M. Popov, *Vestn. LGU* (in press).
- ⁹ G. D. Birkhoff, *Dynamical Systems*, Moscow—Leningrad, 1941.
- ¹⁰ V. I. Arnold, *Functional Analysis and Its Applications*, 1, issue 1, 1 (1967).
- ¹¹ V. F. Lazutkin, *Optics and Spectroscopy*, 24, 453 (1968).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.