

COMPACT APPROXIMATION OF OPERATORS AND APPROXIMATE SOLUTION OF OPERATOR EQUATIONS

G. M. Vainikko

1969

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-196901.66888>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.948 : 513.88

MATHEMATICS

G. M. VAINIKKO

COMPACT APPROXIMATION OF OPERATORS AND APPROXIMATE SOLUTION OF OPERATOR EQUATIONS

(Presented by Academician V. I. Smirnov on 25 III 1969)

The concept of compact approximation of operators was introduced in ^(1,2). The situation considered below is more general in several respects.

1. Let E, F, E_n, F_n ($n = 1, 2, \dots$) be Banach spaces. Suppose there exist operators $p_n \in \mathcal{L}(E, E_n)$, $q_n \in \mathcal{L}(F, F_n)$ with the following properties:

$$p_n E = E_n, \quad q_n F = F_n \quad (n = 1, 2, \dots); \quad (1)$$

$$\|p_n\| \leq c, \quad \|q_n\| \leq c \quad (n = 1, 2, \dots; c = \text{const}); \quad (2)$$

$$\|\xi_n\|_{E_n} \geq \gamma \inf_{x \in E, p_n x = \xi_n} \|x\|_E, \quad \|\eta_n\|_{F_n} \geq \gamma \inf_{y \in F, q_n y = \eta_n} \|y\|_F \quad (3)$$

for any $\xi_n \in E_n$, $\eta_n \in F_n$ ($n = 1, 2, \dots; \gamma = \text{const} > 0$);

$$\lim_{n \rightarrow \infty} \|p_n x\| > 0, \quad \lim_{n \rightarrow \infty} \|q_n y\| > 0 \quad \text{for } \forall x \in E, y \in F (x \neq 0, y \neq 0). \quad (4)$$

Definition 1. A sequence of operators $T_n \in \mathcal{L}(E_n, F_n)$ **compactly approximates** an operator $T \in \mathcal{L}(E, F)$ with respect to $\{p_n\}$ and $\{q_n\}$, if the following conditions are satisfied:

- a) $\|q_n T x - T_n p_n x\| \rightarrow 0$ as $n \rightarrow \infty$ for every $x \in E$;
- b) for any sequence $\{\xi_n\}$ ($\xi_n \in E_n$, $\|\xi_n\| \leq 1$, $n = 1, 2, \dots$) there exist such $x_n \in E$, $y_n \in F$ and $c' = \text{const}$ that $p_n x_n = \xi_n$, $q_n y_n = T_n \xi_n$, $\|y_n\| \leq c' \|T_n \xi_n\|$ ($n = 1, 2, \dots$), and the sequence $\{y_n - T x_n\}$ is compact in F .

Remark 1. If $T \in \mathcal{L}(E, F)$ is completely continuous, then condition b) is equivalent to the following condition (cf. ⁽¹⁾): for any sequence $\{\xi_n\}$ ($\xi_n \in E_n$, $\|\xi_n\| \leq 1$, $n = 1, 2, \dots$) there exist such $y_n \in F$ that $q_n y_n = T_n \xi_n$ ($n = 1, 2, \dots$), and the sequence $\{y_n\}$ is compact in F .

Consider the equations

$$Tx = y \quad (y \in F), \quad (5)$$

$$T_n \xi_n = q_n y. \quad (6)$$

Theorem 1. Suppose the following conditions are satisfied:

- 1) the sequence $T_n \in \mathcal{L}(E_n, F_n)$ compactly approximates $T \in \mathcal{L}(E, F)$;
- 2) T has an inverse $T^{-1} \in \mathcal{L}(F, E)$;
- 3) the operators T_n are such that from

$$\inf_{\xi_n \in E_n, \|\xi_n\|=1} \|T_n \xi_n\| > 0$$

it follows that there exists an inverse $T_n^{-1} \in \mathcal{L}(F_n, E_n)$.

Then equation (6) has, for all sufficiently large n , a unique solution $\xi_n^* \in E_n$, and $\|\xi_n^* - p_n x^*\| \rightarrow 0$ as $n \rightarrow \infty$, where $x^* = T^{-1}y$ is the solu-

* By $\mathcal{L}(X, Y)$ is denoted the space of linear continuous operators from X to Y .

** The formula $\varphi_n \xi_n = p_n^{-1}(\xi_n)$ defines an isomorphism $\varphi_n \in \mathcal{L}(E_n, E/p_n^{-1}(0))$, for which $\|\varphi_n\| \leq 1/\gamma$, $\|\varphi_n^{-1}\| \leq c$ ($n = 1, 2, \dots$); here $p_n^{-1}(\xi_n)$ is the complete preimage of the element $\xi_n \in E_n$. Similarly, F_n is isomorphic to the quotient space $F/q_n^{-1}(0)$.

solution of equation (5). The estimate is valid

$$c_1 \|q_n T x^* - T_n p_n x^*\| \leq \|\xi_n^* - p_n x^*\| \leq c_2 \|q_n T x^* - T_n p_n x^*\| \quad (c_1, c_2 = \text{const} > 0). \quad (7)$$

Proof. From 1) it follows that

$$\|T_n\| \leq 1/c_1 = \text{const} \quad (n = 1, 2, \dots). \quad (8)$$

Let us show that, for sufficiently large n , there exist $T_n^{-1} \in \mathcal{L}(F_n, E_n)$ and

$$\|T_n^{-1}\| \leq c_2 = \text{const} \quad (n = n_0, n_0 + 1, \dots). \quad (9)$$

Suppose that for some $\xi_n \in E_n$ ($\|\xi_n\| = 1$, $n = 1, 2, \dots$) we have $\|T_n \xi_n\| \rightarrow 0$ as $n \rightarrow \infty$. Choose $x_n \in E$, $y_n \in F$ ($n = 1, 2, \dots$) such that $p_n x_n = \xi_n$, $q_n y_n = T_n \xi_n$, $\|y_n\| \leq c' \|T_n \xi_n\| \rightarrow 0$, and $\{y_n - T x_n\}$ is compact in F . Then $\{x_n\}$ is compact in E , and from condition a) of Definition 1 we conclude that $\|q_n T x_n - T_n p_n x_n\| \rightarrow 0$ as $n \rightarrow \infty$. Since $\|T_n p_n x_n\| = \|T_n \xi_n\| \rightarrow 0$, it follows that $\|q_n T x_n\| \rightarrow 0$, and also $\|q_n T x'\| \rightarrow 0$ as $n \rightarrow \infty$ for every limit point x' of the sequence $\{x_n\}$; by (4), $T x' = 0$ and $x' = 0$. Hence $x_n \rightarrow 0$ as $n \rightarrow \infty$, which contradicts (2) and the equalities $\|p_n x_n\| = \|\xi_n\| = 1$. Together with 3), this contradiction proves (9).

From inequalities (8), (9) and the equality

$$T_n(\xi_n^* - p_n x^*) = q_n T x^* - T_n p_n x^*$$

the estimate (7) follows. The convergence $\|\xi_n^* - p_n x^*\| \rightarrow 0$ follows from (7) and condition a) of Definition 1. Theorem 1 is proved.

Inequalities (9), together with condition a) of Definition 1, mean, in the terminology of $(^3, ^4)$, that the sequence $T_n \in \mathcal{L}(E_n, F_n)$ stably approximates $T \in \mathcal{L}(E, F)$. In $(^3, ^4)$ it is proved that stability of the approximation is not only a sufficient but also a necessary condition for the convergence $\|\xi_n^* - p_n x^*\| \rightarrow 0$ for any right-hand side $y \in F$ of equation (5). Theorem 1 can be reformulated as follows: if the sequence $T_n \in \mathcal{L}(E_n, F_n)$ compactly approximates $T \in \mathcal{L}(E, F)$ and if conditions 2) and 3) of Theorem 1 are fulfilled, then the approximation is stable. The converse assertion is true only in the following weakened form.

Remark 2. Let the sequence $\tilde{T}_n \in \mathcal{L}(E_n, F_n)$ stably approximate $T \in \mathcal{L}(E, F)$. Let $TE = F$, and let T admit a compact approximation by some sequence of operators $T_n \in \mathcal{L}(E_n, F_n)$, for which condition 3) of Theorem 1 is fulfilled. Then there exist $\tilde{q}_n \in \mathcal{L}(F, F_n)$ with properties (1)–(4) such that, with respect to $\{p_n\}$ and $\{\tilde{q}_n\}$, the sequence $\tilde{T}_n$ compactly approximates T .

2. Consider the case where the operators $T : E \rightarrow F$ and $T_n : E_n \rightarrow F_n$ in equations (5) and (6) are nonlinear.

Lemma 1. Let the operator T_n be Fréchet differentiable in the ball $\{\xi_n \in E_n : \|\xi_n - \xi_n^0\| \leq \delta_0\}$, where $\delta_0 > 0$, $\xi_n^0 \in E_n$. Let $T'_n(\xi_n^0) \in \mathcal{L}(E_n, F_n)$ have an inverse $[T'_n(\xi_n^0)]^{-1} \in \mathcal{L}(F_n, E_n)$, and suppose that for some δ and θ ($0 < \delta \leq \delta_0$, $0 \leq \theta < 1$) the inequalities

$$\sup_{\xi_n \in E_n, \|\xi_n - \xi_n^0\| \leq \delta} \|[T'_n(\xi_n^0)]^{-1} [T'_n(\xi_n) - T'_n(\xi_n^0)]\| \leq \theta, \quad (10)$$

$$\alpha_n \equiv \|[T'_n(\xi_n^0)]^{-1} [T_n \xi_n^0 - q_n y]\| \leq \delta(1 - \theta). \quad (11)$$

Then equation (6) has, in the ball $\|\xi_n - \xi_n^0\| \leq \delta$, a unique solution ξ_n^* , and the estimate is valid

$$\alpha_n/(1 + \theta) \leq \|\xi_n^* - \xi_n^0\| \leq \alpha_n/(1 - \theta). \quad (12)$$

The proof is analogous to the proof of Theorem 2 from (5).

Theorem 2. Suppose that the following conditions are fulfilled:

- 1) equation (5) has a solution x^* ;
- 2) the operator T is Fréchet differentiable at the point x^* , and $T'(x^*) \in \mathcal{L}(E, F)$ has an inverse $[T'(x^*)]^{-1} \in \mathcal{L}(F, E)$;
- 3) the operators T_n ($n = 1, 2, \dots$) are Fréchet differentiable in the corresponding balls $\|\xi_n - p_n x^*\| \leq \delta_0$ ($\delta_0 = \text{const} > 0$), and for any $\varepsilon > 0$ there exist n_ε and η_ε ($0 < \eta_\varepsilon \leq \delta_0$) such that $\|T'_n(\xi_n) - T'_n(p_n x^*)\| \leq \varepsilon$ for $n \geq n_\varepsilon$, $\|\xi_n - p_n x^*\| \leq \eta_\varepsilon$;
- 4) $\|q_n T x^* - T_n p_n x^*\| \rightarrow 0$ as $n \rightarrow \infty$ ($q_n T x^* = q_{ny}$);
- 5) the sequence $T'_n(p_n x^*) \in \mathcal{L}(E_n, F_n)$ compactly approximates the operator $T'(x^*) \in \mathcal{L}(E, F)$;
- 6) the operators $T'_n(p_n x^*)$ are such that from

$$\inf_{\xi_n \in E_n, \|\xi_n\|=1} \|T'_n(p_n x^*)\xi_n\| > 0$$

there follows the existence of an inverse $[T'_n(p_n x^*)]^{-1} \in \mathcal{L}(F_n, E_n)$.

Then there exist N and $\delta > 0$ such that, for $n \geq N$, equation (6) has in the ball $\|\xi_n - p_n x^*\| \leq \delta$ a unique solution ξ_n^* . As $n \rightarrow \infty$, the convergence $\|\xi_n^* - p_n x^*\| \rightarrow 0$ holds with the estimate (7).

Proof is based on Lemma 1, in which we put $\xi_n^0 = p_n x^*$. From 2), 5), and 6) we conclude that, for sufficiently large n , the inverses $[T'_n(p_n x^*)]^{-1} \in \mathcal{L}(F_n, E_n)$ exist, and their norms are bounded in the aggregate. Fix θ ($0 < \theta < 1$). Using condition 3), we find $\delta > 0$ such that, for sufficiently large n , (10) will be satisfied. By virtue of 4), for sufficiently large n (11) is also fulfilled. From (12) we obtain the estimate (7); from (7) and 4) follows the convergence $\|\xi_n^* - p_n x^*\| \rightarrow 0$ as $n \rightarrow \infty$. Theorem 2 is proved.

3. Let now $E = F$, $E_n = F_n$, $p_n = q_n$ ($n = 1, 2, \dots$). In studying the closeness of solutions of the operator equations

$$x = Tx, \quad (13)$$

$$\xi_n = T_n \xi_n \quad (14)$$

with completely continuous operators T and T_n , it is natural to use the concept of the rotation of vector fields ⁽⁶⁾. We shall assume that the Banach spaces E and E_n ($n = 1, 2, \dots$) are real. Let Ω be a bounded domain (a connected open set) in E ; denote $\Omega_n = p_n \Omega$. Then Ω_n is a bounded domain in E_n . By $\bar{\Omega}$, $\dot{\Omega}$, $\bar{\Omega}_n$, $\dot{\Omega}_n$ we denote the closures and boundaries of the domains Ω and Ω_n . It is clear that $p_n \bar{\Omega} \subset \bar{\Omega}_n$.

Definition 2. A sequence of operators $T_n : \bar{\Omega}_n \rightarrow E_n$ **compactly approximates** the completely continuous operator $T : \bar{\Omega} \rightarrow E$ with respect to $\{p_n\}$, if the following conditions are fulfilled:

- a) $\|p_n T x - T_n p_{nx}\| \rightarrow 0$ as $n \rightarrow \infty$ for every $x \in \bar{\Omega}$;
- b) for any sequence $\{\xi_n\}$ ($\xi_n \in \bar{\Omega}_n$, $n = 1, 2, \dots$) there exist such $y_n \in E$ that $p_n y_n = T_n \xi_n$ ($n = 1, 2, \dots$) and the sequence $\{y_n\}$ is compact in E .

Below we assume that (4) is fulfilled in the following strengthened form:

$$\lim_{n \rightarrow \infty} \|p_{nx}\| \geq c_0 \|x\| \quad \text{for all } x \in E \quad (c_0 = \text{const} > 0). \quad (4')$$

Lemma 2. Suppose the following conditions are fulfilled:

- 1) the sequence of completely continuous operators $T_n : \bar{\Omega}_n \rightarrow E_n$ compactly approximates the completely continuous operator $T : \bar{\Omega} \rightarrow E$;
- 2) the operators T_n ($n = 1, 2, \dots$) are such that from $\|\xi_n - p_{nx}\| \rightarrow 0$ ($\xi_n \in \bar{\Omega}_n$, $x \in \bar{\Omega}$) it follows that $\|T_n \xi_n - T_n p_{nx}\| \rightarrow 0$ as $n \rightarrow \infty$;
- 3)

$$\lim_{n \rightarrow \infty} \inf_{\xi_n \in \dot{\Omega}_n} \|p_{nx} - \xi_n\| > 0 \quad \text{for every } x \notin \dot{\Omega} \quad (x \in E);$$

- 4) the operator T has no fixed points on the boundary $\dot{\Omega}$.

Then, for all sufficiently large n , the operator T_n has no fixed points on the boundary $\dot{\Omega}_n$, and the equality of rotations holds:

$$\gamma(\xi_n - T_n \xi_n; \dot{\Omega}_n) = \gamma(x - Tx; \dot{\Omega}) \quad (n = n_0, n_0 + 1, \dots).$$

Remark 3. Condition 3) of Lemma 2 is fulfilled if (2) and (4') hold with $c = c_0 = 1$, and Ω is a ball in E .

Theorem 3. Suppose conditions 1)–3) of Lemma 2 are satisfied, and suppose equation (13) has an isolated solution $x^* \in \Omega$ of nonzero index ⁽⁶⁾, unique in $\bar{\Omega}$. Then equation (14), for sufficiently large n , has at least one solution $\xi_n^* \in \bar{\Omega}_n$, and $\|\xi_n^* - p_n x^*\| \rightarrow 0$ as $n \rightarrow \infty$ for all solutions $\xi_n^* \in \bar{\Omega}_n$.

4. As an application, we consider the method of mechanical quadratures for solving integral equations. Let D be a metric compact set, ρ a metric in D , and ν a measure defined on some algebra of subsets of D . Assume that $|\nu|(D) < \infty$ (where $|\nu|$ is the total variation of ν) and that the ball $S(t_0, r) = \{t \in D : \rho(t, t_0) < r\}$ is ν -measurable for any $t_0 \in D, r > 0$, with $|\nu|(S(t_0, r)) > 0$. Clearly, every function continuous on D is ν -integrable. Consider the integral equation

$$x(t) = \int_D K(t, s, x(s)) \nu(ds), \quad (15)$$

a convergent quadrature process

$$\int_D z(s) \nu(ds) = \sum_{j=1}^n \alpha_{jn} z(s_{jn}) + \Phi_n(z) \quad (n = 1, 2, \dots)$$

$$(\Phi_n(z) \rightarrow 0 \text{ for continuous functions } z(s))$$

and the system of equations

$$\xi_{in} = \sum_{j=1}^n \alpha_{jn} K(s_{in}, s_{jn}, \xi_{jn}) \quad (i = 1, \dots, n). \quad (16)$$

Theorem 4. Suppose equation (15) has a solution $x^*(t)$ (continuous on D), and suppose the kernel $K(t, s, x)$ is continuous and has a continuous derivative $\partial K(t, s, x)/\partial x$ for $t, s \in D, |x - x^*(s)| \leq \delta_0$ ($\delta_0 = \text{const} > 0$). Suppose the equation

$$y(t) = \int_D H(t, s) y(s) \nu(ds) \quad (H(t, s) = \partial K(t, s, x^*(s))/\partial x)$$

has only the zero solution. Then there exist N and $\delta > 0$ such that, for $n \geq N$, the system of equations (16) has a unique solution $(\xi_{1n}^*, \dots, \xi_{nn}^*)$ such that $|\xi_{jn}^* - x^*(s_{jn})| \leq \delta$ ($j = 1, \dots, n$). The convergence

$$\max_{1 \leq j \leq n} |\xi_{jn}^* - x^*(s_{jn})| \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (17)$$

holds, with the estimate

$$c_1 \varepsilon_n \leq \max_{1 \leq j \leq n} |\xi_{jn}^* - x^*(s_{jn})| \leq c_2 \varepsilon_n \quad (c_1, c_2 = \text{const} > 0),$$

where

$$\varepsilon_n = \max_{1 \leq i \leq n} |\Phi_n(z_{in})|, \quad z_{in}(s) = K(s_{in}, s, x^*(s)).$$

The proof is based on Theorem 2, which is applied with $E = F = C(D)$, $E_n = F_n = m_n$, $p_{nx} = q_{nx} = (x(s_{1n}), \dots, x(s_{nn}))$.

Theorem 3 makes it possible to establish the convergence (17), assuming only the continuity of the kernel $K(t, s, x)$ and the existence of a solution $x^*(t)$ of nonzero index (6) of equation (15).

Similar results in the case of Lebesgue measure on $D = [a, b] \subset R^1$ were obtained by another method in ⁵, and for linear equations in a case similar to the one considered, in ¹.

Tartu State University

Received

26 II 1969

References

- ¹ G. M. Vainikko, *Uch. zap. Tartu State Univ.*, 220 (1968).
- ² G. M. Vainikko, *Zh. Vychisl. Mat. i Mat. Fiz.*, 9, No. 4, 739 (1969).
- ³ N. N. Gudovich, *Zh. Vychisl. Mat. i Mat. Fiz.*, 6, No. 5 (1966).
- ⁴ S. G. Krein, *Linear Differential Equations in Banach Space*, "Nauka," 1967.
- ⁵ G. M. Vainikko, *Zh. Vychisl. Mat. i Mat. Fiz.*, 7, No. 4 (1967).
- ⁶ M. A. Krasnosel'skii, *Topological Methods in the Theory of Nonlinear Integral Equations*, 1956.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.