

EMBEDDING THEOREMS FOR FUNCTIONS DEFINED IN UNBOUNDED DOMAINS, AND THEIR APPLICATION TO THE SPECTRAL THEORY OF ELLIPTIC SELF-ADJOINT OPERATORS

J. V. Rybalov

1969

SovietRxiv

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.514

MATHEMATICS

EMBEDDING THEOREMS FOR FUNCTIONS DEFINED IN UNBOUNDED DOMAINS, AND THEIR APPLICATION TO THE SPECTRAL THEORY OF ELLIPTIC SELF-ADJOINT OP- ERATORS

Yu. V. Rybalov

(Presented by Academician I. M. Vinogradov on 26 VI 1968)

As K. Clark showed ⁽¹⁾, the embedding $W_2^1(\Omega)$ into $L_2(\Omega)$ is not completely continuous if the unbounded domain Ω has infinite measure. A consequence of this is the non-discreteness of the spectrum of the Laplace operator in a domain containing a countable number of identical nonintersecting squares (see ⁽²⁾).

In this paper, for a sufficiently general unbounded domain Ω , we consider the class $W_{p,\alpha}^l(\Omega, g)$, which coincides with the class $W_p^l(\Omega)$ in the case of a bounded domain (for $\Omega = E_n$, $W_{p,\alpha}^l(\Omega, g)$ is the class of L. D. Kudryavtsev $W_{p,\alpha}^l(E)$ ⁽³⁻⁵⁾), and it is shown that the embedding $W_{p,\alpha}^l(\Omega, g)$ into $W_{p,\alpha_k}^k(\Omega, g)$ is completely continuous ($0 \leq k < l$) for a certain exactly determined exponent of the weight degree α_k . With the help of this result an operator $a(X, D)$ is constructed, differing from the elliptic self-adjoint operator $L(X, D)$ by a positive factor of power growth at infinity, and it is proved that the operator $a(X, D)$ has a discrete spectrum in an unbounded domain not containing a sphere of positive radius.

Let the unbounded domain $\Omega \subset E_n$, $n \geq 1$, let g be a bounded interior subdomain of Ω , let α be a real number, and $1 \leq p \leq \infty$.

We introduce classes of functions as linear normed spaces of functions for which the following norms are meaningful and finite:

$$|f; L_{p,\alpha}(\Omega)| = |(1 + |X|)^{-\alpha} f; L_p(\Omega)|,$$

$$|f; L_{p,c(X)}(\Omega)| = |c^{1/p}(X)f; L_p(\Omega)|,$$

$$|f; W_{p,\alpha}^l(\Omega, g)| = |f; L_p(g)| + \sum_{|r|=l} |D^r f; L_{p,\alpha}(\Omega)| =$$

$$= |f; L_p(g)| + |f; L_{p,\alpha}^l(\Omega)| \quad (|f; W_{p,\alpha}^0(\Omega, g)| = |f; L_{p,\alpha}(\Omega)|),$$

$$|f; W_{p,\alpha,\alpha_0}^l(\Omega)| = |f; L_{p,\alpha_0}(\Omega)| + |f; L_{p,\alpha}^l(\Omega)|,$$

where $D^r f$ is a generalized derivative of order

$$|r| = \sum_{i=1}^n r_i.$$

Denote by $\overset{\circ}{B}$ the closure in the norm of the class B of functions from $C_0^\infty(\Omega)$. By definition, a domain Ω with locally Lipschitz boundary belongs to the class Z_m if

$$\Omega = G + \sum_{j=1}^N \Omega_j,$$

where G is a finite domain, $N < \infty$, and the domain Ω_j ($j = 1, \dots, N$) has the following properties after a nonsingular linear transformation, its own for each j :

- 1) for $X \in \Omega_j$, the coordinates x_i have $|x_i| \leq c < \infty$ ($i = m+1, \dots, n$), while the remaining coordinates $x_1, \dots, x_m$ are unbounded;
- 2) if one passes from the orthogonal coordinates X to cylindrical coordinates

$$Y: \quad y_1 = |X_m| = \left(\sum_{i=1}^m x_i^2 \right)^{1/2}, \quad y_2 = \varphi, \quad y_3 = \theta_1, \dots, y_m = \theta_{m-2}, \quad y_{m+1} = x_{m+1}, \dots$$

$$\dots, y_n = x_n \quad (y_1 = x_1 \text{ for } m = 1),$$

then for some $i > 1$, depending on j , the domain Ω_j is written in the form

$$\Omega_j = \{Y: a_k^{(j)} < y_k < b_k^{(j)} (k \neq i), \quad a_i^{(j)} < y_i < \psi_i^{(j)}(P_i) (\psi_i^{(j)}(P_i) \geq b_i^{(j)})\},$$

where $a_k^{(j)}, b_k^{(j)}$ are constants, $|a_k^{(j)}| < \infty$ ($k = 1, \dots, n$), $b_1^{(j)} = \infty$, $|b_k^{(j)}| < \infty$ ($k = 2, \dots, n$), $P_i = (y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n)$, and $\psi_i^{(j)}(P_i)$ is a single-valued continuous bounded function of P_i .

Theorem 1. Let $f \in W_{p,\alpha}^l(\Omega, g)$, $\Omega \in Z_m$, $1 \leq m \leq n$; let α be a real number; $1 \leq p \leq \infty$; l, k natural numbers, $0 \leq k < l$; $\beta = \max(\alpha, m/p - 1 + \varepsilon)$, $\varepsilon > 0$; $\alpha_k = \beta + l - k$; $|r| = k$.

Then

$$|D^r f; L_{p,\alpha_k}(\Omega)| \ll |f; W_{p,\alpha}^l(\Omega, g)|^*.$$

Corollary 1. Let $\Omega \in Z_m$, $1 \leq m \leq n$; let α be a real number; $1 \leq p \leq \infty$; l a natural number;

$$\beta = \max(\alpha, m/p - 1 + \varepsilon), \quad \varepsilon > 0; \quad \alpha_0 = \beta + l.$$

Then the classes $W_{p,\alpha}^l(\Omega, g)$, $W_{p,\alpha,\alpha_0}^l(\Omega)$ coincide up to equivalence of norms.

Theorem 1 for $\Omega = E_n$ is contained in the works ⁽³⁻⁵⁾.

Theorem 2. Let $\Omega \in Z_m$, $1 \leq m \leq n$; let α be a real number; $1 \leq p \leq \infty$; l, k natural numbers, $0 \leq k < l$; $\beta = \max(\alpha, m/p - 1 + \varepsilon)$, $\varepsilon > 0$; $\alpha_k = \beta + l - k$.

Then $W_{p,\alpha}^l(\Omega, g)$ is embedded in $W_{p,\alpha_k+\varepsilon}^k(\Omega, g)$ completely continuously.

Theorem 3. Let $\alpha \geq 0$; $1 \leq p \leq \infty$; l, k be natural numbers, $0 \leq k < l$; $\varepsilon > 0$; $\alpha_k = \alpha + l - k$.

Then the embedding $\dot{W}_{p,\alpha}^l(\Omega, g)$, $\dot{W}_{p,\alpha_k+\varepsilon}^k(\Omega, g)$ is completely continuous.

Corollary. Let the unbounded domain Ω contain no sphere of positive radius; $\alpha \geq 0$; $1 \leq p < \infty$; l, k be natural numbers, $0 \leq k < l$; $\varepsilon > 0$; $\alpha_k = \alpha + l - k$.

Then the embedding $\dot{L}_{p,\alpha}^l(\Omega)$ in $\dot{L}_{p,\alpha_k+\varepsilon}^k(\Omega)$ is completely continuous.

The stated theorems are sharp in the sense that the exponents of the weight cannot be decreased by $\varepsilon > 0$.

Theorem 4. Let the unbounded domain Ω contain no sphere of positive radius. Suppose, further, that $\alpha \geq 0$; $\varepsilon > 0$; $\alpha_0 = \alpha + l + \varepsilon$; $\beta = \alpha$ for $\alpha > n/2 - 1$, $\beta = \alpha + \varepsilon$ for $\alpha \leq n/2 - 1$; $L(X, D)$ is a linear elliptic self-adjoint operator of order $2l$ (D is the differentiation operator in the sense of distributions)

$$\begin{aligned} L(X, D) &= (-1)^l \sum_{\sum \alpha_i = l} \frac{l!}{\alpha_1! \dots \alpha_n!} \frac{\partial^l}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}} \left(a_{\alpha_1 \dots \alpha_n}(X) \frac{\partial^l}{\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}} \right) + b(X), \\ &\sum_{\sum \alpha_i = l} \frac{l!}{\alpha_1! \dots \alpha_n!} a_{\alpha_1 \dots \alpha_n}(X) (\xi_1^{\alpha_1} \dots \xi_n^{\alpha_n})^2 \geq \\ &\geq \frac{B}{(1 + |X|^2)^\alpha} \sum_{\sum \alpha_i = l} \frac{l!}{\alpha_1! \dots \alpha_n!} (\xi_1^{\alpha_1} \dots \xi_n^{\alpha_n})^2 \quad \text{for } X \in \Omega, \quad \vec{\xi} \in E_n, \\ a(X, D) &= c^{-1}(X) L(X, D); \end{aligned}$$

$a_{\alpha_1 \dots \alpha_n}(X)$, $b(X)$, $c(X)$ are infinitely differentiable coefficients in Ω ;

$$|a_{\alpha_1 \dots \alpha_n}(X)| \leq A/(1 + |X|^2)^\alpha; \quad 0 \leq$$

* By the expression $A \ll B$ here and below is meant $A \leq cB$, where the positive constant c does not depend

$$\leq b(X) \leq A/(1 + |X|^2)^{\beta+l}; \quad 0 < c(X) \leq A/(1 + |X|^2)^{\alpha_0};$$

A , B are positive constants.

Define the operator T in $L_{2,c(X)}(\Omega)$ as follows:

$$D(T) = \mathring{L}_{2,\alpha}^l(\Omega) \cap \{f \in L_{2,c(X)}(\Omega) : a(X, D)f \in L_{2,c(X)}(\Omega)\},$$

$$Tf = a(X, D)f, \quad f \in D(T).$$

Then T is a closed linear operator; the spectrum $\sigma(T)$ is discrete, has no finite limit points, and is situated in the half-plane $\operatorname{Re} \lambda > 0$; for $\lambda \notin \sigma(T)$ the resolvent

$$R_\lambda(T) = (\lambda I - T)^{-1}$$

is completely continuous in $L_{2,c(X)}(\Omega)$.

We note that the operator $L(X, D)$ is equivalent to the operator $a(X, D)$ in the sense that the equations $L(X, D)u = f$ for $f \in L_{2,c^{-1}(X)}(\Omega)$ and

$$a(X, D)u = c^{-1}(X)f = g$$

for $g \in L_{2,c(X)}(\Omega)$ are equivalent (a generalized solution of one is a generalized solution of the other).

If, in the definition of the spectrum of the operator A , one replaces the resolvent

$$R_\lambda(A) = (\lambda I - A)^{-1}$$

by the resolvent

$$R_{\lambda,c}(A) = (\lambda c(X)I - A)^{-1},$$

then we obtain the definition of the c -spectrum of the operator A .

Corollary. Let the hypotheses of Theorem 4 be satisfied. Define the operator T_1 as follows:

$$D(T_1) = \mathring{L}_{2,\alpha}^l \cap \{f \in L_{2,c(X)}(\Omega) : L(X, D)f \in L_{2,c^{-1}(X)}(\Omega)\},$$

$$T_1 f = L(X, D)f, \quad f \in D(T).$$

Then T_1 is a closed linear operator, the c -spectrum $\sigma_c(T_1)$ is discrete, has no finite limit points, and is situated in the half-plane $\operatorname{Re} \lambda > 0$; for $\lambda \notin \sigma_c(T_1)$ the resolvent

$$R_{\lambda,c}(T_1) = (\lambda c(X)I - T_1)^{-1}$$

is completely continuous from $L_{2,c^{-1}(X)}(\Omega)$ into $L_{2,c(X)}(\Omega)$.

For $n = 1$ and $l = 1$, the c -spectrum of the operator $L(X, D)$, under other restrictions on the coefficients, was first considered by K. Friedrichs ⁽⁶⁾.

Theorem 5. Let the hypotheses of Theorem 4 be satisfied, with the sole exception that

$$\alpha_0 = \beta + l.$$

Then the equation

$$L(X, D)u = f$$

for a function f such that

$$(1 + |X|)^{\alpha_0} f \in L_2(\Omega),$$

has, moreover, a unique generalized solution

$$u \in \mathring{L}_{2,\alpha}^l(\Omega),$$

and the inequality

$$|u; \mathring{L}_{2,\alpha}^l(\Omega)| \leq |(1 + |X|)^{\alpha_0} f; L_2(\Omega)|$$

holds.

The operator $L(X, D)$ for $l = 1$ was considered earlier by L. D. Kudryavtsev ⁽⁷⁻⁹⁾ in connection with the variational method for solving elliptic equations posed in a half-space.

In conclusion, the author expresses his gratitude to his scientific supervisor Prof. L. D. Kudryavtsev for posing the problem and for his attention to the work.

Moscow

Institute of Physics and Technology

Received

20 VI 1968

REFERENCES

1. C. Clark, *Pacific J. Math.*, **19**, No. 2, 243 (1966).
2. A. M. Molchanov, *Tr. Mosk. matem. obshch.*, **2**, 169 (1953).

3. L. D. Kudryavtsev, *DAN*, **153**, No. 3, 34 (1963).
4. L. D. Kudryavtsev, *Matem. sborn.*, **69** (111), No. 4 (1966).
5. L. D. Kudryavtsev, *Matem. sborn.*, **70** (112), No. 1, 1 (1966).
6. K. Friedrichs, *Stud. and Essays present. to R. Courant*, N. Y., 1948, p. 145.
7. L. D. Kudryavtsev, *DAN*, **157**, No. 1, 45 (1964).
8. L. D. Kudryavtsev, Soviet-American Symposium on Partial Differential Equations, Novosibirsk, 1963.
9. L. D. Kudryavtsev, *Izv. AN SSSR, ser. matem.*, **31**, 1179 (1967).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.