

BIFURCATION POINTS OF THE HAMMERSTEIN EQUATION

P. P. Zabreiko, A. I. Povolotskii

1970

SovietRxiv

View the original and related papers at <https://sovietsrxiv.org/items/ru-197001.14186>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.948.33

MATHEMATICS

P. P. ZABREIKO, A. I. POVOLOTSKII

BIFURCATION POINTS OF THE HAMMERSTEIN EQUATION

(Presented by Academician I. N. Vekua on March 10, 1970)

The paper considers the problem of bifurcation points of the equation

$$\int_{\Omega} k(t, s) f[s, x(s)] ds = \lambda x(t). \quad (1)$$

Here Ω is a bounded closed subset of a finite-dimensional space. In what follows it is assumed that the function $f(s, u)$, defined on $\Omega \times R^n$ with values in R^n , satisfies the Carathéodory conditions and is potential ($f(s, u) = \text{grad } G(s, u)$; $G(s, 0) = 0$); the matrix $k(t, s)$ ($t, s \in \Omega$) is symmetric, measurable in the aggregate of the variables, and the linear operator defined by it

$$Kx(t) = \int_{\Omega} k(t, s)x(s) ds \quad (2)$$

in the Hilbert space of vector functions $H = \mathcal{L}_2$ is self-adjoint and has no more than a finite number of negative eigenvalues (each of finite multiplicity). The paper is a natural continuation of work ⁽¹⁾ on eigenvectors of the Hammerstein operator.

Equation (1) has the form $Ax = \lambda x$. Suppose that the Hammerstein operator A acts in some real Banach space E . If the condition $f(s, 0) = 0$ is fulfilled, then equation (1) has the zero solution for all real λ . A number λ_0 is called a **bifurcation point** of equation (1) if for any $\varepsilon, \delta > 0$ there exists a λ , $|\lambda - \lambda_0| < \varepsilon$, to which there corresponds a nonzero solution $x(t)$ of equation (1), with $\|x\|_E < \delta$. The paper also considers asymptotic bifurcation points (in this case fulfillment of the condition $f(s, 0) = 0$ is not assumed). A number λ_0 is called an **asymptotic bifurcation point** of equation (1) if for any $\varepsilon, \rho > 0$ there exists a λ , $|\lambda - \lambda_0| < \varepsilon$, to which there corresponds a solution $x(t)$ of equation (1), with $\|x\|_E > \rho$.

The problem of bifurcation points of equation (1), in the case where the role of E is played by various $\mathcal{L}_p$, was considered in a number of works by M. A.

Krasnosel' skii ^(2,3) and others, M. M. Vainberg ⁽⁴⁾ and others. A detailed analysis of the properties of operators in general Banach spaces of measurable functions (the so-called ideal spaces ^(5,6)) made it possible to obtain essentially new results for this problem; these results are presented in the present note.

1. Let E be an ideal space of measurable vector functions finite almost everywhere on Ω (see ^(1,5)). By E' we denote the ideal space dual to E ; by E^0 , the set of vector functions from E with absolutely continuous norm; by E'/E , the space of multipliers from E into E' . The spaces E^0 and E'/E are themselves ideal. Examples of ideal spaces are the space M_{u_0} (u_0 a nonnegative measurable function) of vector functions for which the norm $\|x\|_{M_{u_0}} = \inf\{\lambda : |x| \leq \lambda u_0\}$ is meaningful, the spaces $\mathcal{L}_p$, and Orlicz spaces.

A set in E is called w -bounded if for every $\varepsilon > 0$ this set has an ε -net T such that $|x| \leq u_0$ for all $x \in T$ and for some $u_0 \in E$. An operator acting from one ideal space E_1 into another E_2 is called w -bounded if it maps every norm-bounded set into a w -bounded set.

Let H_1 be the linear span of the eigenvectors of the operator (2) in H corresponding to negative eigenvalues. Denote $J = -P_1 + P_2$, where P_1 and P_2 are the projection operators, respectively, onto H_1 and $H_2 = H \ominus H_1$.

We shall state the conditions that will be assumed to hold in what follows. Let the superposition operator $fx(s) = f[s, x(s)]$ act from E into E' , and let the linear operator K (and consequently also $\tilde{K} = JK$) act from E' into E , with $E \subseteq H$. In this case in (1) the operator $C = \tilde{K}^{1/2}$ acts from H into E , while the adjoint operator C' acts from E' into H . The operator $G[s, x(s)]$ acts from E into $\mathcal{L}_1$, and therefore on H the Golomb functional is defined by

$$\Phi(y) = \int_{\Omega} G[s, Cy(s)] ds. \quad (3)$$

This functional turns out to be differentiable on H , and its gradient is the Krasnosel' skii operator $\Gamma = C'fC$.

If the operator K is positive definite in H , then $\tilde{K} = K$ and $\tilde{K}^{1/2} = K^{1/2}$. In this case the problem of bifurcation points of equation (1) reduces to the problem of bifurcation points of the equation $\Gamma y = \lambda y$ with the Krasnosel' skii operator Γ , acting in H . If, however, K has a finite number of negative eigenvalues, then the problem reduces to the problem of bifurcation points of the equation $J\Gamma y = \lambda y$. This is also true for asymptotic bifurcation points.

2. Suppose that

$$f(s, u) = a_0(s)u + h(s, u)u, \quad (4)$$

where $a_0(s) \in E'/E$, and $h(s, u)$ ($h(s, 0) = 0$) is a function satisfying the Carathéodory conditions and defining the superposition operator $hx = h[s, x(s)]$,

acting from E into E'/E . By K_0 denote the linear integral operator with kernel $k(t, s)a_0(s)$.

Assume that one of the following conditions is fulfilled:

- a) the operator K is completely continuous;
- b) the operator f is w -bounded (or the operator G is w -bounded, and the operator f is uniformly continuous), $a_0(s) \in (E'/E)^0$, and the operator h is continuous at the zero point.
- c) $E = M_{u_0}$.

If any of these conditions is fulfilled, the Krasnosel' skii operator $\Gamma = C'fC$ has as its potential the weakly continuous and smooth (uniformly differentiable on each pair) Golomb functional Φ . Moreover, the operator Γ has at the zero point the completely continuous Fréchet derivative $B_0 = C'A_0C$, where A_0 is the operator of multiplication by the function $a_0(s)$. Therefore, using the theorem of M. A. Krasnosel' skii ⁽²⁾ and the theorem from ⁽³⁾, we obtain the following statements.

Theorem 1. Let the operator K be positive definite and $a_0(s) \neq 0$. Then every eigenvalue of the operator K_0 is a bifurcation point for equation (1).

Theorem 2. Let the operator K have a finite number of negative eigenvalues and $a_0(s) = c \neq 0$. Then for $c < 0$ every positive, and for $c > 0$ every negative, eigenvalue of the operator K_0 is a bifurcation point for equation (1).

3. Suppose now that

$$f(s, u) = a_\infty(s)u + \omega(s, u), \quad (5)$$

where $a_\infty(s) \in E'/E$, and $\omega(s, u)$ is a function satisfying the Carathéodory conditions, with

$$\lim_{u \rightarrow \infty} \frac{\omega(s, u)}{u} = 0, \quad (6)$$

$$|\omega(s, u)| \leq a(s) + b(s)|u|, \quad (7)$$

where $a(s) \in E'$, $b(s) \in E'/E$ (under these assumptions, obviously, the superposition operator $\omega x = \omega[s, x(s)]$ acts from E into E'). Denote by K_∞ the linear integral operator with kernel $k(t, s)a_\infty(s)$.

Assume that one of the following conditions is satisfied:

- a) the operator K is completely continuous;
- b) the operator f is w -bounded (or the operator G is w -bounded, and the operator f is uniformly continuous); $a_\infty(s) \in (E'/E)^0$; $b(s) \in (E'/E)^0$;

c) $E = M_{u_0}$.

Each of these conditions guarantees that the Krasnosel'skii operator Γ has a weakly continuous and smooth potential—the Golomb functional Φ . Moreover, under any of these conditions the operator Γ has the completely continuous asymptotic derivative $B_\infty = C' A_\infty C$, where A_∞ is the operator of multiplication by the function $a_\infty(s)$. Therefore, using somewhat modified results from (3), we obtain the following assertions.

Theorem 3. *Let the operator K be positive definite and $a_\infty(s) \neq 0$. Then every eigenvalue of the operator K_∞ is a bifurcation point for equation (1).*

Theorem 4. *Let the operator K have a finite number of negative eigenvalues and let $a_\infty(s) = c \neq 0$. Then for $c < 0$ every positive eigenvalue, and for $c > 0$ every negative eigenvalue of the operator K_∞ , is a bifurcation point for equation (1).*

4. Let the superposition operator $fx = f[s, x(s)]$ act from one ideal space E_1 into another E_2 . In the case of a regular (see (5)) space E_2 , the requirement that it be w -bounded is equivalent to the relation

$$\lim_{\text{mes } D \rightarrow 0} \sup_{\|x\| \leq r} \|P_{Dfx}\| = 0 \quad (8)$$

(P_D is the operator of multiplication by the characteristic function of the set $D \subset \Omega$), and the requirement of uniform continuity is equivalent to the relation

$$\lim_{\text{mes } D, \rho \rightarrow 0} \sup_{\|x\|, \|y\| \leq r; \|x-y\| \leq \rho} \|P_{Dfx} - P_{Dfy}\| = 0. \quad (9)$$

Therefore, in the case under consideration, uniform continuity is a weaker restriction. If the space E_2 does not possess the regularity property, the requirements of w -boundedness and uniform continuity are independent.

Consider, in particular, the case when E_1 is the Orlicz space $\mathcal{L}_M$, generated by the N -function $m(s, u)$, and E_2 is the Orlicz space $\mathcal{L}_N$, generated by the N -function $n(s, u)$.

Lemma 1. *The superposition operator $fx = f[s, x(s)]$ acting from $\mathcal{L}_M$ into $\mathcal{L}_N$ is w -bounded if there exists a function $\Psi(u)$ satisfying the condition*

$$\lim_{u \rightarrow \infty} \frac{\Psi(u)}{u} = \infty, \quad (10)$$

such that for every $r > 0$ one can specify $a_r \in \mathcal{L}_1$, $b_r, \lambda_r > 0$, such that

$$\Psi\{n[s, \lambda_r f(s, u)]\} \leq a_r(s) + b_r m(s, u/r). \quad (11)$$

Lemma 2. The superposition operator acting from $\mathcal{L}_m$ to $\mathcal{L}_N$, $fx = f[s, x(s)]$, is uniformly continuous if and only if, for some $c > 0$ and arbitrary $\varepsilon, r > 0$, one can specify $a_{\varepsilon, r} \in \mathcal{L}_1$, $b_{\varepsilon, r} > 0$ such that

$$n \left[s, \frac{f(s, u) - f(s, v)}{\varepsilon} \right] \leq a_{\varepsilon, r}(s) + m[s, b_{\varepsilon, r}(u - v)] + c \left[m \left(s, \frac{u}{r} \right) + m \left(s, \frac{v}{r} \right) \right]. \quad (12)$$

Lemmas 1 and 2 also cover the cases when the functions $n(s, u)$ do not satisfy the Δ_2 -condition.

5. In conclusion, we note that the results of the paper carry over to equations with a Lebesgue integral with respect to an arbitrary measure, in particular to infinite systems; in this case, instead of Banach spaces, one may consider locally convex spaces.

Leningrad State Pedagogical Institute
named after A. I. Herzen

Received
28 IV 1969

CITED LITERATURE

1. P. P. Zabreiko, A. I. Povolotskii, DAN, 183, No. 4 (1968).
2. M. A. Krasnosel'skii, *Topological Methods in the Theory of Nonlinear Integral Equations*, 1956.
3. M. A. Krasnosel'skii, A. I. Povolotskii, DAN, 91, No. 1, 19 (1953).
4. M. M. Vainberg, *Variational Methods for the Study of Nonlinear Operators*, 1956.
5. P. P. Zabreiko, Tr. seminara po funktsional' n. analizu, vol. 8 (1966).
6. P. P. Zabreiko, *Studies in the Theory of Integral Operators in Ideal Spaces of Functions*, Doctoral dissertation, Voronezh, 1968.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.