

MULTIPLIER TRANSFORMATIONS FOR PSEUDODIFFER- ENTIAL OPERATORS IN (L_p)

K. Tel'ner

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.14251>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.432

MATHEMATICS

K. TELFER

MULTIPLIER TRANSFORMATIONS FOR PSEUDODIFFERENTIAL OPERATORS IN L_p

(Presented by Academician V. M. Smirnov on 10 X 1969)

1. By a multiplier transformation we shall mean an operation that assigns to every pseudodifferential (ps.d.) operator K with symbol $K(\xi, x)$, i.e.

$$(Ku)(x) = (2\pi)^{-m/2} \int_{R^m} K(\xi, x) e^{i\langle \xi, x \rangle} \hat{u}(\xi) d\xi, \quad (1)$$

a new ps.d. operator ΦK with symbol $\varphi(\xi, x)K(\xi, x)$, i.e.

$$(\Phi Ku)(x) = (2\pi)^{-m/2} \int_{R^m} \varphi(\xi, x) K(\xi, x) e^{i\langle \xi, x \rangle} \hat{u}(\xi) d\xi. \quad (2)$$

Here R^m is m -dimensional Euclidean space; x and ξ are points of R^m ; $\langle \xi, x \rangle = \xi_1 x_1 + \dots + \xi_m x_m$; $\hat{u}$ is the Fourier transform of the function $u \in S$, where S is the space of rapidly decreasing functions. For symbols of ps.d. operators homogeneous of degree 0 (with respect to ξ), an operator of the form (1) is a singular integral (s.i.) operator.

The question is posed: under what conditions does the transformation $K \rightarrow \Phi K$ preserve the continuity of ps.d. operators in various function spaces? For the spaces $L_2(R^m)$ and $W_2^s(R^m)$ of Sobolev-Slobodetskii this problem was investigated in papers ⁽¹⁻⁵⁾. The purpose of the present note is to report two results on multiplier transformations in the spaces $L_p(R^m)$.

2. We shall give a description of the function spaces in terms of which the conditions of Theorems 1 and 2 will be expressed.

- a) Introduce the rings $\mathfrak{A}_l = \{\xi \in R^m : 2^{l-2} \leq |\xi| \leq 2^{l+2}\}$, $l = 0, \pm 1, \dots$. By $L_p^{\alpha, \beta}$ ($\alpha, \beta \geq 0$) we denote the space of all generalized functions $\varphi(\xi)$ belonging to $L_p^\alpha(\mathfrak{A}_l)$ (see ⁽¹⁾) for every l , and for which the seminorm $\|\varphi\|_{L_p^{\alpha, \beta}}$ is finite, where

$$\|\varphi\|_{L_p^{\alpha,\beta}}^p = \sum_{l=-\infty}^{\infty} 2^{l\beta p} \|\varphi\|_{L_p^{\alpha}(\mathfrak{A}_l)}^p.$$

If $\alpha = N$ is an integer, then $\|\varphi\|_{L_p^{N,\beta}}$ is equivalent to the integral seminorm

$$\left(\sum_{|\nu|=N} \int_{R^m} |\xi|^{|\beta p|} |D^\nu \varphi(\xi)|^p d\xi \right)^{1/p}.$$

b) By $L_{\infty;x}(L_p^{\alpha,\beta}; \xi)$ we shall denote the space of all functions $\varphi(\xi, x)$ belonging for almost all $x \in R^m$ to the space $L_p^{\alpha,\beta}$ (with respect to ξ) and for which the seminorm is finite

$$\|\varphi\|_{L_{\infty;x}(L_p^{\alpha,\beta}; \xi)} \equiv \text{vrai sup}_{x \in R^m} \|\varphi(\cdot, x)\|_{L_p^{\alpha,\beta}}.$$

c) Let S^{m-1} be the unit sphere of the space R^m ; θ its points. By $W_p^\alpha(S^{m-1})$ we denote the Sobolev-Slobodetskii space of functions $\varphi(\theta)$ defined on the sphere S^{m-1} . The space $L_{\infty;x}(W_{p;\theta}^\alpha(S^{m-1}))$ is defined by analogy with item b).

3. We formulate the results of the note.

Theorem 1. Suppose that the following two conditions are satisfied:

$$(I) \quad \varphi(\xi, x) \in L_{\infty;x}(W_{2;\xi}^\alpha(R^m)) \quad \text{for some } \alpha > m/2.$$

$$(II) \quad \varphi(\xi, x) \in L_{\infty;x}(L_{2;\xi}^{m/2+\beta,\beta}) \quad \text{for some } \beta > 0.$$

If $|1/p - 1/2| < \beta/m$ ($1 < p < \infty$), then the following conclusion is valid:

For every operator K of the form (1) with locally summable symbol $K(\xi, x)$, continuous in $L_p(R^m)$, the operator ΦK of the form (2) is also continuous in $L_p(R^m)$. Moreover

$$\|\Phi K\|_{L_p \rightarrow L_p} \leq C \left(\|\varphi\|_{L_\infty(W_2^\alpha)} + \|\varphi\|_{L_\infty(L_2^{m/2+\beta,\beta})} \right) \|K\|_{L_p \rightarrow L_p}.$$

Theorem 2. Let the function $\varphi(\xi, x)$ be homogeneous of degree 0 in ξ . If *

$$\varphi(\theta, x) \in L_{\infty;x}(W_{2;\theta}^\alpha(S^{m-1})), \quad \text{then, for } \alpha > (m-1)/2 + m|1/p - 1/2|$$

$$(1 < p < \infty)$$

the conclusion of Theorem 1 is valid. Moreover

$$\|\Phi K\|_{L_p \rightarrow L_p} \leq C \|\varphi\|_{L_\infty(W_2^s)} \|K\|_{L_p \rightarrow L_p}.$$

4. In the space $L_2(R^m)$ the multiplier problem formulated above is included in the theory of double operator integrals ^(1, 4), the use of which leads to various results. In studying the same problem in the spaces $L_p(R^m)$, the general methods of Hilbert space applied in ⁽¹⁻⁴⁾ are unsuitable. However, some elements of the method of integral sums developed in ⁽¹⁾ can be used in proving Theorems 1 and 2. The idea consists in approximating the multipliers $\varphi(\xi, x)$ by functions of a special form—piecewise-polynomial approximations corresponding to suitable decompositions of the space R^m . This idea is combined with the application of new results of Petre ⁽⁶⁾ and Littman, McCarthy, and Rivière ⁽⁷⁾ on multipliers of Fourier integrals and on estimates of Littlewood–Paley type.
5. Since the symbol of the identity operator is equal to $K(\xi, x) \equiv 1$, our results contain the following assertion.

Corollary. *Under the conditions of Theorems 1 and 2, a ps.d. (s.i.) operator with symbol $\varphi(\xi, x)$ is bounded in $L_p(R^m)$.*

We note that the boundedness criterion for a s.i. operator in $L_p(R^m)$ that follows as a special result from Theorem 2, for $p < 2$, somewhat improves the result of S. G. Mikhlin ⁽⁸⁾, Theorem V.1.26). For $p > 2$ the conditions of Theorem 2 are more restrictive than the conditions of S. G. Mikhlin; this is connected with the fact that the latter are not of multiplier character.

Other boundedness conditions for ps.d. operators in $L_p(R^m)$ were indicated, in the case $p = 2$, by L. Hörmander ⁽⁹⁾; from this V. M. Kagan ⁽¹⁰⁾ derived an analogous criterion for arbitrary p . Both results likewise do not have multiplier character. Kagan's conditions and Theorem 1 of the present work do not cover one another.

* The restriction of the function $\varphi(\xi, x)$ to the sphere S^{m-1} (with respect to the first variable) will be denoted by $\varphi(\theta, x)$.

The author expresses sincere gratitude to M. Z. Solomyak for posing the problem and for his attention to the work.

Leningrad State University
named after A. A. Zhdanov

Received
1 X 1969

REFERENCES

1. M. Sh. Birman, M. Z. Solomyak, in: *Problems of Mathematical Physics*, vol. 2, L., 1967, p. 26.
2. M. Sh. Birman, M. Z. Solomyak, *Vestn. LGU*, No. 13, 21 (1967).
3. M. Sh. Birman, M. Z. Solomyak, *Vestn. LGU*, No. 1, 35 (1969).
4. M. Sh. Birman, M. Z. Solomyak, *Izv. Vyssh. Uchebn. Zaved., Matematika*, No. 9 (88) (1969).
5. K. Tölner, *Vestn. LGU*, No. 1 (1970).
6. J. Peetre, *Ricerche di Mat.*, 15, fasc. 1°, 3 (1966).
7. W. Littman, Ch. McCarthy, N. Rivière, *Studia Math.*, 30, No. 2, 193 (1968).
8. S. G. Mikhlin, *Multidimensional Singular Integrals and Integral Equations*, Moscow, 1963.
9. L. Hörmander, in: *Pseudo-differential Operators*, Moscow, 1967, p. 297.
10. V. M. Kagan, *Izv. Vyssh. Uchebn. Zaved., Matematika*, No. 6 (73), 35 (1968).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.