

ON A COMPACTIFICATION OF NONCOMPACT SYMMETRIC RIEMANNIAN SPACES

G. F. Kushner

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.20831>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 519.46

MATHEMATICS

G. F. KUSHNER

ON A COMPACTIFICATION OF NONCOMPACT SYMMETRIC RIEMANNIAN SPACES

(Presented by Academician I. G. Petrovskii on 10 VII 1969)

1. Let $\mathcal{E}$ be a noncompact symmetric Riemannian space of rank l , with maximal connected group of motions G and stationary subgroup U of some point x_0 of it.

Denote the Lie algebras of the groups G and U by $\mathfrak{G}$ and $\mathfrak{U}$; denote by $\mathcal{E}\langle\cdot, \cdot\rangle$ the invariant scalar product in the algebra $\mathfrak{G}$ that gives rise to the Riemannian metric. The orthogonal complement $\mathcal{L}$ of the subalgebra $\mathfrak{U}$ in the algebra $\mathfrak{G}$ is identified, as usual, with the tangent space of the space $\mathcal{E}$ at the point x_0 .

Let $\mathfrak{h}$ be some Cartan subalgebra of the space $\mathcal{E}$ contained in $\mathcal{L}$, $\Sigma = \Sigma(\mathfrak{h})$ its system of roots, and Z the set of integers. The set

$$I = I(\mathfrak{h}) = \left\{ \mu \in \mathfrak{h} \mid \frac{2\langle\mu, \alpha\rangle}{\langle\alpha, \alpha\rangle} \in Z \text{ for all } \alpha \in \Sigma \right\} \quad (1)$$

will be called the lattice of the space $\mathcal{E}$.

It is obvious that any two lattices of the space $\mathcal{E}$ are conjugate with respect to the adjoint group $\text{Ad}(G)$ of the group G .

Let $(\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$, where $1 \leq k \leq l$, be an orthonormal root of pairwise commuting vectors of the space $\mathcal{L}$. Choose some Cartan subalgebra $\mathfrak{h}$ of the space $\mathcal{E}$ so that

$$\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1} \in \mathfrak{h}\mathcal{L}, \quad (2)$$

and select from the system of roots $\Sigma = \Sigma(\mathfrak{h})$ and the lattice $I = I(\mathfrak{h})$ the subsystem

$$\Sigma^{(k)} = \Sigma(\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}) = \{\alpha \in \Sigma \mid \langle\alpha, \mathfrak{h}_s\rangle = 0 \text{ for all } 0 \leq s \leq k-1\} \quad (3)$$

and the sublattice

$$I^{(k)} = I(\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}) = \{\mu \in I \mid \langle \mu, \mathfrak{h}_s \rangle = 0 \text{ for all } 0 \leq s \leq k-1\}. \quad (4)$$

We shall call the tuple $(\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$ admissible if, for every s ($1 \leq s \leq k-1$), $\mathfrak{h}_s \in ((I(\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})))$ (the linear span of I).

2. To the space $\mathcal{E}$ with fixed point x_0 , by means of its root system Σ and lattice I , one can associate a family of symmetric Riemannian spaces, which we shall call the family of boundary spaces, or ε -spaces, of the space $\mathcal{E}$.

An arbitrary ε -space of the space $\mathcal{E}$ is constructed as follows. Let $(\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$, where $1 \leq k \leq l$, be an arbitrary admissible tuple; let $\mathfrak{h}$ be some Cartan subalgebra of the space $\mathcal{E}$ satisfying (2); let $G^{(k)} = G(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$ be the noncompact semisimple Lie group generated by the root subsystem (3); $U^{(k)} = U(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}) = G^{(k)} \cap U$, and $H_{(k)} = H_{(k)}(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$ the connected noncompact commutative group corresponding to the orthogonal complement $\mathfrak{h}_{(k)} = \mathfrak{h}(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$ of the space $((\Sigma^{(k)}))$ in the space $((I^{(k)}))$ (3,4).

The direct product $\widehat{G}^{(k)}$ of the groups $G^{(k)}$ and $H_{(k)}$

$$\widehat{G}^{(k)} = \widehat{G}(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}) = G^{(k)} \cdot H_{(k)} \quad (5)$$

does not depend on the choice of the Cartan subalgebra $\mathfrak{h}$ satisfying (2), and gives rise to the symmetric Riemannian space

$$S^{(k)} = S(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}) = \widehat{G}^{(k)} / U^{(k)} \quad (1 \leq k \leq l), \quad (6)$$

which is called the ε -space of the k -th degree of the space $\mathcal{E}$, corresponding to its admissible tuple $(\mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$.

The set of all ε -spaces (6), supplemented for generality of the discussion by the space

$$S^{(0)} = \mathcal{E}, \quad (7)$$

is the family of ε -spaces of the space $\mathcal{E}$.

Every ε -space $S^{(k)}$ which does not degenerate into a point has non-positive curvature and is homeomorphic to Euclidean space. As a Riemannian space, it decomposes into the direct product

$$S^{(k)} = \mathcal{E}^{(k)} \times E^{(k)} \quad (8)$$

of the noncompact symmetric space

$$\mathcal{E}^{(k)} = \mathcal{E}(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}) = G^{(k)} / U^{(k)}$$

and the Euclidean space

$$E^{(k)} = E(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}) = H_{(k)}/\{e\}$$

(e is the identity element of the group G).

To each k -pencil $F^{(k)}$ of geodesic ε -spaces of

$$S^{(k)} = S(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1})$$

(¹) there corresponds, in a one-to-one manner, the ε -space

$$S^{(k)}(x_0^{(k)}, F^{(k)}) = S(x_0, \mathfrak{h}_0, \dots, \mathfrak{h}_{k-1}, \mathfrak{h}_k), \quad (9)$$

where $\mathfrak{h}_k$ is the tangent vector with which the geodesic $\gamma^{(k)} \in F^{(k)}$ passes through the point $x_0^{(k)}$ with stationary subgroup $U^{(k)}$. The space (9) is called the ε -space of the k -pencil $F^{(k)}$ of geodesics of the space $S^{(k)}$.

3. From the set of all projective representations of the motion group G of the space $\mathcal{E}$, one can single out a subset θ of its so-called admissible representations, or ∂ -representations. A real projective representation φ of the group G is called its ∂ -representation if in its space P_φ there exists a point $\varphi(x_0)$ satisfying the conditions:

$$U \subseteq G(\varphi, x_0), \quad P_\varphi = P(\varphi, x_0), \quad (10)$$

where $G(\varphi, x_0)$ is the stationary subgroup of the point $\varphi(x_0)$ in the group G , and $P(\varphi, x_0)$ is the projective linear manifold of the space P_φ generated by the orbit $\mathcal{E}_\varphi$ of the group G passing through the point $\varphi(x_0)$. We note that the space $\mathcal{E}_\varphi$ is homeomorphic to the space $\mathcal{E}$. If the ∂ -representation φ satisfies the equality $U = G(\varphi, x_0)$, then the space $\mathcal{E}_\varphi$ is isomorphic to the space $\mathcal{E}$ and is called a projective representation of the space $\mathcal{E}$, while its closure $\overline{\mathcal{E}}_\varphi$ in the space P_φ is called a projective compactification (^{2,3}).

4. Let

$$P_\theta = \prod_{\varphi \in \theta} P_\varphi$$

be the direct product of the spaces of all ∂ -representations of the motion group G of the space $\mathcal{E}$, endowed with the Tikhonov topology. It is obvious that P_θ is a compact metrizable space.

Let $p = (p_\varphi)_{\varphi \in \theta}$ be an arbitrary point of the space P_θ , and let

$$\theta(x_0) = (\varphi(x_0))_{\varphi \in \theta}$$

(see 3). It is not hard to show that the formula

$$\theta(g)p = (\varphi(g)p_\varphi)_{\varphi \in \theta} \quad (p \in P_\theta, g \in G) \quad (11)$$

turns the group G into a topological group of transformations of the space P_θ , in which the stationary subgroup $G(\theta, x_0)$ of the point $\theta(x_0)$ coincides

with stationary subgroup U of the point x_0 . It follows from this that the orbit $\mathcal{E}_\theta$ of the group G , passing through the point $\theta(x_0)$, is a representation of the space $\mathcal{E}$, and its closure $\bar{\mathcal{E}}_\theta$ in the space P_θ is a compactification of this space.

For any e -space (6) and its factors (8) one can also construct representations and compactifications in the space P_θ . This is done as follows. In the space P_θ , by means of a chain of limiting transitions,

$$\theta(x'_0) = \lim_{t \rightarrow +\infty} \theta(\exp \mathfrak{h}_0 t) \theta(x_0), \dots, \theta(x_0^{(k)}) = \lim_{t \rightarrow \infty} \theta(\exp \mathfrak{h}_{k-1} t) \theta(x_0^{(k-1)})$$

one constructs the point $\theta(x_0^{(k)})$. The orbits $S_\theta^{(k)}, \mathcal{E}_\theta^{(k)}, E_\theta^{(k)}$ passing through it of the groups $\bar{G}^{(k)}, G^{(k)}, H_{(k)}$ are isomorphic to the spaces $S^{(k)}, \mathcal{E}^{(k)}, E^{(k)}$ and are called their representations generated by the compactification $\bar{\mathcal{E}}_\theta$; and the closures $\bar{S}_\theta^{(k)}, \bar{\mathcal{E}}_\theta^{(k)}, \bar{E}_\theta^{(k)}$ of these orbits in the space P_θ are compactifications of these spaces, also generated by the compactification $\bar{\mathcal{E}}_\theta$.

Let us describe the structure of the boundary $\mathcal{T}_\theta = \bar{\mathcal{E}}_\theta \setminus \mathcal{E}_\theta$ of the compactification $\bar{\mathcal{E}}_\theta$ and some of its connections with the geometry of the space $\mathcal{E}$. Let $S^{(k)}$ ($k \geq 0$) be an arbitrary e -space of the space $\mathcal{E}$ not degenerating to a point (6,7); let $\gamma^{(k)}$ be a geodesic of the space $S^{(k)}$; let $\mathfrak{h}_k$ be the tangent vector of the geodesic $\gamma^{(k)}$ at some point $x_0^{(k)}$, and let $\exp \mathfrak{h}_k t$ ($-\infty < t < +\infty$) be the group of translations of the space $S^{(k)}$ along the geodesic $\gamma^{(k)}$. The point

$$\theta(x_{\gamma^{(k)}}) = \lim_{t \rightarrow +\infty} \theta(\exp \mathfrak{h}_k t) \theta(x_0^{(k)}) \quad (12)$$

is naturally called the improper point of the geodesic $\gamma^{(k)}$ in the compactification $\bar{S}_\theta^{(k)}$.

Theorem 1. *All geodesics belonging to one and the same $\mathfrak{h}$ -bundle of the e -space $S^{(k)}$ (1) have, in the compactification $\bar{S}_\theta^{(k)}$, a common improper point.*

Let $F^{(k)}$ be an arbitrary k -bundle of geodesics of the space $S^{(k)}$ (1), and let $S^{(k)}(x_0^{(k)}, F^{(k)})$ be the corresponding e -space (9). We shall call the representation $S_\theta^{(k)}(x_0^{(k)}, F^{(k)})$ of the space $S^{(k)}(x_0^{(k)}, F^{(k)})$, generated by the compactification $\bar{\mathcal{E}}_\theta$, the component of the boundary corresponding to the k -bundle $F^{(k)}$.

Theorem 2. *The boundary component*

$$S_\theta^{(k)}(F^{(k)}) = S_\theta^{(k)}(x_0^{(k)}, F^{(k)})$$

of any k -bundle $F^{(k)}$ of geodesics of the space $S^{(k)}$ does not depend on the choice of the initial point $x_0^{(k)}$ and coincides with the set of improper points of all geodesics of this k -bundle in the compactification $\overline{S}_\theta^{(k)}$. It decomposes into the direct product

$$S_\theta^{(k)}(F^{(k)}) = \mathcal{E}_\theta^{(k)}(x_0^{(k)}, F^{(k)}) \times E_\theta^{(k)}(x_0^{(k)}, F^{(k)}) \quad (13)$$

of representations of the noncompact and Euclidean factors of the e -space $S^{(k)}(x_0^{(k)}, F^{(k)})$, and its foliations, generated by this decomposition, do not depend on the choice of the initial point $x_0^{(k)}$.

Let $\Phi^{(k)}$ be the set of all k -bundles of the e -space $S^{(k)}$ ($k \geq 0$), and let $\mathcal{T}_\theta^{(k)} = \overline{S}_\theta^{(k)} / S_\theta^{(k)}$.

Theorem 3. For any e -space $S^{(k)}$ ($k \geq 0$) not degenerating to a point,

$$\mathcal{T}_0^{(k)} = \bigcup_{F^{(k)} \in \Phi^{(k)}} \overline{S}_\theta^{(k)}(F^{(k)}), \quad (14)$$

where

$$\overline{S}_\theta^{(k)}(F_1^{(k)}) \cap \overline{S}_\theta^{(k)}(F_2^{(k)}) = \emptyset \quad \text{for } F_1^{(k)} \neq F_2^{(k)}. \quad (15)$$

It follows directly from Theorem 3 that

Theorem 4. The family of representations of all k -spaces of the space $\mathcal{E}$ of nonzero degrees (6), generated by the compactification $\mathcal{E}_\theta$, is a decomposition of its boundary $\mathcal{T}_\theta$.

5. Let us compare the compactification $\mathcal{E}_\theta$ with the compactification $\overline{\mathcal{E}}$ of F. I. Karpelevich¹.

Theorem 5. For any noncompact symmetric Riemannian space $\mathcal{E}$, the natural mapping $\theta^{-1} : \mathcal{E}_\theta \rightarrow \mathcal{E}$ can be extended to a continuous surjection $\overline{\theta}^{-1} : \overline{\mathcal{E}}_\theta \rightarrow \overline{\mathcal{E}}$.

Theorem 6. If the rank of the space $\mathcal{E}$ is greater than 1, then it is impossible to extend the natural mapping $\theta : \mathcal{E} \rightarrow \mathcal{E}_\theta$ to a continuous mapping $\overline{\theta} : \overline{\mathcal{E}} \rightarrow \overline{\mathcal{E}}_\theta$.

Remark 1. For spaces $\mathcal{E}$ of rank 0 or 1, the compactifications $\mathcal{E}_\theta$ and $\overline{\mathcal{E}}$ are equivalent.

Remark 2. From Theorems 5 and 6 it is clear that the compactification $\overline{\mathcal{E}}_\theta$ has a more developed boundary than the compactification $\overline{\mathcal{E}}$.

The author thanks P. K. Rashevskii for posing the problem and for guidance in its solution.

Astrakhan
State Pedagogical Institute

Received
2 VI 1969

REFERENCES

- ¹ F. I. Karpelevich, Tr. Mosk. matem. obshch., **14**, 48 (1965).
- ² P. K. Rashevskii, Matem. sborn. **50** (92), 2, 171 (1960).
- ³ I. Satake, Sborn. per. Matematika, **5**, 3, 45 (1961).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.