

ON THE APPLICATION OF THE METHOD OF FAST MATRIX MULTIPLICATION TO THE PROBLEM OF FINDING THE TRANSITIVE CLOSURE OF A GRAPH

M. E. Furman

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.25427>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 518.5

MATHEMATICS

M. E. FURMAN

ON THE APPLICATION OF THE METHOD OF FAST MATRIX MULTIPLICATION TO THE PROBLEM OF FINDING THE TRANSI- TIVE CLOSURE OF A GRAPH

(Presented by Academician I. G. Petrovskii, 10 III 1970)

This algorithm makes it possible to find the transitive closure of a graph in $cn^{\log_2 7} \lg n^*$ operations, which is asymptotically better than the previously known estimate $cn^3 / \lg n$ ⁽¹⁾.

Lemma. *Boolean $n \times n$ matrices can be multiplied in $cn^{\log_2 7}$ operations.*

Indeed, we shall regard these matrices as arithmetic ones consisting of zeros and ones, and multiply them by the method ⁽²⁾ in $cn^{\log_2 7}$ operations. Then, in cn^2 operations, we inspect the elements of the product and replace every nonzero element by one. It is obvious that in this way we obtain the Boolean product of the matrices.

Theorem. *The transitive closure of a graph of order n can be obtained in $cn^{\log_2 7} \lg n$ operations.*

Proof. Let A_0 be the adjacency matrix of the given graph. We shall obtain the adjacency matrix of the transitive closure by the following iterations: $A_{n+1} = A_n \vee A_n^2$.

One iteration requires $cn^{\log_2 7}$ operations, and the number of iterations required is $\lg K$, where K is the maximum distance between two vertices (the diameter of the original graph), since each iteration halves the diameter of the graph. But $K \leq n$; thus, we required fewer than $cn^{\log_2 7} \lg n$ operations. This method is more advantageous than the method ⁽¹⁾ for values of n on the order of 10^{19} .** However, if better estimates are obtained for the number of operations required for matrix multiplication, then this algorithm will also become advantageous for realistic n .

Moscow State Pedagogical Institute
named after V. I. Lenin

Received
4 III 1970

REFERENCES

¹ V. L. Arlazarov, E. A. Dinitz, M. A. Kronrod, I. A. Faradzhev. DAN, 194, No. 3 (1970).

² V. Strassen, *Numerische Math.*, **13**, No. 4, 354 (1969).

* $\lg x$ is the least integer greater than or equal to $\log_2 x$; n is the number of vertices of the graph.

** With equal constants entering as factors in the estimate of the number of operations.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.