

ENTROPY OF A SHIFT AND MARKOV MEASURES IN THE PATH SPACE OF A COUNTABLE GRAPH

B. M. Gurevich

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.37210>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 519.217+519.53

MATHEMATICS

B. M. GUREVICH

ENTROPY OF A SHIFT AND MARKOV MEASURES IN THE PATH SPACE OF A COUNTABLE GRAPH

(Presented by Academician A. N. Kolmogorov, 4 XII 1969)

Let G be a directed graph with a countable set of vertices, which henceforth will be identified with the natural numbers. Denote the adjacency matrix of the graph G by $\Pi(G) = \Pi = (\pi_{ij})$. We shall assume that the graph is connected, i.e., any two of its vertices can be joined by a path. The length of a path is the number of edges in it. If the lengths of all closed paths (cycles) of the graph G are relatively prime, then the graph is called **aperiodic**. An infinite two-sided sequence of vertices $\{i_k, -\infty < k < \infty\}$ forms an infinite path if, for every k , there is an edge leading from i_k to i_{k+1} , i.e. $\pi_{i_k i_{k+1}} = 1$. Denote by $\Omega(G)$ the totality of all infinite paths of the graph G .

Let M be the set of numbers of the form $0, 1/n$ ($n = 1, 2, \dots$), and let $\Omega(M)$ be the set of infinite two-sided sequences of elements of M . If M is regarded as a compact subset of the line, then $\Omega(M)$ with the topology of the direct product will be a compact topological space. The mapping $n \mapsto 1/n$ induces an embedding of $\Omega(G)$ into $\Omega(M)$ and thereby defines a topology in $\Omega(G)$. The same topology can be obtained if one introduces the discrete topology on the set N of natural numbers and regards $\Omega(G)$ as a subset of the space $\Omega(N)$ of two-sided sequences of natural numbers with the topology of the direct product. Obviously, $\Omega(G)$ is closed in $\Omega(N)$ and not closed in $\Omega(M)$. Denote its closure in $\Omega(M)$ by $\overline{\Omega}(G)$.

In each of the spaces mentioned one can define the shift, which takes the sequence $\omega = \{\omega_k\}$ to $\omega' = \{\omega'_k\}$, where $\omega'_k = \omega_{k-1}$, and which is a homeomorphism. The shift $T(G)$ in the space $\Omega(G)$ is called a **topological Markov chain** (see ^(1,2)).

From the theorem proved in ⁽²⁾ it follows that the topological entropy (see ⁽³⁾) of the shift in the compact space $\overline{\Omega}(G)$ coincides with the upper bound of the metric entropies over all normalized invariant Borel measures concentrated on $\Omega(G)$.

Denote by $h(G)$ the common value of these two quantities and define $\lambda(G)$ by

the condition $h(G) = \log \lambda(G)$ (the base of the logarithms is the same as in the definition of entropy). A measure μ in the space $\Omega(G)$ for which the metric entropy of the shift $h_\mu(T(G))$ is equal to $h(G)$ will be called a **measure with maximal entropy**. The example contained in ⁽²⁾ shows that, in the case under consideration, unlike the case of a finite graph (see ^(4,5)), there need not exist a measure with maximal entropy. In this note we give necessary and sufficient conditions for the existence and uniqueness of such a measure.

1. Special graphs. First consider a graph G whose adjacency matrix $\Pi(G) = \Pi = (\pi_{ij})$ is determined by a nondecreasing sequence of natural numbers $\{k_s, s = 1, 2, \dots\}$ according to the rule: $\pi_{ij} = 1$ if

$$i \neq \sum_{s=1}^n k_s, \quad j = i + 1$$

or

$$i = \sum_{s=1}^n k_s, \quad j = 1 + \sum_{s=1}^m k_s, \quad n \geq 1, \quad m \geq 0$$

(we agree that $\sum_{s=1}^0 k_s = 0$), and $\pi_{ij} = 0$ in all other cases. Graphs of the described—

graphs of this form will be called **special**. Such graphs, in connection with the problem of interest to us, were first considered by E. I. Dinaburg ^{(6)*}.

Theorem 1. *If G is a special graph defined by the sequence $\{k_s\}$, and $r(G) = r$ is the radius of convergence of the series*

$$\varphi(t) = \sum_{s=1}^{\infty} t^{k_s},$$

then $\lambda(G) = r^{-1}$ for $\varphi(r) \leq 1$ and $\lambda(G) = t_0^{-1}$ for $\varphi(r) > 1$, $\varphi(t_0) = 1$, $t_0 > 0$.

The proof is obtained by comparing the results of ^(2,6).

Let us note that the shift in the path space of a special graph serves as a topological analogue of a special automorphism (see, for example, ⁽⁷⁾), constructed from a Bernoulli automorphism with a countable number of states and a function depending only on the state at time zero and equal to k_n on the n -th state.

Theorem 2. *Let, in the notation of the preceding theorem, $r > 0$. Then, for the existence of a measure with maximal entropy, it is necessary and sufficient that one of the following two conditions hold:*

$$1) \varphi(r) > 1; \quad 2) \varphi(r) = 1, \quad \varphi'(r) < \infty.$$

If a measure with maximal entropy exists, then it is unique, and after the introduction of this measure the shift $T(G)$ becomes a special automorphism constructed from a Bernoulli automorphism with states $E_1, E_2, \dots$ and probabilities $p(E_n) = (\lambda(G))^{-k_n}$, $n = 1, 2, \dots$, and the function $f(E_n) = k_n$.

In the proof the following simple

Lemma. *Let*

$$\psi(x) = \sum_{n=1}^{\infty} a_n x^n, \quad \text{where } a_n \geq 0.$$

Then

$$x[\psi'(x)]^2 - \psi(x)\psi'(x) - x\psi(x)\psi''(x) \leq 0,$$

and equality is possible only in the case when $\psi(x) = cx^k$.

2. The general case. We now pass to the study of an arbitrary graph G , assuming only that $\lambda(G) < \infty$. From this assumption it follows that for any vertex i and any $n > 0$ there is only a finite number of cycles of length n passing through i (see ⁽²⁾, Lemma 3). Fixing i , enumerate, in increasing order of their lengths, all cycles passing through i exactly once (cycles of the same length may be numbered in an arbitrary order). Let $k_s(i)$ be the length of the s -th cycle in our enumeration, and let $G(i)$ be the special graph defined by the sequence $\{k_s(i)\}$. We shall call $G(i)$ the **special representation** of the graph G corresponding to the vertex i .

Using the special representation and the results of Section 1, one can establish the following facts:

Theorem 3. *Let G be a connected graph and $\lambda(G) < \infty$. Then: 1) if a measure with maximal entropy exists, then it is unique; 2) if i is an arbitrary vertex of the graph G and $G(i)$ is the special representation corresponding to it, then $T(G)$ and $T(G(i))$ possess measures with maximal entropy simultaneously; 3) if μ_0 is a measure with maximal entropy, then μ_0 is a Markov measure positive on every open set (the latter means that the random variables $x_n(\omega) = \omega_n$, $\omega = (\dots, \omega_{-1}, \omega_0, \omega_1, \dots) \in \Omega(G)$, form a Markov chain).*

Corollary. *If G is a connected aperiodic graph, $\lambda(G) < \infty$, and μ_0 is a measure with maximal entropy, then $(T(G), \mu_0)$ is a K -automorphism.*

Let us give one more fact, useful for applications and following from the results of ⁽²⁾.

Theorem 4. *If G is a connected graph, then for every $c > 0$ there exists an ergodic Markov measure, positive on every open set,*

* Taking the occasion, I note at the request of the author of ⁽⁶⁾ that, for the validity of assertion 4) from that paper, the condition $f(k) > c \ln k$ must be replaced by $f(k) < c(\ln k)^a$, $a > 1$.

measure μ_c such that $h_\mu(T(G)) > h(T(G)) - c$ when $\lambda(G) < \infty$, and $h_{\mu_c}(T(G)) > c$ when $\lambda(G) = \infty$.

3. Relation with the matrix $\Pi(G)$. As in the case of a finite graph G (see ^(4, 5)), the entropy $h(G)$ and the measure with maximal entropy, if it exists, can be directly related to the matrix $\Pi(G)$. Suppose that $\lambda(G) < \infty$. Then all powers $\Pi^n = (\pi_{ij}^{(n)})$, $n > 0$, of the matrix $\Pi = \Pi(G)$ are defined (see ⁽²⁾, Lemma 3), and the radius of convergence $R(G)$ of the series

$$\sum_n \pi_{ij}^{(n)} t^n$$

does not depend on i and j (see ⁽⁸⁾).

Theorem 5. *If G is a connected graph, $\lambda(G) < \infty$, and $R = R(G)$, then*

$$h(G) = \log \lambda(G) = -\log R.$$

The number $R(G)$ is called the convergence parameter of the matrix $\Pi(G)$, and $[R(G)]^{-1}$ serves as an analogue of the maximal eigenvalue (see ⁽⁹⁾). If $R(G) = R$ and $\pi_{ij}^n R^n \rightarrow 0$, i.e. the matrix $\Pi(G)$ is, by definition, R -recurrent and R -positive, then R^{-1} is a genuine eigenvalue: there exist sequences of positive numbers $x(G) = x = (x_1, x_2, \dots)$, $y(G) = y = (y_1, y_2, \dots)$, such that

$$\sum_j \pi_{ij} x_j = (1/R) x_i,$$

$$\sum_i \pi_{ij} y_i = (1/R) y_j,$$

and the vectors x, y are determined uniquely up to a factor, and

$$\sum_i x_i y_i < \infty$$

(see ⁽⁹⁾).

Theorem 6. *If G is a connected graph and $\lambda(G) = 1/R < \infty$, then for the existence of a measure μ_0 with maximal entropy it is necessary and sufficient that the matrix $\Pi(G)$ be R -recurrent and R -positive. Under these conditions μ_0 is the Markov measure corresponding to transition probabilities*

$$p_{ij} = x_j \pi_{ij} / \lambda(G) x_i, \quad i, j = 1, 2, \dots,$$

and stationary probabilities

$$p_i = x_i y_i,$$

where the vectors $x(G) = x$, $y(G) = y$ are normalized so that

$$\sum_i x_i y_i = 1.$$

4. Application to Y -systems. The results of § 2 find application in the theory of classical dynamical systems. In particular, with their help the following can be proved.

Theorem 7. *Let T be a C^2 -diffeomorphism of a closed n -dimensional Riemannian manifold M , possessing an integral invariant and satisfying the condition Y of D. V. Anosov ⁽¹⁰⁾. Then on M there exists a unique invariant, relatively T -normalized, Borel measure μ_0 , for which the metric entropy $h_{\mu_0}(T)$ assumes the maximum of the possible values, equal to the topological entropy (see ^(3, 11)). The transformation T of the space M with measure μ_0 is a K -automorphism.*

Remark. The existence of a measure with maximal entropy for a Y -diffeomorphism was earlier proved by R. Bowen ⁽¹²⁾.

The author thanks E. I. Dinaburg for useful conversations.

Moscow State University
named after M. V. Lomonosov

Received
21 XI 1969

CITED LITERATURE

1. V. M. Alekseev, UMN, **23**, no. 2 (1968).
2. B. M. Gurevich, DAN, **187**, no. 4 (1969).
3. R. L. Adler, A. G. Konheim, M. H. McAndrew, Trans. Am. Math. Soc., **114**, no. 2 (1965).

4. W. Parry, *ibid.*, **112**, no. 1 (1964).
5. R. L. Adler, B. Weiss, *Proc. Nat. Acad. Sci. U.S.A.*, **57**, no. 6 (1967).
6. E. I. Dinaburg, *UMN*, **23**, no. 4 (1968).
7. B. M. Gurevich, *DAN*, **153**, no. 4 (1963).
8. D. Vere-Jones, *Quart. J. Math. (Oxford)*, Ser. 13 (1962).
9. D. Vere-Jones, *Pacific J. Math.*, **22**, no. 2 (1967).
10. D. V. Anosov, *Tr. Matem. inst. im. V. A. Steklova AN SSSR*, **90** (1967).
11. E. I. Dinaburg, *DAN*, **190**, no. 1 (1970).
12. R. Bowen, Preprint. Univ. of Warwick, Coventry, 1969.

* In the presence of an integral invariant, each leaf of the expanding and contracting foliations corresponding to T is everywhere dense in M (see ⁽¹⁰⁾). Only this is needed for the proof of the last assertion of Theorem 7.

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.