

ON THE INDEX OF A PSEUDODIFFERENTIAL OPERATOR WITH A FINITE GROUP OF SHIFTS

A. B. Antonevich

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.39059>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 513.88 : 513.83 + 517.948

MATHEMATICS

A. B. ANTONĖVICH

ON THE INDEX OF A PSEUDODIFFERENTIAL OPERATOR WITH A FINITE GROUP OF SHIFTS

(Presented by Academician I. G. Petrovskii, 20 VI 1969)

Let G be a finite group of order N ; let X be a connected G -manifold and let the action of the group G on X be effective. In the space $C_\infty(X)$, by the formula

$$T_g u(x) = u(g^{-1}x), \quad g \in G, \quad x \in X, \quad u \in C_\infty(X),$$

we define a representation T_g of the group G .

By a pseudodifferential operator of order r with a finite group of shifts we shall mean an operator A acting in $C_\infty(X)$ by the formula

$$Au = \sum_{g \in G} A_g T_g u,$$

where the A_g are pseudodifferential operators of order r . The operator A belongs to the operators with a deviating argument. In the present paper conditions for the Noether property of the operator A are indicated and a formula for its index is given.

Let $\tilde{A}$ be the operator acting in the space $C_\infty(X \times G)$ by the formula

$$\tilde{A}u(x, g') = \sum_{g \in G} T_{g'} A_g T_g^{-1} u(x, g'g).$$

If functions from $C_\infty(X \times G)$ are regarded as vector-functions on X , then the operator $\tilde{A}$ is a G -invariant pseudodifferential operator ⁽¹⁾.

Theorem 1. *In order that the extension of the operator A be a Noether operator from $H_l(X)$ into $H_{l-r}(X)$, it is necessary and sufficient that the operator $\tilde{A}$ be elliptic.*

Proof. In the space $C_\infty(X \times G)$ define a representation $\tilde{T}_g$ of the group G

$$\tilde{T}_{g''}u(x, g') = u(g''^{-1}x, g''^{-1}g').$$

The space $C_\infty(X \times G)$ decomposes into a direct sum of a finite number of invariant subspaces M^i ($i = 1, \dots, k$) such that in M^i there acts a representation which is a multiple of the irreducible representation D^i of the group G . The operator $\tilde{A}$ commutes with all operators $\tilde{T}_{g''}$ and, consequently, the subspaces M^i are invariant with respect to the operator $\tilde{A}$, i.e. the operator $\tilde{A}$ decomposes into the direct sum of operators A_i , $i = 1, \dots, k$. Let M^1 and A_1 be the subspace and the operator corresponding to the identity representation of the group G . The mapping $p : M^1 \rightarrow C_\infty(X)$, acting by the formula

$$pu(x) = u(x, e),$$

is an isomorphism and carries the operator A_1 into A . The remaining operators A_i can also be realized in $C_\infty^{n_i}(X)$ as pseudodifferential operators with shift (n_i is the dimension of the representation D^i).

Let now $\tilde{A}$ be an elliptic operator. Then it can be extended to a Noetherian operator acting from $H_l^N(X)$ to $H_{l-r}^N(X)$, and consequently the operator A is also extended to a Noetherian operator from $H_l(X)$ to $H_{l-r}(X)$. Suppose that $\tilde{A}$ is not an elliptic operator. Then one can construct a function $u_\lambda \in M^1$ such that, as $\lambda \rightarrow \infty$, the a priori estimate is not satisfied for u_λ , which contradicts the Noetherian property of the operator A . The theorem is proved.

In work ¹ the analytic index of a G -invariant operator is defined as an element of the ring $R(G)$ of characters of representations of the group G :

$$\text{index } B = [\text{Ker } B] - [\text{Coker } B] \in R(G),$$

where $[\text{Ker } B]$, $[\text{Coker } B]$ are the characters of the representations induced by the action of the group G in $\text{Ker } B$ and $\text{Coker } B$. The value of $\text{index } B$ at an element g is called the Lefschetz number and is denoted by $L(g, B)$. In ¹⁻³ a method is indicated for computing the index for pseudodifferential operators.

Theorem 2. *The index of a pseudodifferential operator A with a finite group of shifts is expressed by the formula*

$$\text{ind } A = \frac{1}{N} \sum_{g \in G} L(g, \tilde{A}). \quad (1)$$

Proof. From the decomposition of the operator $\tilde{A}$ we obtain

$$\text{index } \tilde{A} = \sum_{i=1}^k \text{index } A_i, \quad (2)$$

and since in M^i there acts a representation that is a multiple of the irreducible representation D^i , we have

$$\text{index } A_i = \frac{\text{ind } A_i}{n_i} \chi_i, \quad (3)$$

where χ_i is the character of the irreducible representation D^i , $\text{ind } A_i = \dim \text{Ker } A_i - \dim \text{Coker } A_i$. By virtue of the orthogonality of the characters of irreducible representations, from formulas (2) and (3) we obtain

$$\text{ind } A_i = \frac{n_i}{N} \sum_{g \in G} L(g, \tilde{A}) \overline{\chi_i}(g). \quad (4)$$

From (4), for $i = 1$, we obtain the assertion of the theorem.

Theorem 3. *If every transformation $g \in G$, $g \neq e$, has a finite number of fixed points, then*

$$\text{ind } A = \frac{1}{N} \text{ind } \tilde{A}. \quad (5)$$

Proof. If g has a finite number of fixed points, then in work ² $L(g, \tilde{A})$ is expressed in terms of the trace of the operator $\tilde{T}_g$ in the fiber over a fixed point. Since in the fiber over a fixed point $\tilde{T}_g$ acts as the regular representation, $\text{Tr } T_g = 0$, $L(g, \tilde{A}) = 0$, and formula (1) gives (5).

The results of the article are easily transferred to operators in sections of vector bundles. In the case of one-dimensional singular integral operators and the cyclic group G , formula (5) was obtained by other methods in works ^{4,5}.

Belorussian State University
named after V. I. Lenin
Minsk

Received
11 VI 1969

CITED LITERATURE

- ¹ M. F. Atiyah, I. M. Singer, *UMN*, **23**, no. 5 (143), 99 (1968).
- ² M. F. Atiyah, G. B. Segal, *UMN*, **23**, no. 6 (144), 135 (1968).
- ³ M. F. Atiyah, I. M. Singer, *UMN*, **24**, no. 1 (145), 127 (1969).
- ⁴ G. S. Litvinchuk, *Izv. AN SSSR, ser. matem.*, **31**, 563 (1967).
- ⁵ G. S. Litvinchuk, *Izv. AN SSSR, ser. matem.*, **32**, 1414 (1968).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.