

ON THE HADAMARD WELL-POSEDNESS OF THE CAUCHY PROBLEM FOR AN EVOLUTION EQUATION

S. G. Krein, G. I. Laptev, G. A. Tsvetkova

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.40672>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.948:513.88

MATHEMATICS

S. G. KREIN, G. I. LAPTEV, G. A. TSVETKOVA

ON THE HADAMARD WELL-POSEDNESS OF THE CAUCHY PROBLEM FOR AN EVOLUTION EQUATION

(Presented by Academician I. N. Vekua on 27 X 1969)

The uniformly well-posed Cauchy problem for an evolution equation in a Banach space has been well studied (see ⁽¹⁾). If one considers systems of partial differential equations with constant coefficients of the form $\partial u / \partial t = A(D)u$, then necessary and sufficient conditions for the uniform well-posedness of the Cauchy problem for them in the space $\mathcal{L}_2(R^n)$ were obtained by H. O. Kreiss ⁽²⁾ and are complicated and difficult to verify. On the other hand, if a system is Petrovskii correct, i.e., if all eigenvalues of its characteristic matrix lie in some fixed half-plane, then, as I. G. Petrovskii showed ⁽³⁾, the Cauchy problem for it is Hadamard well-posed. Thus, Hadamard well-posed Cauchy problems form a considerably broader class than uniformly well-posed ones.

In the present paper a definition is given of Hadamard well-posedness of the abstract Cauchy problem for an evolution equation in a Banach space, and necessary and sufficient conditions for Hadamard well-posedness are found.

In a Banach space E with norm $\|\cdot\|$, consider the differential equation

$$dx/dt = Ax(t) \quad (0 \leq t < \infty) \quad (1)$$

with a linear closed operator A , having an everywhere dense domain of definition $D(A)$ in E .

A solution of equation (1) on the half-axis $[0, \infty)$ is a function $x(t)$ with values in $D(A)$, strongly continuously differentiable on $[0, \infty)$ and satisfying equation (1) for all $t \in [0, \infty)$. By the Cauchy problem is meant the problem of finding a solution on $[0, \infty)$ satisfying the initial condition

$$x(0) = x_0 \in D(A). \quad (2)$$

Let in the space E there be given a linear set E_1 , contained in $D(A)$, on which another norm $\|\cdot\|_1$ and a seminorm $\|\cdot\|_2$ are defined.

Definition 1. The Cauchy problem (1), (2) is **Hadamard well-posed with type ω on the set E_1** if there exist constants C_1 and C_2 such that for any $x_0 \in E_1$ there exists a solution $x(t)$ satisfying the conditions

$$\|x(t)\| \leq C_1 e^{\omega t} \|x_0\|_1, \quad \|Ax(t)\| \leq C_2 e^{\omega t} \|x_0\|_2,$$

and this solution is unique.

If the Cauchy problem is Hadamard well-posed, then by a change of the unknown function it is reduced to the same problem with type $\omega = 0$. In what follows we shall consider Hadamard well-posed problems for which

$$\|x(t)\| \leq C_1 \|x_0\|_1, \quad \|Ax(t)\| \leq C_2 \|x_0\|_2 \quad (x_0 \in E_1). \quad (3)$$

We make the main assumption.

I. The positive half-axis does not belong to the point spectrum of the operator A , i.e., for $\lambda > 0$ there exists the operator $J_\lambda = (A - \lambda I)^{-1}$ (generally speaking, unbounded).

Lemma 1. *Under assumption I, for a given $x_0 \in D(A)$ there cannot exist more than one solution of problem (1), (2), bounded on the half-axis.*

Corollary. *Under assumption I, the uniqueness requirement in the definition of Hadamard well-posedness may be discarded.*

Definition 2. The maximal domain of Hadamard well-posedness is the linear set D of all elements x_0 for which there exists a solution of problem (1), (2), bounded together with its derivative on the half-axis $[0, \infty)$.

This definition is justified by the following consideration: by virtue of Lemma 1, on D the norm and seminorm are uniquely determined:

$$\|x_0\|_I = \sup_{0 \leq t < \infty} \|x(t)\|, \quad \|x_0\|_{II} = \sup_{0 \leq t < \infty} \|Ax(t)\|,$$

where $x(t)$ is the solution referred to in Definition 2. It is obvious that, in the norms $\|\cdot\|_I$ and $\|\cdot\|_{II}$, problem (1), (2) is Hadamard well-posed on D . Furthermore, if problem (1), (2) is Hadamard well-posed with type $\omega = 0$ on some set E_1 , then $E_1 \subset D$ and $\|x\|_I \leq C_1 \|x\|_1$ and $\|x\|_{II} \leq C_2 \|x\|_2$.

Under assumption I, let us study the properties of the operators J_ν ($\nu > 0$).

1°. The operator J_ν ($\nu > 0$) is defined on D , and the formula

$$J_\nu x_0 = - \int_0^\infty e^{-\nu \tau} x(\tau) d\tau \quad (x_0 \in D, \nu > 0), \quad (4)$$

holds, where $x(t)$ is the corresponding solution of problem (1), (2).

2°. The set D is invariant with respect to the operators J_ν ($\nu > 0$), and the solution $y(t)$ of equation (1) with the initial condition $y(0) = J_\nu x_0$ is determined by the formula

$$y(t) = - \int_0^\infty e^{-\nu\tau} x(t + \tau) d\tau. \quad (5)$$

Moreover,

$$\|y(t)\| \leq \frac{1}{\nu} \|x_0\|_I, \quad \|Ay(t)\| \leq \frac{1}{\nu} \|x_0\|_{II}.$$

3°. Hilbert's identity holds:

$$J_\nu J_\lambda x_0 = \frac{1}{\lambda - \nu} (J_\lambda - J_\nu) x_0 \quad (x_0 \in D; \lambda, \nu > 0),$$

as do the formulas for derivatives

$$J_\lambda^n x_0 = \frac{1}{(n-1)!} \frac{d^{n-1} J_\lambda x_0}{d\lambda^{n-1}} \quad (x_0 \in D, \lambda > 0, n \geq 1). \quad (6)$$

4°. For $x_0 \in D$, $\lambda > 0$, and $n \geq 0$, the inequalities

$$\|J_\lambda^n x_0\| \leq \lambda^{-n} \|x_0\|_I, \quad \|J_\lambda^n A x_0\| \leq \lambda^{-n} \|x_0\|_{II} \quad (7)$$

hold.

5°. The equalities (cf. (5))

$$\|x_0\|_I = \sup_{n \geq 0} \sup_{\lambda > 0} \|\lambda^n J_\lambda^n x_0\|, \quad (8)$$

$$\|x_0\|_{II} = \sup_{n \geq 0} \sup_{\lambda > 0} \|\lambda^n J_\lambda^n A x_0\| \quad (9)$$

hold.

From formula (5) it follows that the solution $z(t)$ of equation (1) with initial value $z(0) = \nu J_\nu x_0 + x_0$, for $x_0 \in D(A)$ and $\nu > 0$, is representable in the form

$$z(t) = - \int_0^\infty e^{-\nu\tau} A x(t + \tau) d\tau.$$

Hence, from (7) it follows

6°. The inequality holds

$$\|J_\lambda^n (\nu J_\nu x_0 + x_0)\| \leq \frac{C_2}{\nu \lambda^n} \|x_0\|_1, \quad x_0 \in D.$$

7°. For $x_0 \in D$ the relation

$$\lim_{\nu \rightarrow \infty} \|\lambda^n J_\lambda^n (\nu J_\nu A x_0 + A x_0)\| = 0 \quad (10)$$

holds uniformly with respect to $\lambda > 0$ and $n \geq 0$.

Let us introduce the set G , consisting of all elements $y \in E$ for which

$$\|y\|_G = \sup_{n \geq 0} \sup_{\lambda > 0} \|\lambda^n J_\lambda^n y\| < \infty. \quad (11)$$

From the closedness of the operators J_λ it follows that G , with respect to $\|\cdot\|_G$, is a Banach space. From 1°, 2°, 4°, and 5° it follows that $D \subset G$ and $\|x\|_1 = \|x\|_G$ for $x \in D$. For every $y \in G$ the function $J_\nu y$ can be analytically continued by means of the expansion

$$J_\nu y = \sum_{k=0}^{\infty} (\nu - \lambda)^k J_\lambda^{k+1} y \quad (\lambda > 0), \quad (12)$$

which converges in the disk $|\nu - \lambda| < \lambda$. Varying λ , we obtain an analytic continuation of $J_\nu x$ into the right half-plane.

The following assertion is essential:

Lemma 2. *The operator J_μ ($\mu > 0$) is a bounded operator on G in the norm (11), and, moreover, for it the relation*

$$\|J_\mu x\|_G \leq \frac{1}{\mu} \|x\|_G \quad (13)$$

holds.

For the proof, the quantity $\|\lambda^n J_\lambda^n (\mu J_\mu x)\|$ for $\mu < \lambda$ can be estimated with the aid of the expansion (12):

$$\|\lambda^n J_\lambda^n (\mu J_\mu x)\| \leq \frac{\mu}{\lambda} \sum_{k=0}^{\infty} \left(1 - \frac{\mu}{\lambda}\right)^k \|\lambda^{n+k+1} J_\lambda^{n+k+1} x\| \leq \|x\|_G.$$

For $\lambda < \nu$ the same estimate can be obtained by expanding the function $\lambda^n J_\lambda^n x$ in a series at the point $\lambda = \mu$. From these estimates the inequality (13) follows.

Let F now denote the collection of all elements of G for which

$$\|\mu J_\mu y + y\|_G \rightarrow 0 \quad \text{as } \mu \rightarrow \infty. \quad (14)$$

By virtue of the uniform boundedness of the operators μJ_μ , F is a closed subspace of G . From the commutativity of the operators J_ν and J_μ , and the boundedness of J_ν in G , it follows that $J_\nu F \subset F$. The operators J_ν bounded in F form a pseudo-resolvent, and since they vanish only at zero, they are the resolvent of some operator $\widehat{A}$. By virtue of (14), the domain of definition of this operator is dense in F . It is not difficult to verify that the operator $\widehat{A}$ is the restriction of the operator A to the space F , i.e., it is defined on all elements $x_0 \in F \cap D(A)$ for which $Ax_0 \in F$, and on them coincides with A .

By what has been said and by Lemma 2, for the operator $\hat{A}$ the conditions of the Hille-Yosida theorem are satisfied, and consequently it is the infinitesimal generator of a contraction semigroup $U(t)$ of operators in F (see (1,4)). For every $x_0 \in D(\hat{A})$ there exists a solution $x(t) = U(t)x_0$ of the equation

$$x' = \hat{A}x$$

in F , satisfying the condition $x(0) = x_0$. This solution will also be a solution of problem (1), (2), since the norm in F is stronger than the norm in E . Further:

$$\sup_{0 \leq t < \infty} \|x(t)\| \leq \sup_{0 \leq t < \infty} \|U(t)x_0\|_G = \|x_0\|_G < \infty,$$

$$\sup_{0 \leq t < \infty} \|Ax(t)\| \leq \sup_{0 \leq t < \infty} \|U(t)\hat{A}x_0\|_G = \|Ax_0\|_G < \infty,$$

i.e. $x_0 \in D$.

Thus, $D(\hat{A}) \subset D$. On the other hand, by properties 1⁰–7⁰, $D \subset D(\hat{A})$, whence $D = D(\hat{A})$.

From all that has been said the following main results follow.

Theorem 1. *Under assumption I the maximal domain of Hadamard correctness consists of all elements x_0 from $D(A)$ possessing the properties:*

- 1) $x_0 \in G$ and $\|\mu J_\mu x_0 + x_0\|_G \rightarrow 0$ as $\mu \rightarrow \infty$;
- 2) $Ax_0 \in G$ and $\|\mu J_\mu Ax_0 + Ax_0\|_G \rightarrow 0$ as $\mu \rightarrow \infty$.

Theorem 2. *Let I be satisfied. In order that the Cauchy problem (1), (2) be Hadamard correct on a set E_1 with type $\omega = 0$, the following conditions are necessary and sufficient:*

- 1) for each $x_0 \in E_1$ all operators J_λ ($\lambda > 0$) and all their possible products are defined;
- 2) the inequalities are valid ($x_0 \in E_1$):

$$\|J_\lambda^n x_0\| \leq C_1 \lambda^{-n} \|x_0\|, \quad \|J_\lambda^n Ax_0\| \leq C_2 \lambda^{-n} \|x_0\|_2;$$

- 3) uniformly with respect to $n \geq 0$ and $\lambda > 0$ the relations hold

$$\lim_{\mu \rightarrow \infty} \|\lambda^n J_\lambda^n (\mu J_\mu x_0 + x_0)\| = 0,$$

$$\lim_{\mu \rightarrow \infty} \|\lambda^n J_\lambda^n (\mu J_\mu Ax_0 + Ax_0)\| = 0.$$

Theorem 3. *Let I be satisfied and let the Cauchy problem be Hadamard correct with type $\omega = 0$ on the set E_1 . Then there exists a space F , algebraically and topologically embedded in the space E and containing the set E_1 , such that for the equation*

$$dx/dt = \hat{A}x,$$

where $\hat{A}$ is the restriction of A in the space F , the Cauchy problem is uniformly correct and has type $\omega = 0$.

Voronezh State University
named after the Lenin Komsomol

Latvian State University
named after P. Stuchka
Riga

Received
13 X 1969

CITED LITERATURE

- ¹ S. G. Krein, *Linear Differential Equations in Banach Space*, "Nauka," 1967.
- ² H. O. Kreiss, Sborn. per. Matematika, 7, No. 2, 39 (1963).
- ³ I. G. Petrovskii, Bull. Moscow State Univ., Ser. Math. and Mech., 1, 7, 1 (1938).
- ⁴ E. Hille, R. S. Phillips, *Functional Analysis and Semi-Groups*, IL, 1962.
- ⁵ W. Feller, Ann. Math., 58, 1, 166 (1953).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.