

ON QUASICONFORMAL HARMONIC MAPPINGS OF EUCLIDEAN SPACE

O. V. Titov

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.44758>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.54

MATHEMATICS

O. V. TITOV

ON QUASICONFORMAL HARMONIC MAPPINGS OF EUCLIDEAN SPACE

(Presented by Academician M. A. Lavrent'ev on 13 III 1970)

Let $f : \mathbf{R}^n \rightarrow \mathbf{R}^n$ be a C^∞ mapping of n -dimensional Euclidean space $\mathbf{R}^n$ into itself, given by n coordinate functions $f_1(x), \dots, f_n(x)$, where $x = (x_1, \dots, x_n)$. The mapping f is called quasiconformal if the inequality

$$\left(\sum_{i,j=1}^n \left| \frac{\partial f_i}{\partial x_j} \right|^2 \right)^{1/2} \leq C |J_f|^{1/n}, \quad (1)$$

holds everywhere, where C is some constant, and J_f is the Jacobian of the mapping f .

Combining two theorems of Liouville, one can prove the following assertion: any nonconstant quasiconformal mapping $f : \mathbf{R}^{2n} \rightarrow \mathbf{R}^{2n}$ which is at the same time holomorphic in the complex space $\mathbf{C}^n$, for $n > 1$, is a nondegenerate linear mapping.

In the present note we study a more general class of quasiconformal mappings $f : \mathbf{R}^n \rightarrow \mathbf{R}^n$ that are gradients of harmonic functions. Such mappings will be called harmonic.

Theorem. *A harmonic quasiconformal mapping $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ that does not reduce to a constant is nondegenerate linear.*

Lemma 1. *If a quasiconformal mapping $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ is given by analytic functions $f_i(x)$ ($i = 1, 2, 3$) and is different from a constant, then it is a homeomorphism. The Jacobian J_f therefore cannot assume values of different signs.*

Let S be the set of zeros of J_f . As an analytic set, S is the union of a finite number of manifolds of class C^∞ (see ⁽¹⁾). By virtue of inequality (1), the mapping f takes each connected component of S to a point. In the works ^(2,3) it is shown that the full preimage of a point under a quasiconformal mapping f , for any $\alpha > 0$, has zero α -dimensional Hausdorff measure. Consequently, each connected component of S consists of a single point, i.e. S is the union of a

finite number of points. The set B_f of branch points of the mapping f belongs to S and, consequently, is also finite.

We show that $B_f = \emptyset$. If B_f is nonempty, surround any point $x \in B_f$ by such a neighborhood U that it contains no other points of B_f . It is known (see ^(2,3)) that $f(U)$ is a certain neighborhood of the point $f(x)$. On the one hand, the fundamental group $\pi_1(f(U) \setminus f(x))$ is trivial; on the other hand, in $f(U)$ there certainly are points different from $f(x)$ whose preimage consists of more than one point, which indicates the nontriviality of $\pi_1(f(U) \setminus f(x))$. This contradiction proves that $B_f = \emptyset$.

Thus, the mapping f is locally homeomorphic, and then, by the theorem of V. A. Zorich ⁽⁴⁾, it is homeomorphic globally as well. Finally, it is well known that under these conditions $f(\mathbf{R}^3) = \mathbf{R}^3$ (see ⁽⁵⁾ or ⁽⁶⁾). The lemma is proved.

It remains to show that all the functions $f_i(x)$ ($i = 1, 2, 3$) in the formulation of the theorem are polynomials. Indeed, consider the mapping—

the mapping $h = \sigma f \sigma$, where $\sigma(x) = x|x|^{-2}$ is inversion. Since $f(\infty) = \infty$, we have $h(0) = 0$, and, by the multidimensional analogue of Mori's theorem ⁽⁷⁾, there exist constants C_1 and C_2 such that, in some neighborhood of zero,

$$|h(x)| > C_1|x|^{C_2}.$$

Returning to the mapping f , we obtain that for sufficiently large $|x|$

$$|f(x)| < \frac{1}{C_1}|x|^{C_2}.$$

This inequality is certainly also valid if in it $|f(x)|$ is replaced by $|f_i(x)|$ ($i = 1, 2, 3$). Since all f_i are harmonic functions, it follows from the inequalities obtained that they are polynomials.

It also follows easily from inequality (1) that all degrees of the polynomials f_i are the same. We shall denote the common value of these degrees by m . We may write

$$f_i = p_i + q_i \quad (i = 1, 2, 3),$$

where p_i are homogeneous polynomials of degree m , while the degrees of q_i are strictly less than m . It is obvious that the mapping $p : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ defined by the coordinate functions p_i ($i = 1, 2, 3$) is harmonic.

Lemma 2. Let $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ be a quasiconformal homeomorphism, with $f(0) = 0$. Denote by $m(R)$ and $M(R)$, respectively, the minimum and maximum of $|f(x)|$ on the sphere $|x| = R$. Then there exists a constant C , depending only on f , such that

$$M(R) \leq Cm(R). \quad (2)$$

The proof follows easily from the estimate given by Gehring in ⁽⁸⁾ for the modulus of a ring.

Lemma 3. Let $f : \mathbf{R}^3 \rightarrow \mathbf{R}^3$ be a homeomorphism given by polynomials, and let $f(0) = 0$. Take some ray l issuing from the origin 0, and a point $x \in l$ such that $|x| = r$. If $s_l(r)$ is the length of the image of the segment $[0, x]$ under the mapping f , then there exists a constant $C(l)$, depending possibly on l , such that for every $r > 1$

$$s_l(r) \leq C(l)|f(x)|. \quad (3)$$

The proof is elementary.

Without loss of generality we may assume that for the original mapping f , $J_f \geq 0$ and $f(0) = 0$. Let $J_f|_l$ be the restriction of the Jacobian J_f to some ray l issuing from the origin. Then, in the notation of Lemma 3, $J_f|_l$ is a polynomial in r of degree k_l .

The number k_l does not depend on l . Indeed, for rays l_j ($j = 1, 2$) such that $k_{l_1} < k_{l_2}$, take points x_j lying on l_j at distance $r > 1$ from the origin ($j = 1, 2$). Using (2) and (3), we obtain

$$s_{l_1}(r) \geq m(r) \geq CM(r) \geq C|f(x_2)| \geq \frac{C}{C(l_2)} s_{l_2}(r).$$

But

$$s_{l_j}(r) = \int_0^r \mu_{l_j}(t) dt \quad (j = 1, 2),$$

where $\mu_l(t)$ are the coefficients of local stretching in the corresponding directions. From the geometric definition of quasiconformality there follows the existence of constants C_1 and C_2 such that

$$C_1(J_f|_{l_j})^{1/3} \leq \mu_{l_j} \leq C_2(J_f|_{l_j})^{1/3} \quad (j = 1, 2).$$

Hence $k_{l_1} \geq k_{l_2}$, which contradicts the inequality $k_{l_1} < k_{l_2}$.

Denote by k the common value of the k_l , and let us show that $k = 3(m-1)$. On the sphere $|x| = 1$ there exists a point x such that $p(x) \neq 0$. It is easy to find constants $M_1(x)$ and $M_2(x)$ such that, for $r > 1$,

$$M_1(x)r^m \leq |f(rx)| \leq M_2(x)r^m. \quad (4)$$

If l is the ray determined by the vector x , then, using (3) and (4), we obtain

$$\int_0^r (J_f|_l)^{1/3} dt \geq \frac{1}{C_2} |f(rx)| \geq \frac{M_1(x)}{C_2} r^m = \frac{mM_1(x)}{C_2} \int_0^r [t^{3(m-1)}]^{1/3} dt.$$

It follows that $k \geq 3(m-1)$. Similarly one proves that $k \leq 3(m-1)$.

Let us show that everywhere on the unit sphere the Jacobian J_p of the mapping p , defined above by the homogeneous components of highest degree of the polynomials f_i ($i = 1, 2, 3$), does not vanish. Indeed, if at some point x , $|x| = 1$, one had $J_p(x) = 0$, then

$$\lim_{r \rightarrow \infty} \frac{J_f(rx)}{r^{3(m-1)}} = \lim_{r \rightarrow \infty} \det \left(\frac{\partial p_i / \partial x_j|_{rx} + \partial q_i / \partial x_j|_{rx}}{r^{m-1}} \right) = \lim_{r \rightarrow \infty} \det \left(\frac{\partial p_i}{\partial x_j} \Big|_x + \frac{\partial q_i / \partial x_j|_{rx}}{r^{m-1}} \right) = 0,$$

and this contradicts the equality $k = 3(m-1)$.

In view of the compactness of the unit sphere, there exists a constant C such that, for $|x| = 1$,

$$\left(\sum_{i,j=1}^3 \left| \frac{\partial p_i}{\partial x_j} \right|^2 \right)^{1/2} \leq C |J_p|^{1/3}. \quad (5)$$

Since on the right and on the left in (5) there are homogeneous functions of degree $m-1$, this inequality is valid everywhere in $\mathbf{R}^3$, i.e., p is quasiconformal.

By Lemma 1 the mapping p is a homeomorphism, but, according to a result of Levy [9], the Jacobian of a harmonic homeomorphism in $\mathbf{R}^3$ does not vanish at 0. This is possible only when the degree of homogeneity of J_p is zero, i.e., $m = 1$. The theorem is proved.

Remark. All the results presented, except Levy's theorem [9], are also valid for the n -dimensional ($n \geq 3$) case. The question of the nonvanishing of the Jacobian of a harmonic homeomorphism in $\mathbf{R}^n$ ($n > 3$) remains open.

The author expresses gratitude to B. V. Shabat for posing the question and for great assistance in the work.

Moscow State University
named after M. V. Lomonosov

Received
7 III 1970

References

1. R. Thom, in: Collection, *Singularities of differentiable mappings*, 1968.
2. Yu. G. Reshetnyak, *Siberian Mathematical Journal*, 9, No. 2 (1968).
3. Yu. G. Reshetnyak, *ibid.*, 8, No. 3 (1967).
4. V. A. Zorich, *Mathematical Collection*, 74 (116), No. 3 (1967).
5. B. V. Shabat, *Dokl. Akad. Nauk SSSR*, 132, No. 5 (1960).
6. Ch. Loewner, *J. Math. Mech.*, 8 (1959).
7. P. Caraman, *Homeomorfisme cvasiconforme n-dimensionale*, București, 1968.
8. F. W. Gehring, *Trans. Am. Math. Soc.*, 101, No. 3 (1961).
9. H. Levy, *Ann. Math.*, 88, No. 3 (1968).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.