

SPACES DEFINED BY THE VARIATION OF LOCAL APPROXIMATIONS

Yu. A. Brudnyi

1970

SovietRxiv

View the original and related papers at <https://sovietsxiv.org/items/ru-197001.61972>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 517.51

MATHEMATICS

Yu. A. BRUDNYI

SPACES DEFINED BY THE VARIATION OF LOCAL APPROXIMATIONS

(Presented by Academician V. I. Smirnov on 4 VI 1969)

1. A family of spaces defined by the variation of local approximations is obtained by means of a very simple construction. Nevertheless, among the members of this family there are a number of spaces that are constantly used in analysis (Lipschitz spaces, functions of bounded variation, Sobolev spaces, etc.). Thus a somewhat unexpected connection between these spaces is established; in particular, from the theorems given below there follows a number of new properties of these spaces. We note that two representatives of the family of spaces under consideration were studied earlier by F. Riesz (Riesz' s lemma, see ⁽¹⁾, p. 85) and by F. John—L. Nirenberg ⁽²⁾.
2. We shall need several definitions. Below Q_0 denotes a fixed n -cube, and $Q \subset Q_0$ a cube parallel to it. Let X be a Banach space whose elements are functions on the cube $I(Q)$, and suppose that the norm in X has the following property: if $|I_1(Q)| \leq |I_2(Q)|$, $Q \subset Q_0$, and $I_2 \in X$, then $I_1 \in X$ and $\|I_1\| \leq \|I_2\|$. Let P be the projector $L_p(Q_0) \rightarrow P_{k-1}$, where P_{k-1} is the space of polynomials of degree $k-1$ (consisting only of zero when $k=0$), and let P_Q be the "transplant" of P to $L_p(Q)$ by means of a homothety carrying Q_0 onto Q . Finally, let

$$I_f^k(Q) = \left\{ \frac{1}{m(Q)} \int_Q |f - P_Q f|^p dx \right\}^{1/p}. \quad (1)$$

Definition. The space $\mathcal{L}_p^k(X)$ is the linear set of functions $f \in L_p(Q_0)$ for which $I_f^k(Q) \in X$. As the norm of f in this space we take the quantity

$$\|f\|_{L_p(Q_0)} + \|I_f^k\|_X.$$

3. We choose as X the space of functions $I(Q)$ having bounded variation in the following sense:

$$V(I; \Omega) = \sup \left\{ \sum m(Q_s) \left| \frac{I(Q_s)}{\varphi(a_s)} \right|^q \right\}^{1/q} < +\infty \quad (2)$$

for every open $\Omega \subset Q_0$. Here the supremum is taken over all families $\{Q_s\}$ of pairwise nonoverlapping cubes in Ω ; a_s is the side length of Q_s , and φ is a majorant, i.e. a positive nondecreasing function on $(0, +\infty)$ satisfying the condition

$$\sup \frac{\varphi(2\tau)}{\varphi(\tau)} < +\infty.$$

We denote this space by V_q^φ ; its subspace consisting of those I for which (2) is absolutely continuous will be denoted by v_q^φ .

For the formulation of the first theorem, set

$$m_Q(f; \tau) = \text{mes}\{x \in Q \mid |(f - P_Q f)(x)| > \tau\} \quad (3)$$

and by $f_Q^*(\tau)$, $0 < \tau \leq m(Q)$, the right-continuous function inverse to (3).

Theorem 1. If $f \in \mathcal{L}_p^k(V_q^\varphi)$, $1 \leq p, q \leq \infty$, then for every $Q \subset Q_0$ with side a

$$f_Q^*(\tau^n) \leq cV(I_f^k; Q) \int_\tau^a \frac{\varphi(\tau)}{\tau^{1+n/q}} d\tau. \quad (4)$$

From Theorem 1, for $k = 1$ and $\varphi = 1$, we obtain the known results of John–Nirenberg ⁽²⁾, and for $k = 1$ and $q = \infty$, the result of S. Spanne ⁽³⁾.

Corollary 1. If, for some $r \geq q$, the function

$$\bar{\varphi}(\tau) = \tau^\lambda \int_0^\tau \frac{\varphi(u)}{u^{1+a}} du, \quad a = n(q^{-1} - r^{-1}),$$

is defined, then there is a continuous embedding

$$\mathcal{L}_p^k(V_q^\varphi) \subset \mathcal{L}_r^k(V_q^{\bar{\varphi}}).$$

In particular, if $p \leq r$ and φ is a quasi-degree majorant, i.e. $\bar{\varphi} \leq \varphi$, then $\mathcal{L}_p^k(V_q^\varphi)$ is isomorphic to $\mathcal{L}_r^k(V_q^\varphi)$.

Corollary 2. All spaces $\mathcal{L}_p^k(V_q^\varphi)$, $1 \leq p < q$, are isomorphic.

For the formulation of the second theorem, assume that $\varphi(\tau) = \tau^s \psi(\tau)$, where $s \geq 0$ is an integer smaller than k , and ψ is a modulus of continuity. Put

$$\bar{\psi}(\tau) = \int_0^\tau \frac{\psi(u)}{u} du + \tau \int_\tau^1 \frac{\psi(u)}{u^2} du, \quad (5)$$

where, for $s = k - 1$, the second term is omitted.

Theorem 2. If $f \in \mathcal{L}_p^k(V_q^\varphi)$, $\varphi(\tau) = \tau^s \psi(\tau)$, and the function (5) exists, then for every $\varepsilon > 0$

$$f = g + h,$$

where h has support whose measure is less than ε , and g belongs to $C^{s, \bar{\psi}}$, with

$$\sup_{|\alpha|=s} \sup_{x \neq y} \frac{|D^\alpha(g; x) - D^\alpha(g; y)|}{\bar{\psi}(|x - y|)} = O(\varepsilon^{-1/q}) V(I_f^k; Q_0).$$

Remark. One can also give an estimate for higher-order differences of the function $D^\alpha g$ on the set $Q_0 \setminus \text{supp } h$.

4. We indicate the connection between $\mathcal{L}_p^k(V_q^\varphi)$ and known function spaces.

A. Sobolev spaces.

Theorem 3. The space $\mathcal{L}_p^k(V_q^\varphi)$, for $\varphi(\tau) = \tau^k$, $1 < q \leq p$, and $k > n(q^{-1} - p^{-1})$, is isomorphic to the space $W_q^k \cap L_p$.

Remark. For $k = n(q^{-1} - p^{-1})$ only the proper embedding into $W_q^k \cap L_p$ holds.

For $n = 1$, $p = \infty$, and $k = 1$, we obtain from Theorem 3 the known lemma of F. Riesz (see ⁽¹⁾, p. 85). From Theorem 1 we now obtain the embedding theorem of S. L. Sobolev ⁽⁴⁾ with a certain refinement in the limiting case (cf. ⁽⁵⁾), and from Theorem 2, a certain new property of functions from Sobolev spaces, namely that, after modification on a set of small measure, they coincide with functions from $C^{k-1,1}$.

B. Lipschitz spaces. In the works ⁽⁶⁻⁹⁾ it was shown that if $\varphi(\tau) = \tau^{s+\alpha}$, $0 \leq s < k - 1$, $0 < \alpha < 1$, or $\varphi(\tau) = \tau^{s+\alpha}$, $s = k - 1$, $\alpha = 1$, then $\mathcal{L}_p^k(V_\infty^\varphi)$ is isomorphic to the space $C^{s, \alpha}$. It also follows from the results of ⁽⁹⁾ that if $0 < s \leq k - 1$ and $\alpha = 0$, then $\mathcal{L}_p^k(V_\infty^\varphi)$ is isomorphic to $C^{s-1, 1-0}$, where the space $C^{l, 1-0}$ is defined by the condition that the second-

differences of step h of the higher derivatives are majorized by the quantity $O(|h|)$. For $q < \infty$ we have the following result.

Theorem 4. If $\varphi(\tau) = \tau^\alpha$, $0 < \alpha < k$, then for $p \leq q$ the continuous embeddings

$$L_p \cap B_q^\alpha \subset \mathcal{L}_p^k(V_q^\varphi) \subset H_p^\alpha,$$

hold, and for $q \leq p$ the embeddings

$$B_p^\alpha \subset \mathcal{L}_p^k(V_q^\varphi) \subset H_q^\alpha \cap L_p.$$

Here H_p^α , B_p^α are the well-known spaces of S. M. Nikol'skii and O. V. Besov (see the survey¹⁰).

B. Functions of bounded variation. For $\varphi(\tau) = \tau^{1/q}$ and $n = 1$, the space $\mathcal{L}_\infty(V^\varphi)$ is isomorphic to the space of functions of bounded q -variation in the sense of Wiener–L. Young. Therefore also for $n > 1$ it is natural to call the functions of the space $\mathcal{L}_\infty(V_q^\varphi)$, where $\varphi(\tau) = \tau^{1/q}$, functions of bounded q -variation. The space $\mathcal{L}_1(V_1^\varphi)$ for $\varphi(\delta) = \tau$ is isomorphic to the space of functions having bounded variation in the sense of Tonelli. More generally, the following holds.

Theorem 5. The space $\mathcal{L}_1^k(V_1^\varphi)$ for $\varphi(\tau) = \tau^k$ is isomorphic to the space BV^k of functions in $L_1(Q_0)$ whose generalized k -derivatives are Borel measures.

Remark. At the same time, the space $\mathcal{L}_1^k(v_1^\varphi)$, $\varphi(\tau) = \tau^k$, is isomorphic to the space W_1^k .

5. One can choose X in $\mathcal{L}_p^k(X)$ also in such a way as to obtain any Lipschitz space in the sense of Calderon (see¹¹). In particular, one can obtain the spaces H_p^α of S. M. Nikol'skii or $B_{p,q}^\alpha$ of O. V. Besov.
6. In conclusion we state an interpolation theorem for operators acting from the scale L_p into the space $\mathcal{L}_p^k(X)$.

Let X_0, X_1 be two spaces of functions of the cube $I(Q)$, described in item 2. Denote by X_s , $0 < s < 1$, the space of those functions $I(Q)$ for each of which there are a number λ and functions $I_i \in X_i$, $i = 0, 1$, such that

$$|I(Q)| \leq \lambda |I_0(Q)|^{1-s} |I_1(Q)|^s, \quad Q \subset Q_0. \quad (6)$$

We take the lower bound of λ in (6), under $\|I_i\|_{X_i} \leq 1$, as the norm in X_s .

Theorem 6. Let a linear operator T act continuously from L_{p_i} to $\mathcal{L}_{p_i}^k(X_i)$, and let its norm be a_i , $i = 0, 1$. Then T acts continuously from L_p to $\mathcal{L}_p^k(X_s)$, where $1/p = (1-s)/p_0 + s/p_1$, and its norm does not exceed $a_0^{1-s} a_1^s$.

Dnepropetrovsk Chemical-Technological
Institute

Received
23 V 1969

CITED LITERATURE

- ¹ F. Riesz, B. Sz.-Nagy, *Lectures on Functional Analysis*, IL, 1964.
- ² F. John, Z. Nirenberg, *Comm. Pure Appl. Math.*, **14**, 415 (1961).
- ³ S. Spanne, *Ann. Scuola Norm. Super. Pisa*, **19**, No. 4, 593 (1965).
- ⁴ S. L. Sobolev, *Some Applications of Functional Analysis in Mathematical Physics*, L., 1950.
- ⁵ V. I. Yudovich, *DAN*, **138**, No. 3, 805 (1961).
- ⁶ S. Campanato, *Ann. Scuola Norm. Super. Pisa*, **17**, 175 (1963).
- ⁷ N. Meyers, *Proc. Am. Math. Soc.*, **15**, 717 (1964).
- ⁸ S. Campanato, *Ann. Scuola Norm. Super. Pisa*, **18**, No. 1, 137 (1964).
- ⁹ Yu. A. Brudnyi, *Studies in the Theory of Local Best Approximation*, dissertation, Dnepropetrovsk, 1965.
- ¹⁰ S. M. Nikol'skii, *UMN*, **16**, No. 5 (101), 63 (1961).
- ¹¹ A. P. Calderon, *Studia Math.*, **24**, 113 (1964).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.