

THE GAMMA-CORRELATION PROCESS AND ITS PROPERTIES

I. O. Sarmanov

1970

SovietRxiv

View the original and related papers at <https://sovietrxiv.org/items/ru-197001.65576>

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.

Abstract

Full Text

UDC 519.272.129

MATHEMATICS

I. O. SARMANOV

THE GAMMA-CORRELATION PROCESS AND ITS PROPERTIES

(Presented by Academician Yu. V. Linnik on 17 VII 1969)

1. Theorem. The function $p(t_1, x; t_2, y)$, defined for $0 < t_1 < t_2$, $x, y \geq 0$ by the bilinear expansion

$$p(t_1, x; t_2, y) = p(t_1, x)p(t_2, y) \sum_{k=0}^{\infty} a_k(t_1, t_2) L_k^{\alpha(t_1)}\left(\frac{x}{\beta(t_1)}\right) L_k^{\alpha(t_2)}\left(\frac{y}{\beta(t_2)}\right), \quad (1)$$

where

$$p(t, x) = x^{\alpha(t)} e^{-x/\beta(t)} / [\beta(t)]^{\alpha(t)+1} \Gamma(\alpha(t) + 1); \quad (2)$$

$$a_k(t_1, t_2) =$$

$$= [\omega(t_1)/\omega(t_2)]^k \left[\frac{\Gamma(\alpha(t_2) + 1) \Gamma(\alpha(t_1) + k + 1)}{\Gamma(\alpha(t_1) + 1) \Gamma(\alpha(t_2) + k + 1)} \right]^{1/2}; \quad (3)$$

$\beta(t) > 0$; $\alpha(t) > -1$ is a monotonically increasing continuous function, $\omega(t)$ is a continuous strictly increasing function, and $L_k^{\alpha}(x/\beta)$ are Laguerre polynomials, is a two-dimensional density defining a Markov process $\xi(t) \geq 0$ for $t > 0$.

Proof. By virtue of the orthogonality and normality of the Laguerre polynomials

$$L_k^{\alpha(t)}\left(\frac{x}{\beta(t)}\right) = \left[\frac{\Gamma(\alpha(t) + 1) \Gamma(\alpha(t) + k + 1)}{k!} \right]^{1/2} \sum_{r=0}^k \frac{(-1)^r \binom{k}{r}}{\Gamma(\alpha(t) + r + 1)} \left(\frac{x}{\beta(t)}\right)^r \quad (4)$$

with weight (2) on the half-axis $0 \leq x < \infty$, the function (1) satisfies the following Markov equation for the two-dimensional density:

$$p(t_1, x; t_2, y) = \int_0^\infty \frac{p(t_1, x; t, z) p(t, z; t_2, y)}{p(t, z)} dz, \quad (5)$$

where

$$p(t, z) = \int_0^\infty p(t_1, x; t, z) dx$$

is the marginal density (2) of the gamma distribution, $0 < t_1 < t < t_2$.

The series (1) for $t_1 < t_2$ converges in the mean, since the squares of the coefficients of the series

$$a_k^2(t_1, t_2) = [\omega(t_1)/\omega(t_2)]^{2k}$$

are less than the terms of a geometric progression with ratio $\omega^2(t_1)/\omega^2(t_2) < 1$.

Consequently, to prove the validity of the assertion of the theorem, it remains to show the nonnegativity of the sum of the series (1) for $t_1 < t_2$ and $x, y \geq 0$.

Noting that the Fourier transform of polynomial (4) with weight (2) has the form¹

$$[\Gamma(\alpha + k + 1)/\Gamma(\alpha + 1)k!]^{1/2} \frac{(-i\tau\beta)^k}{(1 - i\tau\beta)^{\alpha+k+1}},$$

we find the two-dimensional Fourier transform of function (1)

$$\varphi(\tau_1, \tau_2) = \frac{[1 - i\tau_2\beta(t_2)]^{\alpha(t_1) - \alpha(t_2)}}{[1 - i\tau_1\beta(t_1) - i\tau_2\beta(t_2) - (1 - \omega(t_1)/\omega(t_2))\tau_1\tau_2\beta(t_1)\beta(t_2)]^{\alpha(t_1)+1}}. \quad (6)$$

Expression (6) is a two-dimensional characteristic function, since the numerator is a one-dimensional characteristic function of the gamma distribution with non-negative parameter $\alpha(t_2) - \alpha(t_1)$, and the second factor is the two-dimensional characteristic function of the symmetrized gamma correlation studied in [2]. Consequently, (1) is a two-dimensional density. The theorem is proved.

2. In [3] it is noted that the correlation function of a continuous Markov process for $t_1 < t_2$ must have the form $R(t_1, t_2) = \psi(t_1)/\psi(t_2)$, where $\psi(t)$ is a continuous strictly increasing function.

¹The visible page contains the marker "(1)" after "has the form" ; it is preserved here as a footnote marker.

In our case $\psi(t) = \omega(t)[\alpha(t) + 1]^{1/2}$, and

$$R(t_1, t_2) = a_1(t_1, t_2) = \omega(t_1)[\alpha(t_1) + 1]^{1/2} / \omega(t_2)[\alpha(t_2) + 1]^{1/2}, \quad (7)$$

since, in general, $a_k(t_1, t_2)$ is the correlation coefficient between $L_k^{\alpha(t_1)}(\xi(t_1)/\beta(t_1))$ and $L_k^{\alpha(t_2)}(\xi(t_2)/\beta(t_2))$; hence $a_1(t_1, t_2)$ is the correlation coefficient between $\xi(t_1)$ and $\xi(t_2)$.

Definition. The Markov process $\xi(t)$ specified by density (1) will be called a **gamma-correlation process** or a **gamma process**.

3. If $\alpha(t) = \alpha$, $\beta(t) = \beta$ do not depend on t , and $\omega(t) = e^{\lambda t}$, where λ is a positive constant, then $a_k(t_1, t_2) = \exp\{-\lambda(t_2 - t_1)k\}$, and (1) defines the stationary Markov process studied in [4] (for $\beta = 1$).

If $\alpha = -1/2$ (does not depend on t), $\beta(t) = \omega(t) = t$, then $a_k(t_1, t_2) = (t_1/t_2)^k$, and we obtain a gamma process $\xi(t)$ with density

$$\begin{aligned} & \frac{(x/t_1)^{-1/2} e^{-x/t_1}}{\Gamma(1/2)t_1} \frac{(y/t_2)^{-1/2} e^{-y/t_2}}{\Gamma(1/2)t_2} \sum_{k=0}^{\infty} \left(\frac{t_1}{t_2}\right)^k L_k^{-1/2}\left(\frac{x}{t_1}\right) L_k^{-1/2}\left(\frac{y}{t_2}\right) = \\ & = \frac{\text{ch}[2(xy)^{1/2}/(t_2 - t_1)]}{\pi[t_1(t_2 - t_1)xy]^{1/2}} \exp\left\{-\frac{x/t_1 + y/t_2}{1 - t_1/t_2}\right\}. \end{aligned} \quad (8)$$

It is easy to verify that, if one considers the Wiener process $\eta(t)$ with parameters

$$\mathbf{M}\eta(t) = 0, \quad \mathbf{D}\eta(t) = t, \quad R(t_1, t_2) = (t_1/t_2)^{1/2},$$

then (8) is the two-dimensional density of the process $\xi(t) = \eta^2(t)/2$.

Definition. If, for the process $\xi(t)$ with density (1), the correlation function $R(t_1, t_2) = t_1/t_2$ is equal to the square of the correlation function of the Wiener process, then the process $\xi(t)$ will be called a **gamma process of Wiener type**.

4. Let us consider a more general case of a gamma process of Wiener type. Put $\alpha(t) + 1 = t$, $\omega(t) = t^{1/2}$, $\beta(t) = 1$; then the process will have the basic parameters

$$\mathbf{M}\xi(t) = t, \quad \mathbf{D}\xi(t) = t, \quad R(t_1, t_2) = t_1/t_2. \quad (9)$$

The marginal density has the form

$$p(t, x) = x^{t-1} e^{-x} / \Gamma(t), \quad t > 0, \quad x \geq 0.$$

The two-dimensional density is written in the form of a series

$$p(t_1, x; t_2, y) = \frac{x^{t_1-1} e^{-x}}{\Gamma(t_1)} \frac{y^{t_2-1} e^{-y}}{\Gamma(t_2)} \sum_{k=0}^{\infty} \left(\frac{t_1}{t_2} \right)^{k/2} \left[\frac{\Gamma(t_2) \Gamma(t_1 + k)}{\Gamma(t_1) \Gamma(t_2 + k)} \right]^{1/2} L_k^{t_1-1}(x) L_k^{t_2-1}(y), \quad (10)$$

where $0 < t_1 < t_2$.

Let us note that the trajectories of the process defined by density (10) may, as in the Wiener case, be regarded as issuing from the origin; they all lie in the first quadrant of the plane XOT , where time t is plotted along the abscissa axis. The mean value $M\xi(t) = t$ grows without bound as t increases, whereas in the Wiener case $M\eta(t) = 0$.

In conclusion, we write down some other characteristics of the process (10). The characteristic function of the two-dimensional distribution has the form

$$\varphi(\tau_1, \tau_2) = \frac{1}{(1 - i\tau_2)^{t_2-t_1}} \frac{1}{[1 - i\tau_1 - i\tau_2 - (1 - (t_1/t_2)^{1/2}) \tau_1 \tau_2]^{t_1}}, \quad 0 < t_1 < t_2.$$

The initial moments of the marginal distribution are expressed by the formula

$$M\xi^k(t) = t(t+1) \cdots (t+k-1), \quad k = 1, 2, \dots$$

The conditional moments $m_k(t_1, t_2, x)$ of the quantity $\xi(t_2)$ under the condition $\xi(t_1) = x$ are expressed as follows:

$$m_k(t_1, t_2, x) = \Gamma(t_2 + k) \sum_{\mu=0}^k \frac{\binom{k}{\mu} [x(t_1/t_2)^{1/2}]^\mu}{\Gamma(t_2 + \mu)} \sum_{\nu=0}^{k-\mu} \binom{k-\mu}{\nu} [(t_1/t_2)^{1/2}]^\nu \times \\ \times \frac{\Gamma(t_2 + \mu) \Gamma(t_1 + \mu + \nu)}{\Gamma(t_1 + \mu) \Gamma(t_2 + \mu + \nu)}. \quad (11)$$

With the aid of (11) one can find the conditional variance

$$D(\xi(t_2) | \xi(t_1) = x) = t_2 - t_1 + t_1 [1 - (t_1/t_2)^{1/2}]^2 + 2(t_1/t_2)^{1/2} [1 - (t_1/t_2)^{1/2}] x. \quad (12)$$

Remark. We have put $\beta = 1$ only to shorten the notation; replacing x and y respectively by $x/\beta(t_1)$ and $y/\beta(t_2)$ will not change the correlation function

and will make it possible, without difficulty, to pass to a more general process of Wiener type.

Institute of Water Problems
Academy of Sciences of the USSR
Moscow

Received
2 VII 1969

REFERENCES

1. A. Kratzer, W. Franz, *Transcendental Functions*, IL, 1963.
2. O. V. Sarmanov, DAN, 132, No. 2, 299 (1960).
3. S. N. Bernstein, *Collected Works*, 4, Moscow, 1964.
4. O. V. Sarmanov, *Proceedings of the V. A. Steklov Mathematical Institute, Academy of Sciences of the USSR*, 60, 238 (1961).

Note: Figure translations are in progress. See original paper for figures.

Source: Math-Net.Ru and CyberLeninka. Machine translation. Verify with the original.